

A statewide probability survey of  
Vermont lakes - an adaptation of the  
National Lakes Survey

Neil Kamman

VT Agency of Natural Resources  
Water Quality Division

New England Association of Environmental Biologists  
Bartlett, NH  
March 26, 2008

# Some acknowledgements

- Funding
  - USEPA C.W.A. 106
  - VTANR
- Individuals
  - Hilary Snook EPA R1
  - McKalyn Garrity, Eric Howe, Sarah Wheeler

# Today's discussion

- Purpose and design
- Indicators and field approach
- Progress with data analysis
- The sample draw as a palate for analyzing existing data
- The littoral habitat assessment – analysis of new data
- Outreach

# Purpose

- National Lakes Survey
  - Component of USEPA's probability survey initiative
  - Lakes are one of several resources assessed over 5-year repeating blocks
- New England Lakes and Ponds Project
  - a regional do-over of EMAP NE Lakes leveraging the national draw
- VT Lakes Survey
  - provides a probability survey of our lakes that also contributes to NELP and NLA.
- VT's interest is to capture a second probability survey of our lakes\*, and to evaluate NLA and NELP techniques for adoption as standard VT protocols

\*) The first probability survey was the '98-'99 REMAP Mercury Project

# Design

- 2007: Sampled 33 lakes
  - 10 NLA core lakes
  - 20 additional state lakes from the overdraw.
  - Gives its own probability survey w/ ~85% CL.
  - 3 reference lakes (2 from overdraw, one selected).
- 2008: Sample 25 lakes
  - Begin at # 25 on the sample roster
  - Completes survey to a 50-lake probability survey w/ 95% CL.
  - All VT lakes contributing to New England Lakes and Ponds Regional Study.
- Lake selection carried out by EPA Corvallis on the NHD lakes sample frame as a stratified random draw of 10+ acre waters.

# Variance components capture by the VT design

- Within lakes
  - QA on 6 lakes, including replication, spikes, and blanks.
- Within years
  - One lake resampled in 2007, another in 2008.
- Among years
  - Five lakes sampled in 2007 are resampled in 2008.

2007 – 2008, 50 lakes, ~95% CL, +3 reference lakes

2007, 33 LAKES, ~85% CL

Completed?	PANEL	NAME
<input checked="" type="checkbox"/>	Panel_1	Marshfield Pond
<input checked="" type="checkbox"/>	Panel_1	Island Pond, Resample
<input checked="" type="checkbox"/>	Panel_1	Miles Pond
<input checked="" type="checkbox"/>	Panel_1	Lake Champlain
<input checked="" type="checkbox"/>	Panel_1	Caspian Lake
<input checked="" type="checkbox"/>	Panel_1	Derby, Lake
<input checked="" type="checkbox"/>	Panel_1	Silver Lake
<input checked="" type="checkbox"/>	Panel_1	Parker, Lake
<input checked="" type="checkbox"/>	Panel_1	Island Pond
<input checked="" type="checkbox"/>	Panel_1	Beebe Pond
<input checked="" type="checkbox"/>	OverSamp	Martins Pond
<input checked="" type="checkbox"/>	OverSamp	Echo Lake
<input checked="" type="checkbox"/>	OverSamp	Mud Pond
<input checked="" type="checkbox"/>	OverSamp	Curtis Pond
<input checked="" type="checkbox"/>	OverSamp	Eden, Lake
<input checked="" type="checkbox"/>	OverSamp	Cedar Lake
<input checked="" type="checkbox"/>	OverSamp	Harriman Reservoir
<input checked="" type="checkbox"/>	OverSamp	Lowell Lake
<input checked="" type="checkbox"/>	OverSamp	Spring Lake
<input checked="" type="checkbox"/>	OverSamp	Seymour Lake
<input checked="" type="checkbox"/>	OverSamp	Bomoseen, Lake
<input checked="" type="checkbox"/>	OverSamp	Little Averill Pond
<input checked="" type="checkbox"/>	OverSamp	Clyde Pond
<input checked="" type="checkbox"/>	OverSamp	Coles Pond
<input checked="" type="checkbox"/>	OverSamp	Chandler Pond
<input checked="" type="checkbox"/>	OverSamp	Silver Lake
<input checked="" type="checkbox"/>	OverSamp	Bliss Pond
<input checked="" type="checkbox"/>	OverSamp	Hardwick Lake
<input checked="" type="checkbox"/>	OverSamp	Little Rock Pond
<input checked="" type="checkbox"/>	OverSamp	Jobs Pond
<input checked="" type="checkbox"/>	OverSamp	Richville
<input checked="" type="checkbox"/>	OverSamp	Peacham Pond
<input checked="" type="checkbox"/>	NonTarget	Maidstone Lake


Reference lakes


2008, +25 LAKES

PANEL	NAME
OverSamp	Peacham Pond
OverSamp	Clyde Pond
OverSamp	Little Averill Pond
OverSamp	Bomoseen, Lake
OverSamp	Seymour Lake
OverSamp	Mill Pond
OverSamp	Lily Pond
OverSamp	Elmore, Lake
OverSamp	Indian Bk. Res.
OverSamp	Crystal Lake
OverSamp	Reservoir Pond
OverSamp	Round Pond
OverSamp	Berlin Pond
OverSamp	Norton Pond
OverSamp	Old Marsh Pond
OverSamp	Salem-Little
OverSamp	Sabin Pond
OverSamp	Neal Pond
OverSamp	North Springfield Reservoir
OverSamp	Branch Pond
OverSamp	Iroquois, Lake
OverSamp	Carmi, Lake
OverSamp	Doughty Pond
OverSamp	East Long Pond
OverSamp	Stiles Pond

Replication lake

Resample lakes










# Indicators

- 10 NLA Core lakes (+ reference lakes):
  - All NLA indicators
  - Laboratory splits with DEC lab for all WQ parameters and sediment Hg.
- 40 VT State lakes
  - Water chemistry
  - Macroinvertebrates (NLA)
  - Littoral habitat
  - Sediment mercury
  - Multiprobe
  - Whole phytoplankton
  - VT littoral habitat (subset)

# Field methods





# Progress with data analysis

## Complete\*

- Database construction
- Water chemistry (VT)
- Multiprobe
- Littoral habitat

\*) for 2007 lakes

## In process


- Water chemistry (national lab)
- Whole phytoplankton
- VT littoral habitat (subset)
- Sediment mercury
- Macroinvertebrates (NLA+VT)

# The sample draw as a palate for analyzing existing data

- VT lucked out on the random draw...
- 59 “lakes” provided in total
- 4 rejected as land areas that were misattributed in NHD
- Permission denied on one lake
- Of the remaining lakes in the draw, VT has existing data on *every* one.
- In this section, three thought-experiments:
  - Statewide assessment of trophic condition of VT lakes
  - Statewide assessment of lake trophic trends
  - Use the survey to support nutrient criteria inquiry?

# Statewide assessment of trophic condition and trends

- Most survey lakes have data from 2 or more DEC monitoring programs
- Total count of individual datapoints in existing databases for the survey lakes > 63,000
- Plenty of nutrient data







# Statewide assessment of trophic condition and trends

- Data used:

- Spring Phos. Program: included data for lakes w/4+ years data
- Lay Monitoring Program: included data for lakes w/ at least one summer
- Acid lakes monitoring program
- Kendall-tau statistic used to detect trends, adjusted for experiment-wise error



Trophic State	Secchi Average Summer (m)	Chlorophyll-a Average Summer concentration (ug/L)	TP Spring average concentration (ug/L)	TP Summer photic zone - LMP
Undetermined	--	--	--	--
Hypereutrophic			>100	
Eutrophic	0 - 3.0	> 7.0	> 15	> 15
Mesotrophic	3.0 - 5.5	> 3.5 - 7.0	>7 - 15	>7 - 15
Oligotrophic	> 5.5	0 - 3.5	<7	<7
Dystrophic			<20 (and PtCo > 50)	
Number of NLA draw lakes with data	33	33	49	31



**Figure C.2. Cumulative frequency distributions of total phosphorus (spring and summer), and summer chlorophyll-a and Secchi transparency, based on the National Lakes Survey draw for Vermont. Drop-lines show the percent of lakes exhibiting oligotrophic (blue), mesotrophic (yellow), or eutrophic (green) conditions.**

# Results as compared to the targeted assessment

- Probability predictions accurately predict total assessment from targeted lakes
- A surprising proportion of lakes show detectable trends relative to overall dataset





# Nutrient criteria analyses

- Can the draw be used to assist in the derivation of nutrient criteria?
- Supplement existing analyses of targeted waters, and satisfying the “random” assumption implicit in predicting nutrient responses from stressors using statistics.
- Theoretical example: predicting chlorophyll-a impacts from TP
- For this example, conditional probability analysis was used to evaluate the likelihood of a given chlorophyll-a [conc.] across a range of phosphorus values.

# Conditional probability analysis


- Assume that a mean summer chlorophyll-a above 10 ppb is “undesirable”
- What is the conditional probability of this occurrence given spring total phosphorus?
- Can this inform selection of a potential criterion value?



# Littoral habitat assessment



- Riparian and shoreline zone
  - canopy, cover, slope, and disturbance estimated
- Littoral zone
  - substrate, macrophyte cover, woody debris estimated
- 57 measurements x 10 (or more) locations x 50 lakes (before QC)
  - >28,500 individual data points
  - Thank goodness for Hilary's electronic data capture
- *What a fishing expedition!* Can we use lessons from work of Merrell et al. as a guide?
- In the following slides, results are based on the lakes sampled in 2007.

# Development affects plant growth counter-intuitively





- Merrell et al.\* show that in developed sites, plant cover is more dense.
- They attribute this to removal of tree canopy, permitting additional light penetration to the littoral zone and accentuating nutrient and sediment loss, particularly at shallow sites.

# Hypothesis: canopy cover in the riparian zone influences plant growth





- For survey lakes, where canopy cover is low or absent, macrophyte cover can be significantly greater. There is a breakpoint at 10% canopy.

# Hypothesis: managed sites also have more macrophyte growth






- For survey lakes, sites managed for public recreation display significantly more floating and total plant cover, again implicating light and nutrient effects.

# Development results in loss of coarse woody debris habitat



- Merrell et al. show that in developed sites, there is a significant loss of coarse woody debris, which is a critical component of aquatic habitat. There is also a significant increase in sand.
- This is due to removal of debris and (in most VT lakes) importation of sand for recreational or other purposes.

# Hypothesis: developed sites show less coarse woody debris habitat and more sand



- In the survey lakes, the relative cover of coarse woody debris is significantly reduced, while sand is far more predominant

# Littoral habitat

- Considerable commonalities in findings between the designed work of Merrell et al., and the comparative observational assessment of the NLA.
- More inquiry and analysis of the “Phab” dataset is needed. There are potentially important avenues here for the national analysis.
- To properly explore this, we need to synthesize a “disturbance” variable from the multiple captured parameters.
- Once the full dataset is available, we will comprehensively analyze this dataset.

# Outreach

- Publicized the survey by means of VTANR website.
- Placed column in VTANR's "Out of the Blue" newsletter announcing study, and providing updates.
- Informed VT Federation of Lake Associations of the initiative.
- Solicited assistance from volunteer lakes monitoring community.
- Engaged citizens in sampling. While this often extended the length of the workday, the interaction provided valuable information about the study lakes, as well as educational opportunities.

# In 2008

- Finish field work.
- Develop analytical approach for littoral habitat
- Analysis of state-lake macroinvertebrates by contract (??...in discussion).
- Provide quantified probability estimates of physical, chemical, and biological conditions for VT lakes.
- Look to national analyses as a potential template for reviewing these state-specific data.

# Questions?

Contact:

Neil Kamman

VTDEC-WQ

103 S. Main 10N

Waterbury, VT 05671-0408

[Neil.kamman@state.vt.us](mailto:Neil.kamman@state.vt.us)

# Supplemental slide

- What is we used only simultaneously-collected data from the summer survey to estimate Chla response by TP?

