

# Supplemental Labeling



Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

## Intrepid<sup>®</sup> 2F

EPA Reg. No. 62719-442

### Control or Suppression of Lepidoptera Larvae Infesting Avocado

#### ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Intrepid<sup>®</sup> 2F insecticide before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Intrepid 2F according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Intrepid 2F.

#### Directions for Use

Refer to product label for General Use Precautions, Mixing and Application instructions.

#### Pests, Application Rates, Application Timing and Restrictions:

Pests	Application Rate (fl oz/acre)	Application Timing	Restrictions
lepidopteran larvae including: leafrollers loopers orange tortrix spanworms webbing worms western tussock moth	10 - 16 (0.16 – 0.25 lb ai/acre)	Apply at egg hatch or when first signs of feeding occur. Use the higher rates for heavier infestations and under conditions in which thorough coverage is more difficult. Under heavy infestations, continuous moth flights and/or egg masses and larvae in all stages of development, reapplication can be made at a minimum 6-day re-treatment interval to protect new growth until moth flights and/or hits subside.	<ul style="list-style-type: none"><li>• Do not apply more than 64 fl oz of Intrepid 2F (1 lb ai) per acre or make more than 5 applications per calendar year.</li><li>• <b>Pre-harvest Interval:</b> Do not harvest within 2 days of application.</li></ul>

Because Intrepid 2F must be ingested in order to be effective, it is essential that coverage is thorough and uniform. Higher carrier volumes and higher use rates are recommended for very large trees or dense stands and for heavy target pest infestations. A minimum of six hours drying time is required between the completion of application and the onset of precipitation to ensure optimum performance.

**Ground Application:** Apply a minimum of 50 gallons per acre by conventional ground equipment. Use of higher carrier volume is recommended for heavy infestations and in situations where thorough coverage is difficult to achieve. Equipment and spray volume should be calibrated to assure uniform coverage of infested parts of the crop.

**Aerial Application:** Apply a minimum of 10 gallons per acre. Higher carrier volumes are recommended when environmental conditions are less than ideal for aerial applications.

**Spray Adjuvants:** The addition of agricultural adjuvants to Intrepid 2F sprays will improve initial spray distribution and coverage. Select adjuvants that are recommended and registered for your specific use pattern and follow their use directions. Always add adjuvants last in the mixing process.

**Note:** This product is toxic to aquatic invertebrates. Refer to the Environmental Hazards section of the product label attached to the product container for required protective measures.

®Trademark of Dow AgroSciences LLC

R846-013

EPA accepted: 02/27/08

Initial printing.