

Executive Summary

Central to any study of climate change is the development of an emissions inventory that identifies and quantifies a country's primary anthropogenic¹ sources and sinks of greenhouse gases. This inventory adheres to both 1) a comprehensive and detailed methodology for estimating sources and sinks of anthropogenic greenhouse gases, and 2) a common and consistent mechanism that enables signatory countries to the United Nations Framework Convention on Climate Change (UNFCCC) to compare the relative contribution of different emission sources and greenhouse gases to climate change. Moreover, systematically and consistently estimating national and international emissions is a prerequisite for accounting for reductions and evaluating mitigation strategies.

In 1992, the United States signed and ratified the UNFCCC. The objective of the UNFCCC is “to achieve...stabilization of greenhouse gas concentrations in the atmosphere at a level that would prevent dangerous anthropogenic interference with the climate system. Such a level should be achieved within a time-frame sufficient to allow ecosystems to adapt naturally to climate change, to ensure that food production is not threatened and to enable economic development to proceed in a sustainable manner.”²

Parties to the Convention, by ratifying, committed “to develop, periodically update, publish and make available...national inventories of anthropogenic emissions by sources and removals by sinks of all greenhouse gases not controlled by the Montreal Protocol, using comparable methodologies...”³ The United States views this report as an opportunity to fulfill this commitment.

This chapter summarizes the latest information on U.S. anthropogenic greenhouse gas emission trends from 1990 through 2001. To ensure that the U.S. emissions inventory is comparable to those of other UNFCCC Parties, the estimates presented here were calculated using methodologies consistent with those recommended in the *Revised 1996 IPCC Guidelines for National Greenhouse Gas Inventories* (IPCC/UNEP/OECD/IEA 1997) and the *IPCC Good Practice Guidance and Uncertainty Management in National Greenhouse Gas Inventories* (IPCC 2000). For most source categories, the IPCC methodologies were expanded, resulting in a more comprehensive and detailed estimate of emissions.

Naturally occurring greenhouse gases include water vapor, carbon dioxide (CO₂), methane (CH₄), nitrous oxide (N₂O), and ozone (O₃). Several classes of halogenated substances that contain fluorine, chlorine, or bromine are also greenhouse gases, but they are, for the most part, solely a product of industrial activities. Chlorofluorocarbons (CFCs) and hydrochlorofluorocarbons (HCFCs) are halocarbons that contain chlorine, while halocarbons that contain bromine are

¹ The term “anthropogenic”, in this context, refers to greenhouse gas emissions and removals that are a direct result of human activities or are the result of natural processes that have been affected by human activities (IPCC/UNEP/OECD/IEA 1997).

² Article 2 of the Framework Convention on Climate Change published by the UNEP/WMO Information Unit on Climate Change. See <<http://www.unfccc.int>>.

³ Article 4(1)(a) of the United Nations Framework Convention on Climate Change (also identified in Article 12). See <<http://unfccc.int>>.

referred to as bromofluorocarbons (i.e., halons). Because CFCs, HCFCs, and halons are stratospheric ozone depleting substances, they are covered under the *Montreal Protocol on Substances that Deplete the Ozone Layer*. The UNFCCC defers to this earlier international treaty; consequently these gases are not included in national greenhouse gas inventories.⁴ Some other fluorine containing halogenated substances—hydrofluorocarbons (HFCs), perfluorocarbons (PFCs), and sulfur hexafluoride (SF₆)—do not deplete stratospheric ozone but are potent greenhouse gases. These latter substances are addressed by the UNFCCC and accounted for in national greenhouse gas inventories.

There are also several gases that do not have a direct global warming effect but indirectly affect terrestrial and/or solar radiation absorption by influencing the formation or destruction of other greenhouse gases, including tropospheric and stratospheric ozone. These gases include carbon monoxide (CO), oxides of nitrogen (NO_x), and non-methane volatile organic compounds (NMVOCs). Aerosols, which are extremely small particles or liquid droplets, such as those produced by sulfur dioxide (SO₂) or elemental carbon emissions, can also affect the absorptive characteristics of the atmosphere.

Although the direct greenhouse gases CO₂, CH₄, and N₂O occur naturally in the atmosphere, human activities have changed their atmospheric concentrations. Since the pre-industrial era (i.e., ending about 1750), concentrations of these greenhouse gases have increased by 31, 150, and 16 percent, respectively (IPCC 2001).


Beginning in the 1950s, the use of CFCs and other stratospheric ozone depleting substances (ODSs) increased by nearly 10 percent per year until the mid-1980s, when international concern about ozone depletion led to the entry into force of the *Montreal Protocol*. Since then, the production of ODSs is being phased out. In recent years, use of ODS substitutes such as HFCs and PFCs has grown as they begin to be phased in as replacements for CFCs and HCFCs. Accordingly, atmospheric concentrations of these substitutes have been growing (IPCC 2001).

Recent Trends in U.S. Greenhouse Gas Emissions

In 2001, total U.S. greenhouse gas emissions were 6,936.2 teragrams of carbon dioxide equivalents (Tg CO₂ Eq.)⁵ (13.0 percent above 1990 emissions). Emissions declined for the second time since the base year 1990, decreasing by 1.6 percent (111.2 Tg CO₂ Eq.) from 2000 to 2001, driven primarily by decreases in CO₂ emissions from fossil fuel combustion. The following factors were primary contributors to this decrease: 1) slow economic growth in 2001, leading to decreased demand for electricity fuels, 2) a considerable reduction in industrial output, leading to decreased demand for electricity and fossil fuel, 3) warmer winter conditions compared to 2000, and 4) increased output from nuclear facilities. (See the following section for an analysis of emission trends by general economic sectors.)

Figure ES-1 through Figure ES-3 illustrate the overall trends in total U.S. emissions by gas, annual changes, and absolute change since 1990. Table ES-1 provides a detailed summary of U.S. greenhouse gas emissions and sinks for 1990 through 2001.

Figure ES-1



⁴ Emissions estimates of CFCs, HCFCs, halons and other ozone-depleting substances are included in this document for informational purposes.

⁵ Estimates are presented in units of teragrams of carbon dioxide equivalents (Tg CO₂ Eq.), which weight each gas by its Global Warming Potential, or GWP, value. (See section on Global Warming Potentials, Chapter 1.)

Table ES-1: Recent Trends in U.S. Greenhouse Gas Emissions and Sinks (Tg CO₂ Eq.)

| Gas/Source | 1990 | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 |
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|
| CO₂ | 5,003.7 | 5,334.4 | 5,514.8 | 5,595.4 | 5,614.2 | 5,680.7 | 5,883.1 | 5,794.8 |
| Fossil Fuel Combustion | 4,814.8 | 5,141.5 | 5,325.8 | 5,400.0 | 5,420.5 | 5,488.8 | 5,692.2 | 5,614.9 |
| Iron and Steel Production | 85.4 | 74.4 | 68.3 | 71.9 | 67.4 | 64.4 | 65.8 | 59.1 |
| Cement Manufacture | 33.3 | 36.8 | 37.1 | 38.3 | 39.2 | 40.0 | 41.2 | 41.4 |
| Waste Combustion | 14.1 | 18.5 | 19.4 | 21.2 | 22.5 | 23.9 | 25.4 | 26.9 |
| Ammonia Manufacture & Urea Application | 19.3 | 20.5 | 20.3 | 20.7 | 21.9 | 20.6 | 19.6 | 16.6 |
| Lime Manufacture | 11.2 | 12.8 | 13.5 | 13.7 | 13.9 | 13.5 | 13.3 | 12.9 |
| Natural Gas Flaring | 5.5 | 8.7 | 8.2 | 7.6 | 6.3 | 6.7 | 5.5 | 5.2 |
| Limestone and Dolomite Use | 5.5 | 7.0 | 7.6 | 7.1 | 7.3 | 7.7 | 5.8 | 5.3 |
| Aluminum Production | 6.3 | 5.3 | 5.6 | 5.6 | 5.8 | 5.9 | 5.4 | 4.1 |
| Soda Ash Manufacture and Consumption | 4.1 | 4.3 | 4.2 | 4.4 | 4.3 | 4.2 | 4.2 | 4.1 |
| Titanium Dioxide Production | 1.3 | 1.7 | 1.7 | 1.8 | 1.8 | 1.9 | 1.9 | 1.9 |
| Carbon Dioxide Consumption | 0.9 | 1.1 | 1.1 | 1.2 | 1.2 | 1.2 | 1.2 | 1.3 |
| Ferroalloys | 2.0 | 1.9 | 2.0 | 2.0 | 2.0 | 2.0 | 1.7 | 1.3 |
| Land-Use Change and Forestry (Sink) ^a | (1,072.8) | (1,064.2) | (1,061.0) | (840.6) | (830.5) | (841.1) | (834.6) | (838.1) |
| <i>International Bunker Fuels^b</i> | <i>113.9</i> | <i>101.0</i> | <i>102.3</i> | <i>109.9</i> | <i>112.9</i> | <i>105.3</i> | <i>99.3</i> | <i>97.3</i> |
| CH₄ | 644.0 | 650.0 | 636.8 | 629.5 | 622.7 | 615.5 | 613.4 | 605.9 |
| Landfills | 212.1 | 216.1 | 212.1 | 207.5 | 202.4 | 203.7 | 205.8 | 202.9 |
| Natural Gas Systems | 122.0 | 127.2 | 127.4 | 126.0 | 124.0 | 120.3 | 121.2 | 117.3 |
| Enteric Fermentation | 117.9 | 123.0 | 120.5 | 118.3 | 116.7 | 116.6 | 115.7 | 114.8 |
| Coal Mining | 87.1 | 73.5 | 68.4 | 68.1 | 67.9 | 63.7 | 60.9 | 60.7 |
| Manure Management | 31.3 | 36.2 | 34.9 | 36.6 | 39.0 | 38.9 | 38.2 | 38.9 |
| Wastewater Treatment | 24.1 | 26.6 | 26.8 | 27.3 | 27.7 | 28.2 | 28.3 | 28.3 |
| Petroleum Systems | 27.5 | 24.2 | 23.9 | 23.6 | 22.9 | 21.6 | 21.2 | 21.2 |
| Rice Cultivation | 7.1 | 7.6 | 7.0 | 7.5 | 7.9 | 8.3 | 7.5 | 7.6 |
| Stationary Sources | 8.1 | 8.5 | 8.7 | 7.5 | 7.2 | 7.4 | 7.6 | 7.4 |
| Mobile Sources | 5.0 | 4.9 | 4.8 | 4.7 | 4.6 | 4.5 | 4.4 | 4.3 |
| Petrochemical Production | 1.2 | 1.5 | 1.6 | 1.6 | 1.6 | 1.7 | 1.7 | 1.5 |
| Field Burning of Agricultural Residues | 0.7 | 0.7 | 0.7 | 0.8 | 0.8 | 0.8 | 0.8 | 0.8 |
| Silicon Carbide Production | + | + | + | + | + | + | + | + |
| <i>International Bunker Fuels^b</i> | <i>0.2</i> | <i>0.1</i> | <i>0.1</i> | <i>0.1</i> | <i>0.1</i> | <i>0.1</i> | <i>0.1</i> | <i>0.1</i> |
| N₂O | 397.6 | 430.9 | 441.7 | 440.9 | 436.8 | 430.0 | 429.9 | 424.6 |
| Agricultural Soil Management | 267.5 | 284.1 | 293.2 | 298.2 | 299.2 | 297.0 | 294.6 | 294.3 |
| Mobile Sources | 50.6 | 60.9 | 60.7 | 60.3 | 59.7 | 58.8 | 57.5 | 54.8 |
| Manure Management | 16.2 | 16.6 | 17.0 | 17.3 | 17.3 | 17.4 | 17.9 | 18.0 |
| Nitric Acid | 17.8 | 19.9 | 20.7 | 21.2 | 20.9 | 20.1 | 19.1 | 17.6 |
| Human Sewage | 12.7 | 13.9 | 14.1 | 14.4 | 14.6 | 15.1 | 15.1 | 15.3 |
| Stationary Combustion | 12.5 | 13.2 | 13.8 | 13.7 | 13.7 | 13.7 | 14.3 | 14.2 |
| Adipic Acid | 15.2 | 17.2 | 17.0 | 10.3 | 6.0 | 5.5 | 6.0 | 4.9 |
| N ₂ O Product Usage | 4.3 | 4.5 | 4.5 | 4.8 | 4.8 | 4.8 | 4.8 | 4.8 |
| Field Burning of Agricultural Residues | 0.4 | 0.4 | 0.4 | 0.4 | 0.5 | 0.4 | 0.5 | 0.5 |
| Waste Combustion | 0.3 | 0.3 | 0.3 | 0.3 | 0.2 | 0.2 | 0.2 | 0.2 |
| <i>International Bunker Fuels^b</i> | <i>1.0</i> | <i>0.9</i> | <i>0.9</i> | <i>1.0</i> | <i>1.0</i> | <i>0.9</i> | <i>0.9</i> | <i>0.9</i> |
| HFCs, PFCs, and SF₆ | 94.4 | 99.5 | 113.6 | 116.8 | 127.6 | 120.3 | 121.0 | 111.0 |
| Substitution of Ozone Depleting Substances | 0.9 | 21.7 | 30.4 | 37.7 | 44.5 | 50.9 | 57.3 | 63.7 |
| HCFC-22 Production | 35.0 | 27.0 | 31.1 | 30.0 | 40.2 | 30.4 | 29.8 | 19.8 |
| Electrical Transmission and Distribution | 32.1 | 27.5 | 27.7 | 25.2 | 20.9 | 16.4 | 15.4 | 15.3 |
| Semiconductor Manufacture | 2.9 | 5.9 | 5.4 | 6.5 | 7.3 | 7.7 | 7.4 | 5.5 |
| Aluminum Production | 18.1 | 11.8 | 12.5 | 11.0 | 9.0 | 8.9 | 7.9 | 4.1 |
| Magnesium Production and Processing | 5.4 | 5.6 | 6.5 | 6.3 | 5.8 | 6.0 | 3.2 | 2.5 |
| Total | 6,139.6 | 6,514.9 | 6,707.0 | 6,782.6 | 6,801.3 | 6,849.5 | 7,047.4 | 6,936.2 |
| Net Emissions (Sources and Sinks) | 5,066.8 | 5,450.7 | 5,646.0 | 5,942.0 | 5,970.9 | 6,008.5 | 6,212.7 | 6,098.1 |

+ Does not exceed 0.05 Tg CO₂ Eq.

^a For the most recent years, a portion of the sink estimate is based on historical and projected data; see Chapter 6, Table 6-1 for a complete breakdown. Parentheses indicate negative values (or sequestration).

^b Emissions from International Bunker Fuels are not included in totals.

Note: Totals may not sum due to independent rounding.

Figure ES-2




Figure ES-4




Figure ES-3




Figure ES-4 illustrates the relative contribution of the direct greenhouse gases to total U.S. emissions in 2001. The primary greenhouse gas emitted by human activities in the United States was CO₂, representing approximately 84 percent of total greenhouse gas emissions. The largest source of CO₂, and of overall greenhouse gas emissions, was fossil fuel combustion. Methane emissions resulted primarily from decomposition of wastes in landfills, natural gas systems, and enteric fermentation associated with domestic livestock. Agricultural soil management and mobile source fossil fuel combustion were the major sources of N₂O emissions. The emissions of substitutes for ozone depleting substances and emissions of HFC-23 during the production of HCFC-22 were the primary contributors to aggregate HFC emissions. Electrical transmission and

distribution systems accounted for most SF₆ emissions, while the majority of PFC emissions resulted as a by-product of primary aluminum production.

As the largest source of U.S. greenhouse gas emissions, CO₂ from fossil fuel combustion accounted for a nearly constant 80 percent of global warming potential (GWP) weighted emissions in the 1990s. Emissions from this source category grew by 17 percent (800.1 Tg CO₂Eq.) from 1990 to 2001 and were responsible for most of the increase in national emissions during this period. The most recent annual change in CO₂ emissions from fossil fuel combustion was a reduction of 77.3 Tg CO₂ Eq. (1.4 percent), which is the first decrease in emissions since a decrease from 1990 to 1991; the source's average annual growth rate was 1.3 percent from 1990 through 2001. Historically, changes in emissions from fossil fuel combustion have been the dominant factor affecting U.S. emission trends.

Changes in CO₂ emissions from fossil fuel combustion are influenced by many long-term and short-term factors, including population and economic growth, energy price fluctuations, technological changes, and seasonal temperatures. On an annual basis, the overall consumption of fossil fuels in the United States generally fluctuates in response to changes in general economic conditions, energy prices, weather, and the availability of non-fossil alternatives. For example, in a year with increased consumption of goods and services, low fuel prices, severe summer and winter weather conditions, nuclear plant closures, and lower precipitation feeding hydroelectric dams, there would likely be proportionally greater fossil fuel consumption than a year

with poor economic performance, high fuel prices, mild temperatures, and increased output from nuclear and hydroelectric plants.

In the longer-term energy consumption patterns respond to changes that affect the scale of consumption (e.g., population, number of cars, and size of houses), the efficiency with which energy is used in equipment (e.g., cars, power plants, steel mills, and light bulbs) and consumer behavior (e.g., walking, bicycling, or telecommuting to work instead of driving).

Energy-related CO₂ emissions also depend on the type of fuel or energy consumed and its carbon intensity. Producing a unit of heat or electricity using natural gas instead of coal, for example, can reduce the CO₂ emissions because of the lower carbon content of natural gas. Table ES-2 shows annual changes in emissions during the last six years for coal, petroleum, and natural gas in selected sectors.

Milder weather conditions in summer and winter of 1997 (compared to 1996) moderated the growth of CO₂ emissions from fossil fuel combustion. The shutdown of several nuclear power plants, however, lead electric utilities to increase their consumption of coal and other fuels to offset the lost nuclear capacity.

In 1998, warm winter temperatures contributed to a significant drop in residential and commercial natural gas consumption. This drop in emissions from natural gas used for heating was offset by two factors: 1) electric utility

emissions, which increased in part due to a hot summer and its associated air conditioning demand; and 2) increased motor gasoline consumption for transportation.

In 1999, the increase in emissions from fossil fuel combustion was driven largely by growth in petroleum consumption for transportation. In addition, residential and commercial heating fuel demand partially recovered as winter temperatures dropped relative to 1998, although temperatures were still warmer than normal.⁶ These increases were offset, in part, by a decline in emissions from electric power producers due primarily to: 1) an increase in net generation of electricity by nuclear plants which reduced demand from fossil fuel plants; and 2) moderated summer temperatures compared to the previous year—thereby reducing electricity demand for air conditioning.

Emissions from fuel combustion increased considerably in 2000, due to several factors. The primary reason for the increase was the robust U.S. economy, which produced a high demand for fuels—especially for petroleum in the transportation sector—despite increases in the price of both natural gas and petroleum. Colder winter conditions relative to the previous year triggered a rise in residential and commercial demand for heating. Structural and other economic changes taking place within U.S. industry—especially manufacturing—lead to lower coal consumption. Additionally, electricity generation became more carbon intensive as coal and natural gas consumption offset reduced hydropower output.

Table ES-2: Annual Change in CO₂ Emissions from Fossil Fuel Combustion for Selected Fuels and Sectors (Tg CO₂ Eq. and Percent)

| Sector | Fuel Type | 1996 to 1997 | | 1997 to 1998 | | 1998 to 1999 | | 1999 to 2000 | | 2000 to 2001 | |
|--------------------------------|------------------------------|--------------|-----------|--------------|-----------|--------------|-----------|--------------|-----------|---------------|-------------|
| Electricity Generation | Coal | 44.1 | 3% | 30.6 | 2% | 8.7 | 0% | 89.4 | 5% | (50.1) | (3%) |
| Electricity Generation | Natural Gas | 13.9 | 7% | 29.1 | 13% | 12.0 | 5% | 20.6 | 8% | 4.3 | 2% |
| Electricity Generation | Petroleum | 9.0 | 14% | 29.7 | 40% | (7.6) | (7%) | (5.7) | (6%) | 10.7 | 12% |
| Transportation ^a | Petroleum | 7.3 | 0% | 33.0 | 2% | 58.7 | 4% | 49.8 | 3% | 19.7 | 1% |
| Residential | Natural Gas | (14.0) | (3%) | (23.7) | (9%) | 10.0 | 9% | 13.9 | 3% | (9.6) | (1%) |
| Commercial | Natural Gas | 3.1 | 2% | (10.8) | (6%) | 1.7 | 1% | 9.1 | 5% | 1.6 | 1% |
| Industrial | Coal | 1.2 | 1% | (8.7) | (6%) | (6.1) | (4%) | 2.3 | 2% | (7.7) | (6%) |
| Industrial | Natural Gas | 1.1 | 0% | (11.7) | (2%) | (17.6) | (4%) | 12.5 | 3% | (33.5) | (7%) |
| All Sectors^b | All Fuels^b | 74.2 | 1% | 20.5 | 0% | 68.3 | 1% | 202.8 | 4% | (77.9) | (1%) |

^a Excludes emissions from International Bunker Fuels.

^b Includes fuels and sectors not shown in table.

⁶ Normals are based on data from 1961 through 1990. Source: NOAA (2002)

In 2001, economic growth in the U.S. slowed considerably for the second time since 1990, contributing to a decrease in CO₂ emissions from fossil fuel combustion, also for the second time since 1990. A significant reduction in industrial output contributed to weak economic growth, primarily in manufacturing, and led to lower emissions from the industrial sector. Several other factors also played a role in this decrease in emissions. Warmer winter conditions compared to 2000, along with higher natural gas prices, reduced demand for heating fuels. Additionally, nuclear facilities operated at their highest capacity on record, offsetting electricity produced from fossil fuels. Since there are no greenhouse gas emissions associated with electricity production from nuclear plants, this substitution reduces the overall carbon intensity of electricity generation.

Other significant trends in emissions from additional source categories over the twelve-year period from 1990 through 2001 included the following:

- Carbon dioxide emissions from waste combustion increased by 12.8 Tg CO₂ Eq. (91 percent), as the volume of plastics and other fossil carbon-containing materials in municipal solid waste grew.
- Net CO₂ flux from land use change and forestry decreased by 234.7 Tg CO₂ Eq. (22 percent), primarily due to a decline in the rate of net carbon accumulation in forest carbon stocks.
- Methane emissions from coal mining dropped by 26.4 Tg CO₂ Eq. (30 percent) as a result of the mining of less gassy coal from underground mines and the increased use of methane collected from degasification systems.
- By 1998, all of the three major adipic acid producing plants had voluntarily implemented N₂O abatement technology, and as a result, emissions fell by 10.3 Tg CO₂ Eq. (68 percent). The majority of this decline occurred from 1996 to 1998, despite increased production.
- Nitrous oxide emissions from agricultural soil management increased by 26.8 Tg CO₂ Eq. (10 percent) as crop and forage production, manure production, and fertilizer consumption rose.

- Aggregate HFC and PFC emissions resulting from the substitution of ozone depleting substances (e.g., CFCs) increased by 62.7 Tg CO₂ Eq. This increase was significantly offset, however, by reductions in PFC emissions from aluminum production (14.0 Tg CO₂ Eq. or 77 percent), reductions in emissions of HFC-23 from the production of HCFC-22 (15.2 Tg CO₂ Eq. or 43 percent), and reductions of SF₆ from electric power transmission and distribution systems (16.8 Tg CO₂ Eq. or 52 percent). Reductions in PFC emissions from aluminum production resulted from both industry emission reduction efforts and lower domestic aluminum production. HFC-23 emissions from the production of HCFC-22 decreased because a reduction in the intensity of emissions from that source offset an increase in HCFC-22 production. Reduced emissions of SF₆ from electric power transmission and distribution systems are primarily the result of higher purchase prices for SF₆ and efforts by industry to reduce emissions.


Overall, from 1990 to 2001, total emissions of CO₂ and N₂O increased by 791.1 (16 percent) and 27.0 Tg CO₂ Eq. (7 percent), respectively, while CH₄ emissions decreased by 38.1 Tg CO₂ Eq. (6 percent). During the same period, aggregate weighted emissions of HFCs, PFCs, and SF₆ rose by 16.6 Tg CO₂ Eq. (18 percent). Despite being emitted in smaller quantities relative to the other principal greenhouse gases, emissions of HFCs, PFCs, and SF₆ are significant because many of them have extremely high global warming potentials and, in the cases of PFCs and SF₆, long atmospheric lifetimes. Conversely, U.S. greenhouse gas emissions were partly offset by carbon sequestration in forests, trees in urban areas, agricultural soils, and landfilled yard trimmings, which was estimated to be 12 percent of total emissions in 2001.

Emissions by Economic Sector

Throughout this report, emission estimates are grouped into six sectors (i.e., chapters) defined by the IPCC: Energy, Industrial Processes, Solvent Use, Agriculture, Land-Use Change and Forestry, and Waste. While it is important to use this characterization for consistency with UNFCCC reporting guidelines, it is also useful to allocate emissions into more commonly used sectoral categories. This section reports emissions by the following economic sectors: Residential, Commercial, Industry, Transportation, Electricity Generation, and Agriculture, and U.S. Territories. Table ES-3 summarizes emissions from each of these sectors. Figure ES-5 shows the trend in emissions by sector from 1990 to 2001.

Using this categorization, emissions from electricity generation accounted for the largest portion (33 percent) of U.S. greenhouse gas emissions in 2001. Transportation activities, in aggregate, accounted for the second largest portion (27 percent). Emissions from industry accounted for 19 percent of U.S. greenhouse gas emissions in 2001. In contrast to electricity generation and transportation, emissions from industry have declined over the past decade, as structural changes have occurred in the U.S. economy (i.e., shifts from a manufacturing based to a service-based economy), fuel switching has occurred, and efficiency improvements have been made. The remaining 21 percent of U.S. greenhouse gas emissions were contributed by the

Figure ES-5



residential, agriculture, and commercial economic sectors, plus emissions from U.S. Territories. Residences accounted for about 5 percent, and primarily consisted of CO₂ emissions from fossil fuel combustion. Activities related to agriculture accounted for roughly 8 percent of U.S. emissions, but unlike all other economic sectors, these emissions were dominated by N₂O emissions from agricultural soils instead of CO₂ from fossil fuel combustion. The commercial sector accounted for about 7 percent of emissions, while U.S. territories accounted for 1 percent.

Table ES-3: U.S. Greenhouse Gas Emissions Allocated to Economic Sectors (Tg CO₂ Eq.)

| Sector | 1990 | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 |
|--|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|
| Electricity Generation | 1,862 | 1,990 | 2,064 | 2,130 | 2,217 | 2,227 | 2,332 | 2,298 |
| Transportation | 1,526 | 1,651 | 1,694 | 1,707 | 1,736 | 1,798 | 1,849 | 1,867 |
| Industry | 1,423 | 1,445 | 1,485 | 1,485 | 1,433 | 1,381 | 1,401 | 1,316 |
| Agriculture | 488 | 526 | 526 | 538 | 539 | 540 | 526 | 526 |
| Residential | 335 | 371 | 402 | 385 | 353 | 372 | 390 | 379 |
| Commercial | 472 | 488 | 495 | 494 | 474 | 481 | 498 | 497 |
| U.S. Territories | 34 | 44 | 40 | 43 | 48 | 50 | 52 | 54 |
| Total | 6,140 | 6,515 | 6,707 | 6,783 | 6,801 | 6,850 | 7,047 | 6,936 |
| Sinks | (1,073) | (1,064) | (1,061) | (841) | (830) | (841) | (835) | (838) |
| Net Emissions (Sources and Sinks) | 5,067 | 5,451 | 5,646 | 5,942 | 5,971 | 6,008 | 6,213 | 6,098 |


Note: Totals may not sum. Emissions include CO₂, CH₄, HFCs, PFCs, and SF₆. See Table 1-13 for more detailed data.

Carbon dioxide was also emitted and sequestered by a variety of activities related to forest management practices, tree planting in urban areas, the management of agricultural soils, and landfilling of yard trimmings.

Electricity is ultimately consumed in the economic sectors described above. Table ES-4 presents greenhouse gas emissions from economic sectors with emissions related to electricity generation distributed into end-use categories (i.e., emissions from electricity generation are allocated to the economic sectors in which the electricity is consumed). To distribute electricity emissions among end-use sectors, emissions from the source categories assigned to electricity generation were allocated to the residential, commercial, industry, transportation, and agriculture economic sectors according to retail sales of electricity.⁷ These source categories include CO₂ from fossil fuel combustion and the use of limestone and dolomite for flue gas desulfurization, CO₂ and N₂O from waste combustion, CH₄ and N₂O from stationary sources, and SF₆ from electrical transmission and distribution systems.

When emissions from electricity are distributed among these sectors, industry accounts for the largest share of U.S. greenhouse gas emissions (30 percent) in 2001. Emissions from the residential and commercial sectors also increase substantially due to their relatively large share of electricity consumption (e.g., lighting, appliances, etc.). Transportation activities remain the second largest

Figure ES-6



contributor to emissions. In all sectors except agriculture, CO₂ accounts for more than 75 percent of greenhouse gas emissions, primarily from the combustion of fossil fuels. Figure ES-6 shows the trend in these emissions by sector from 1990 to 2001.

Table ES-4: U.S Greenhouse Gas Emissions by Economic Sector and Gas with Electricity-Related Emissions Distributed (Tg CO₂ Eq.)

| Sector | 1990 | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 |
|--|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|
| Industry | 2,097 | 2,164 | 2,224 | 2,245 | 2,209 | 2,165 | 2,201 | 2,074 |
| Transportation | 1,529 | 1,654 | 1,697 | 1,710 | 1,740 | 1,801 | 1,852 | 1,870 |
| Residential | 943 | 1,020 | 1,081 | 1,074 | 1,076 | 1,095 | 1,155 | 1,139 |
| Commercial | 1,025 | 1,081 | 1,110 | 1,151 | 1,164 | 1,178 | 1,241 | 1,253 |
| Agriculture | 512 | 553 | 555 | 560 | 564 | 560 | 547 | 546 |
| US Territories | 34 | 44 | 40 | 43 | 48 | 50 | 52 | 54 |
| Total | 6,140 | 6,515 | 6,707 | 6,783 | 6,801 | 6,850 | 7,047 | 6,936 |
| Sinks | (1,073) | (1,064) | (1,061) | (841) | (830) | (841) | (835) | (838) |
| Net Emissions (Sources and Sinks) | 5,067 | 5,451 | 5,646 | 5,942 | 5,971 | 6,008 | 6,213 | 6,098 |

See Table 1-11 for more detailed data.

⁷ Emissions were not distributed to U.S. territories, since the electricity generation sector only includes emissions related to the generation of electricity in the 50 states and the District of Columbia.

Box ES-1: Recent Trends in Various U.S. Greenhouse Gas Emissions-Related Data

Total emissions can be compared to other economic and social indices to highlight changes over time. These comparisons include: 1) emissions per unit of aggregate energy consumption, because energy-related activities are the largest sources of emissions; 2) emissions per unit of fossil fuel consumption, because almost all energy-related emissions involve the combustion of fossil fuels; 3) emissions per unit of electricity consumption, because the electric power industry—utilities and nonutilities combined—was the largest source of U.S. greenhouse gas emissions in 2001; 4) emissions per unit of total gross domestic product as a measure of national economic activity; or 5) emissions per capita.

Table ES-5 provides data on various statistics related to U.S. greenhouse gas emissions normalized to 1990 as a baseline year. Greenhouse gas emissions in the U.S. have grown at an average annual rate of 1.1 percent since 1990. This rate is slower than that for total energy or fossil fuel consumption and much slower than that for either electricity consumption or overall gross domestic product. At the same time, total U.S. greenhouse gas emissions have grown at about the same rate as national population during the last decade (see Figure ES-7). Overall, global atmospheric CO₂ concentrations—a function of many complex anthropogenic and natural processes—are increasing at 0.4 percent per year.

Table ES-5: Recent Trends in Various U.S. Data (Index 1990 = 100) and Global Atmospheric CO₂ Concentration

| Variable | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | Growth Rate ^f |
|--|------|------|------|------|------|------|------|------|------|------|------|--------------------------|
| Greenhouse Gas Emissions ^a | 99 | 101 | 103 | 105 | 106 | 109 | 110 | 111 | 112 | 115 | 113 | 1.1% |
| Energy Consumption ^b | 100 | 102 | 104 | 106 | 108 | 112 | 112 | 113 | 115 | 118 | 115 | 1.3% |
| Fossil Fuel Consumption ^b | 100 | 102 | 104 | 106 | 107 | 111 | 112 | 113 | 114 | 118 | 115 | 1.3% |
| Electricity Consumption ^b | 102 | 102 | 106 | 109 | 112 | 115 | 117 | 121 | 124 | 128 | 127 | 2.2% |
| GDP ^c | 100 | 103 | 105 | 110 | 112 | 116 | 122 | 127 | 132 | 137 | 137 | 2.9% |
| Population ^d | 101 | 103 | 104 | 105 | 107 | 108 | 109 | 111 | 112 | 113 | 114 | 1.2% |
| Atmospheric CO ₂ Concentration ^e | 100 | 101 | 101 | 101 | 102 | 102 | 103 | 104 | 104 | 104 | 105 | 0.4% |

^a GWP weighted values

^b Energy content weighted values (EIA 2002a)


^c Gross Domestic Product in chained 1996 dollars (BEA 2002)

^d (U.S. Census Bureau 2002)

^e Mauna Loa Observatory, Hawaii (Keeling and Whorf 2002)

^f Average annual growth rate

Figure ES-7



Global Warming Potentials

Gases in the atmosphere can contribute to the greenhouse effect both directly and indirectly. Direct effects occur when the gas itself absorbs radiation. Indirect radiative forcing occurs when chemical transformations of the substance produce other greenhouse gases, when a gas influences the atmospheric lifetimes of other gases, and/or when a gas affects atmospheric processes that alter the radiative balance of the Earth (e.g., affect cloud formation or albedo).⁸ The IPCC developed the Global Warming Potential (GWP) concept to compare the ability of each greenhouse gas to trap heat in the atmosphere relative to another gas.

The GWP of a greenhouse gas is defined as the ratio of the time-integrated radiative forcing from the instantaneous release of 1 kg of a trace substance relative to that of 1 kg of a reference gas (IPCC 2001). Direct radiative effects occur when the gas itself is a greenhouse gas. The reference gas used is CO₂, and therefore GWP-weighted emissions are measured in teragrams of CO₂ equivalents (Tg CO₂ Eq.).⁹ All gases in this executive summary are presented in units of Tg CO₂ Eq. The relationship between gigagrams (Gg) of a gas and Tg CO₂ Eq. can be expressed as follows:

$$\text{Tg CO}_2 \text{ Eq} = (\text{Gg of gas}) \times (\text{GWP}) \times \left(\frac{\text{Tg}}{1,000\text{Gg}} \right)$$

While any time period can be selected, this report uses the 100 year GWPs recommended by the IPCC, and adopted by the UNFCCC for reporting purposes (IPCC 1996). GWP values are listed in Table ES-6.

Global warming potentials are not provided for CO, NO_x, NMVOCs, SO₂, and aerosols because there is no agreed upon method to estimate the contribution of gases that are short-lived in the atmosphere, spatially variable, and have only indirect effects on radiative forcing (IPCC 1996).

Table ES-6: Global Warming Potentials (100 Year Time Horizon) Used in This Report

| Gas | GWP |
|-----------------------------------|--------|
| Carbon dioxide (CO ₂) | 1 |
| Methane (CH ₄)* | 21 |
| Nitrous oxide (N ₂ O) | 310 |
| HFC-23 | 11,700 |
| HFC-32 | 650 |
| HFC-125 | 2,800 |
| HFC-134a | 1,300 |
| HFC-143a | 3,800 |
| HFC-152a | 140 |
| HFC-227ea | 2,900 |
| HFC-236fa | 6,300 |
| HFC-4310mee | 1,300 |
| CF ₄ | 6,500 |
| C ₂ F ₆ | 9,200 |
| C ₄ F ₁₀ | 7,000 |
| C ₆ F ₁₄ | 7,400 |
| SF ₆ | 23,900 |

Source: IPCC (1996)

* The methane GWP includes the direct effects and those indirect effects due to the production of tropospheric ozone and stratospheric water vapor. The indirect effect due to the production of CO₂ is not included.

Carbon Dioxide Emissions

The global carbon cycle is made up of large carbon flows and reservoirs. Billions of tons of carbon in the form of CO₂ are absorbed by oceans and living biomass (i.e., sinks) and are emitted to the atmosphere annually through natural processes (i.e., sources). When in equilibrium, carbon fluxes among these various reservoirs are roughly balanced. Since the Industrial Revolution, this equilibrium of atmospheric carbon has been disrupted. Atmospheric concentrations of CO₂ have risen about 31 percent (IPCC 2001), principally because of fossil fuel combustion, which accounted for 97 percent of total U.S. CO₂ emissions in 2001. Globally, approximately 23,300 Tg of CO₂ were added to the atmosphere through the combustion of fossil fuels at the end of the 1990s, of which the United States accounted for about 24 percent (see Figure ES-8).¹⁰ Changes in land

⁸ Albedo is a measure of the Earth's reflectivity; see the Glossary (Annex AB) for definition.

⁹ Carbon comprises 12/44ths of carbon dioxide by weight.

¹⁰ Global CO₂ emissions from fossil fuel combustion were taken from Marland et al. (2002). <http://cdiac.esd.ornl.gov/trends/emis/meth_reg.htm>

Box ES-2: The IPCC Third Assessment Report and Global Warming Potentials

In its Third Assessment Report (TAR), the IPCC updated the global warming potentials (GWPs) of several gases relative to the Second Assessment Report (SAR), and new GWPs were calculated for an expanded set of gases. Since the Second Assessment Report, the IPCC has applied an improved calculation of CO₂ radiative forcing and an improved CO₂ response function (presented in WMO 1999). The GWPs in the TAR are drawn from WMO (1999) and the Second Assessment Report, with updates for those cases where significantly different new laboratory or radiative transfer results have been published. Additionally, the atmospheric lifetimes of some gases were recalculated. Because the revised radiative forcing of CO₂ is about 12 percent lower than that in the Second Assessment Report, the GWPs of the other gases relative to CO₂ tend to be larger, taking into account revisions in lifetimes. In addition, the values for radiative forcing and lifetimes have been calculated for a variety of halocarbons that were not presented in the Second Assessment Report. Table ES-7 presents the new Global Warming Potentials, relative to those presented in the Second Assessment Report.

Table ES-7: Comparison of 100 Year GWPs

| Gas | SAR | TAR | Change | |
|-----------------------------------|--------|--------|---------|-------|
| Carbon dioxide (CO ₂) | 1 | 1 | NC | NC |
| Methane (CH ₄)* | 21 | 23 | 2 | 10% |
| Nitrous oxide (N ₂ O) | 310 | 296 | (14) | (5%) |
| HFC-23 | 11,700 | 12,000 | 300 | 3% |
| HFC-32 | 650 | 550 | (100) | (15%) |
| HFC-125 | 2,800 | 3,400 | 600 | 21% |
| HFC-134a | 1,300 | 1,300 | NC | NC |
| HFC-143a | 3,800 | 4,300 | 500 | 13% |
| HFC-152a | 140 | 120 | (20) | (14%) |
| HFC-227ea | 2,900 | 3,500 | 600 | 21% |
| HFC-236fa | 6,300 | 9,400 | 3,100 | 49% |
| HFC-4310mee | 1,300 | 1,500 | 200 | 15% |
| CF ₄ | 6,500 | 5,700 | (800) | (12%) |
| C ₂ F ₆ | 9,200 | 11,900 | 2,700 | 29% |
| C ₄ F ₁₀ | 7,000 | 8,600 | 1,600 | 23% |
| C ₆ F ₁₄ | 7,400 | 9,000 | 1,600 | 22% |
| SF ₆ | 23,900 | 22,200 | (1,700) | (7%) |

Source: (IPCC 2001, 1996)

NC (No Change)

* The methane GWP includes the direct effects and those indirect effects due to the production of tropospheric ozone and stratospheric water vapor. The indirect effect due to the production of CO₂ is not included.

Although the GWPs have been updated by the IPCC, estimates of emissions presented in this report use the GWPs from the Second Assessment Report. The UNFCCC reporting guidelines for national inventories¹¹ were updated in 2002, but continue to require the use of GWPs from the SAR so that current estimates of aggregated greenhouse gas emissions for 1990 through 2001 are consistent with estimates developed prior to the publication of the TAR. Therefore, to comply with international reporting standards under the UNFCCC, official emission estimates are reported by the United States using SAR GWP values. Overall, these revisions to GWP values do not have a significant effect on U.S. emission trends, and all estimates are provided throughout this report in both CO₂ equivalents and unweighted units. A comparison of emission values using the SAR GWPs versus the TAR GWPs can be found in Chapter 1 and in more detail in Annex S.

use and forestry practices can also emit CO₂ (e.g., through conversion of forest land to agricultural or urban use) or can act as a sink for CO₂ (e.g., through net additions to forest biomass).

Figure ES-9 and Table ES-8 summarize U.S. sources and sinks of CO₂. The remainder of this section discusses CO₂ emission trends in greater detail.

¹¹ See <http://unfccc.int/cop8/latest/5_sbsta15add1.pdf>.

Figure ES-8





Figure ES-9



Energy

Energy-related activities, primarily fossil fuel combustion, accounted for the vast majority of U.S. CO₂ emissions for the period of 1990 through 2001. In 2001, approximately 86 percent of the energy consumed in the United States was produced through the combustion of fossil fuels. The remaining 14 percent came from other energy sources such as hydropower, biomass, nuclear, wind, and solar energy (see Figure ES-10 and Figure ES-11). A discussion of specific trends related to CO₂ emissions from energy consumption is presented below.

Table ES-8: U.S. Sources of CO₂ Emissions and Sinks (Tg CO₂ Eq.)

| Source or Sink | 1990 | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 |
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|
| Fossil Fuel Combustion | 4,814.8 | 5,141.5 | 5,325.8 | 5,400.0 | 5,420.5 | 5,488.8 | 5,692.2 | 5,614.9 |
| Electricity Generation | 1,805.0 | 1,931.8 | 2,003.9 | 2,070.8 | 2,160.3 | 2,173.5 | 2,277.8 | 2,242.8 |
| Transportation | 1,470.5 | 1,577.8 | 1,617.4 | 1,626.9 | 1,653.9 | 1,713.0 | 1,762.7 | 1,780.9 |
| Industrial | 955.3 | 1,002.6 | 1,039.5 | 1,050.8 | 1,000.1 | 973.2 | 991.1 | 937.7 |
| Residential | 328.9 | 358.5 | 388.6 | 371.7 | 338.8 | 357.3 | 373.9 | 363.3 |
| Commercial | 221.4 | 226.9 | 236.4 | 237.1 | 219.5 | 221.7 | 234.3 | 235.9 |
| U.S. Territories | 33.7 | 44.0 | 40.1 | 42.8 | 47.9 | 50.2 | 52.3 | 54.4 |
| Iron and Steel Production | 85.4 | 74.4 | 68.3 | 71.9 | 67.4 | 64.4 | 65.8 | 59.1 |
| Cement Manufacture | 33.3 | 36.8 | 37.1 | 38.3 | 39.2 | 40.0 | 41.2 | 41.4 |
| Waste Combustion | 14.1 | 18.5 | 19.4 | 21.2 | 22.5 | 23.9 | 25.4 | 26.9 |
| Ammonia Manufacture & Urea Application | 19.3 | 20.5 | 20.3 | 20.7 | 21.9 | 20.6 | 19.6 | 16.6 |
| Lime Manufacture | 11.2 | 12.8 | 13.5 | 13.7 | 13.9 | 13.5 | 13.3 | 12.9 |
| Natural Gas Flaring | 5.5 | 8.7 | 8.2 | 7.6 | 6.3 | 6.7 | 5.5 | 5.2 |
| Limestone and Dolomite Use | 5.5 | 7.0 | 7.6 | 7.1 | 7.3 | 7.7 | 5.8 | 5.3 |
| Aluminum Production | 6.3 | 5.3 | 5.6 | 5.6 | 5.8 | 5.9 | 5.4 | 4.1 |
| Soda Ash Manufacture and Consumption | 4.1 | 4.3 | 4.2 | 4.4 | 4.3 | 4.2 | 4.2 | 4.1 |
| Titanium Dioxide Production | 1.3 | 1.7 | 1.7 | 1.8 | 1.8 | 1.9 | 1.9 | 1.9 |
| Carbon Dioxide Consumption | 0.9 | 1.1 | 1.1 | 1.2 | 1.2 | 1.2 | 1.2 | 1.3 |
| Ferroalloys | 2.0 | 1.9 | 2.0 | 2.0 | 2.0 | 2.0 | 1.7 | 1.3 |
| Land-Use Change and Forestry (Sink) ^a | (1,072.8) | (1,064.2) | (1,061.0) | (840.6) | (830.5) | (841.1) | (834.6) | (838.1) |
| International Bunker Fuels ^b | 113.9 | 101.0 | 102.3 | 109.9 | 112.9 | 105.3 | 99.3 | 97.3 |
| Total | 5,003.7 | 5,334.4 | 5,514.8 | 5,595.4 | 5,614.2 | 5,680.7 | 5,883.1 | 5,794.8 |
| Net Emissions (Sources and Sinks) | 3,930.9 | 4,270.3 | 4,453.8 | 4,754.7 | 4,783.7 | 4,839.6 | 5,048.5 | 4,956.7 |

^a Sinks are only included in net emissions total, and are based partially on projected activity data. Parentheses indicate negative values (or sequestration).

^b Emissions from International Bunker Fuels are not included in totals.

Note: Totals may not sum due to independent rounding.

Figure ES-10





Figure ES-11



Fossil Fuel Combustion (5,614.9 Tg CO₂ Eq.)

As fossil fuels are combusted, the carbon stored in them is emitted almost entirely as CO₂. The amount of carbon in fuels per unit of energy content varies significantly by fuel type. For example, coal contains the highest amount of carbon per unit of energy, while petroleum and natural gas have about 25 percent and 45 percent less carbon than coal, respectively. From 1990 through 2001, petroleum supplied the largest share of U.S. energy demands, accounting for an average of 39 percent of total energy consumption. Natural gas and coal followed in order of importance, accounting for an average of 24 and 23 percent of total energy consumption, respectively. Most petroleum was consumed in the transportation end-use sector, while the vast majority of coal was used by electric power generators, and natural gas was consumed largely in the industrial and residential end-use sectors.

Emissions of CO₂ from fossil fuel combustion increased at an average annual rate of 1.4 percent from 1990 to 2001. The fundamental factors influencing this trend include (1) a growing domestic economy over the last 11 years, and (2) significant growth in emissions from transportation activities and electricity generation. Between 1990 and 2001, CO₂ emissions from fossil fuel combustion increased from 4,814.8 Tg CO₂ Eq. to 5,614.9 Tg CO₂ Eq.—a 17 percent total increase over the twelve-year period.

The four major end-use sectors contributing to CO₂ emissions from fossil fuel combustion are industrial, transportation, residential, and commercial. Electricity generation also emits CO₂, although these emissions are produced as they consume fossil fuel to provide electricity to one of the four end-use sectors. For the discussion below, electricity generation emissions have been distributed to each end-use sector on the basis of each sector's share of aggregate electricity consumption. This method of distributing emissions assumes that each end-use sector consumes electricity that is generated from the national average mix of fuels according to their carbon intensity. In reality, sources of electricity vary widely in carbon intensity. By assuming the same carbon intensity for each end-use sector's electricity consumption, for example, emissions attributed to the residential end-use sector may be underestimated, while emissions attributed to the industrial end-use sector may be overestimated. Emissions from electricity generation are also addressed separately after the end-use sectors have been discussed.

Note that emissions from U.S. territories are calculated separately due to a lack of specific consumption data for the individual end-use sectors. Table ES-9, Figure ES-12, and Figure ES-13 summarize CO₂ emissions from fossil fuel combustion by end-use sector.

Table ES-9: CO₂ Emissions from Fossil Fuel Combustion by End-Use Sector (Tg CO₂ Eq.)

| End-Use Sector | 1990 | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 |
|-------------------------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|
| Industrial | 1,632.1 | 1,710.3 | 1,767.4 | 1,796.8 | 1,764.6 | 1,744.8 | 1,779.5 | 1,684.5 |
| Combustion | 955.3 | 1,002.6 | 1,039.5 | 1,050.8 | 1,000.1 | 973.2 | 991.1 | 937.7 |
| Electricity | 676.8 | 707.7 | 727.9 | 746.0 | 764.5 | 771.7 | 788.4 | 746.8 |
| Transportation | 1,473.5 | 1,580.9 | 1,620.4 | 1,630.0 | 1,657.0 | 1,716.2 | 1,766.1 | 1,784.4 |
| Combustion | 1,470.5 | 1,577.8 | 1,617.4 | 1,626.9 | 1,653.9 | 1,713.0 | 1,762.7 | 1,780.9 |
| Electricity | 3.0 | 3.0 | 3.0 | 3.1 | 3.1 | 3.2 | 3.4 | 3.6 |
| Residential | 918.8 | 996.4 | 1,056.6 | 1,048.0 | 1,051.6 | 1,069.4 | 1,127.3 | 1,111.1 |
| Combustion | 328.9 | 358.5 | 388.6 | 371.7 | 338.8 | 357.3 | 373.9 | 363.3 |
| Electricity | 589.9 | 637.8 | 668.1 | 676.4 | 712.8 | 712.1 | 753.5 | 747.8 |
| Commercial | 756.6 | 810.0 | 841.2 | 882.5 | 899.4 | 908.2 | 966.9 | 980.5 |
| Combustion | 221.4 | 226.9 | 236.4 | 237.1 | 219.5 | 221.7 | 234.3 | 235.9 |
| Electricity | 535.2 | 583.1 | 604.8 | 645.4 | 679.9 | 686.5 | 732.6 | 744.6 |
| U.S. Territories | 33.7 | 44.0 | 40.1 | 42.8 | 47.9 | 50.2 | 52.3 | 54.4 |
| Total | 4,814.8 | 5,141.5 | 5,325.8 | 5,400.0 | 5,420.5 | 5,488.8 | 5,692.2 | 5,614.9 |
| Electricity Generation | 1,805.0 | 1,931.8 | 2,003.9 | 2,070.8 | 2,160.3 | 2,173.5 | 2,277.8 | 2,242.8 |

Note: Totals may not sum due to independent rounding. Emissions from fossil fuel combustion by electricity generation are allocated based on aggregate national electricity consumption by each end-use sector.

Figure ES-12





Figure ES-13



Industrial End-Use Sector. Industrial CO₂ emissions, resulting both directly from the combustion of fossil fuels and indirectly from the generation of electricity that is consumed by industry, accounted for 30 percent of CO₂ from fossil fuel combustion in 2001. About half of these emissions resulted from direct fossil fuel combustion to produce steam and/or heat for industrial processes. The other half of the emissions resulted from consuming electricity for motors, electric furnaces, ovens, lighting, and other applications.

Transportation End-Use Sector. Transportation activities (excluding international bunker fuels) accounted for 32 percent of CO₂ emissions from fossil fuel combustion in 2001.¹² Virtually all of the energy consumed in this end-use sector came from petroleum products. Just over half of the emissions resulted from gasoline consumption for personal vehicle use. The remaining emissions came from other transportation activities, including the combustion of diesel fuel in heavy-duty vehicles and jet fuel in aircraft.

Residential and Commercial End-Use Sectors. The residential and commercial end-use sectors accounted for 20 and 17 percent, respectively, of CO₂ emissions from fossil fuel combustion in 2001. Both sectors relied heavily on electricity for meeting energy demands, with 67 and 76 percent, respectively, of their emissions attributable to electricity consumption for lighting, heating, cooling, and

operating appliances. The remaining emissions were due to the consumption of natural gas and petroleum for heating and cooking.

Electricity Generation. The United States relies on electricity to meet a significant portion of its energy demands, especially for lighting, electric motors, heating, and air conditioning. Electricity generators consumed 34 percent of U.S. energy from fossil fuels and emitted 40 percent of the CO₂ from fossil fuel combustion in 2001. The type of fuel combusted by electricity generators has a significant effect on their emissions. For example, some electricity is generated with low CO₂ emitting energy technologies, particularly non-fossil options such as nuclear, hydroelectric, or geothermal energy. However, electricity generators rely on coal for over half of their total energy requirements and accounted for 93 percent of all coal consumed for energy in the United States in 2001. Consequently, changes in electricity demand have a significant impact on coal consumption and associated CO₂ emissions.

Waste Combustion (26.9 Tg CO₂ Eq.)

The burning of garbage and non-hazardous solids, referred to as municipal solid waste (MSW), as well as the burning of hazardous waste, is usually performed to recover energy from the waste materials. Carbon dioxide emissions arise from the organic (i.e., carbon) materials found in these

¹² If emissions from international bunker fuels are included, the transportation end-use sector accounted for 33 percent of U.S. emissions from fossil fuel combustion in 2001.

wastes. Within MSW, many products contain carbon of biogenic origin, and the CO₂ emissions from their combustion are accounted for under the Land-Use Change and Forestry chapter. Several components of MSW, such as plastics, synthetic rubber, synthetic fibers, and carbon black, are of fossil fuel origin, and are included as sources of CO₂ emissions.

Natural Gas Flaring (5.2 Tg CO₂ Eq.)

Flaring of natural gas from oil wells releases CO₂. Natural gas is flared to relieve rising pressure or to dispose of small quantities of gas that are not commercially marketable. In 2001, flaring accounted for approximately 0.1 percent of U.S. CO₂ emissions.

Biomass Combustion (183.7 Tg CO₂ Eq.)

Biomass refers to organically-based carbon fuels (as opposed to fossil-based). Biomass in the form of fuel wood and wood waste was used primarily in the industrial sector, while the transportation sector was the predominant user of biomass-based fuels, such as ethanol from corn and woody crops.

Although these fuels do emit CO₂, in the long run the CO₂ emitted from biomass consumption does not increase atmospheric CO₂ concentrations if the biogenic carbon emitted is offset by the growth of new biomass. For example, fuel wood burned one year but re-grown the next only recycles carbon, rather than creating a net increase in total atmospheric carbon. Net carbon fluxes from changes in biogenic carbon reservoirs in wooded or croplands are accounted for in the estimates for Land-Use Change and Forestry.

The industrial sector accounted for 69 percent of gross CO₂ emissions from biomass combustion, and the residential sector accounted for 18 percent. Ethanol consumption by the transportation sector accounted for 6 percent.

Industrial Processes

Emissions are produced as a by-product of many non-energy-related activities. For example, industrial processes can chemically transform raw materials, which often release waste gases such as CO₂. The major production processes that emit CO₂ include iron and steel production, cement manufacture, lime manufacture, limestone and dolomite use, soda ash manufacture and consumption, CO₂ consumption, titanium dioxide production, ferroalloy production, and ammonia manufacturing and urea application. Carbon

dioxide emissions from these sources were approximately 148.0 Tg CO₂ Eq. in 2001, accounting for about 3 percent of total CO₂ emissions.

Iron and Steel Production (59.1 Tg CO₂ Eq.)

Iron is produced by first reducing iron oxide (i.e., iron ore) with metallurgical coke in a blast furnace to produce pig iron. Carbon dioxide is produced as the metallurgical coke used in the blast furnace process is oxidized. Steel, which contains less than 2 percent carbon by weight, is produced from pig iron in a variety of specialized steel making furnaces. The majority of CO₂ emissions from the iron and steel processes come from the use of coke in the production of pig iron, with smaller amounts evolving from the removal of carbon from pig iron used to produce steel.

Cement Manufacture (41.4 Tg CO₂ Eq.)

Clinker is an intermediate product in the formation of finished Portland and masonry cement. Heating calcium carbonate (CaCO₃) in a cement kiln forms lime and CO₂. The lime combines with other materials to produce clinker, and the CO₂ is released into the atmosphere.

Ammonia Manufacture and Urea Application (16.6 Tg CO₂ Eq.)

In the United States, roughly 98 percent of synthetic ammonia is produced by catalytic steam reforming of natural gas, and the remainder is produced using naphtha (i.e., a petroleum fraction) or the electrolysis of brine at chlorine plants (EPA 1997). The two fossil fuel-based reactions produce carbon monoxide and hydrogen gas. This carbon monoxide is transformed into CO₂ in the presence of a catalyst. The CO₂ is generally released into the atmosphere, but some of the CO₂, together with ammonia, is used as a raw material in the production of urea [CO(NH₂)₂], which is a type of nitrogenous fertilizer. The carbon in the urea that is produced and assumed to be subsequently applied to agricultural land as a nitrogenous fertilizer is ultimately released into the environment as CO₂.

Lime Manufacture (12.9 Tg CO₂ Eq.)

Lime is used in steel making, construction, flue gas desulfurization, and water and sewage treatment. It is manufactured by heating limestone (mostly calcium carbonate, CaCO₃) in a kiln, creating quicklime (calcium oxide, CaO) and CO₂, which is normally emitted to the atmosphere.

Limestone and Dolomite Use (5.3 Tg CO₂ Eq.)

Limestone (CaCO₃) and dolomite (CaMg(CO₃)) are basic raw materials used in a wide variety of industries, including construction, agriculture, chemical, and metallurgy. For example, limestone can be used as a purifier in refining metals. In the case of iron ore, limestone heated in a blast furnace reacts with impurities in the iron ore and fuels, generating CO₂ as a by-product. Limestone is also used in flue gas desulfurization systems to remove sulfur dioxide from the exhaust gases.

Aluminum Production (4.1 Tg CO₂ Eq.)

Carbon dioxide is emitted when alumina (aluminum oxide, Al₂O₃) is reduced to aluminum. The reduction of the alumina occurs through electrolysis in a molten bath of natural or synthetic cryolite. The reduction cells contain a carbon lining that serves as the cathode. Carbon is also contained in the anode, which can be a carbon mass of paste, coke briquettes, or prebaked carbon blocks from petroleum coke. During reduction, some of this carbon is oxidized and released to the atmosphere as CO₂.

Soda Ash Manufacture and Consumption (4.1 Tg CO₂ Eq.)

Commercial soda ash (sodium carbonate, Na₂CO₃) is used in many consumer products, such as glass, soap and detergents, paper, textiles, and food. During the manufacturing of soda ash, some natural sources of sodium carbonate are heated and transformed into a crude soda ash, in which CO₂ is generated as a by-product. In addition, CO₂ is often released when the soda ash is consumed.

Titanium Dioxide Production (1.9 Tg CO₂ Eq.)

Titanium dioxide (TiO₂) is a metal oxide manufactured from titanium ore, and is principally used as a pigment. It is used in white paint and as a pigment in the manufacture of white paper, foods, and other products. Two processes, the chloride process and the sulfate process, are used for making TiO₂. Carbon dioxide is emitted from the chloride process, which uses petroleum coke and chlorine as raw materials.

Ferroalloy Production (1.3 Tg CO₂ Eq.)

Carbon dioxide is emitted from the production of several ferroalloys through the use of metallurgical coke as a raw material. Ferroalloys are composites of iron and other elements, often including silicon, manganese, and chromium. When incorporated in alloy steels, ferroalloys alter the material properties of the steel.

Carbon Dioxide Consumption (1.3 Tg CO₂ Eq.)

Many segments of the economy consume CO₂, including food processing, beverage manufacturing, chemical processing, and a host of industrial and other miscellaneous applications. Carbon dioxide may be produced as a by-product from the production of certain chemicals (e.g., ammonia), from select natural gas wells, or by separating it from crude oil and natural gas. For the most part, the CO₂ used in these applications is eventually released to the atmosphere.

Land-Use Change and Forestry

When humans alter the terrestrial biosphere through land use, changes in land-use, and forest management practices, they also alter the natural carbon fluxes between biomass, soils, and the atmosphere. Forest management practices, tree planting in urban areas, the management of agricultural soils, and landfilling of yard trimmings have resulted in a net uptake (sequestration) of carbon in the United States, which offset about 14 percent of total U.S. gross CO₂ emissions in 2001. Forests (including vegetation, soils, and harvested wood) accounted for approximately 91 percent of total 2001 sequestration, urban trees accounted for 7 percent, agricultural soils (including mineral and organic soils and the application of lime) accounted for 2 percent, and landfilled yard trimmings accounted for 1 percent of the total sequestration in 2001. The net forest sequestration is a result of net forest growth and increasing forest area, as well as a net accumulation of carbon stocks in harvested wood pools. The net sequestration in urban forests is a result of net tree growth in these areas. In agricultural soils, mineral soils account for a net carbon sink that is approximately one and a third time larger than the sum of emissions from organic soils and liming. The mineral soil carbon sequestration is largely due to conversion of cropland to permanent pastures and hay production, a reduction in summer fallow areas in semi-arid areas, an increase in the adoption of conservation tillage practices, and an increase in the amounts of organic fertilizers (i.e., manure and sewage sludge) applied to agriculture lands. The landfilled yard trimmings net sequestration is due to the long-term accumulation of yard trimming carbon in landfills.

Methane Emissions


According to the IPCC, CH₄ is more than 20 times as effective as CO₂ at trapping heat in the atmosphere. Over the last two hundred and fifty years, the concentration of CH₄ in the atmosphere increased by 150 percent (IPCC 2001). Experts believe that over half of this atmospheric increase was due to emissions from anthropogenic sources, such as landfills, natural gas and petroleum systems, agricultural activities, coal mining, stationary and mobile combustion, wastewater treatment, and certain industrial processes (see Figure ES-14 and Table ES-10).

Landfills (202.9 Tg CO₂ Eq.)

Landfills are the largest anthropogenic source of CH₄ emissions in the United States. In an environment where the oxygen content is low or zero, anaerobic bacteria can decompose organic materials, such as yard waste, household waste, food waste, and paper, resulting in the generation of CH₄ and biogenic CO₂. Site-specific factors, such as waste composition, moisture, and landfill size, influence the level of methane generation.

Methane emissions from U.S. landfills have decreased by 4 percent since 1990. The generally declining emission estimates are a result of two offsetting trends: (1) the amount of municipal solid waste in landfills contributing to CH₄ emissions has increased, thereby increasing the potential

Figure ES-14



for emissions; and (2) the amount of landfill gas collected and combusted by landfill operators has also increased, thereby reducing emissions. Additionally, a regulation promulgated in March 1996 requires the largest U.S. landfills to begin collecting and combusting their landfill gas to reduce emissions of NMVOCs.

Natural Gas and Petroleum Systems (138.6 Tg CO₂ Eq.)

Methane is the major component of natural gas. Fugitive emissions of CH₄ occur throughout the production, processing, transmission, and distribution of natural gas. Because natural gas is often found in conjunction with

Table ES-10: U.S. Sources of Methane Emissions (Tg CO₂ Eq.)

| Source | 1990 | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 |
|--|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|
| Landfills | 212.1 | 216.1 | 212.1 | 207.5 | 202.4 | 203.7 | 205.8 | 202.9 |
| Natural Gas Systems | 122.0 | 127.2 | 127.4 | 126.0 | 124.0 | 120.3 | 121.2 | 117.3 |
| Enteric Fermentation | 117.9 | 123.0 | 120.5 | 118.3 | 116.7 | 116.6 | 115.7 | 114.8 |
| Coal Mining | 87.1 | 73.5 | 68.4 | 68.1 | 67.9 | 63.7 | 60.9 | 60.7 |
| Manure Management | 31.3 | 36.2 | 34.9 | 36.6 | 39.0 | 38.9 | 38.2 | 38.9 |
| Wastewater Treatment | 24.1 | 26.6 | 26.8 | 27.3 | 27.7 | 28.2 | 28.3 | 28.3 |
| Petroleum Systems | 27.5 | 24.2 | 23.9 | 23.6 | 22.9 | 21.6 | 21.2 | 21.2 |
| Rice Cultivation | 7.1 | 7.6 | 7.0 | 7.5 | 7.9 | 8.3 | 7.5 | 7.6 |
| Stationary Sources | 8.1 | 8.5 | 8.7 | 7.5 | 7.2 | 7.4 | 7.6 | 7.4 |
| Mobile Sources | 5.0 | 4.9 | 4.8 | 4.7 | 4.6 | 4.5 | 4.4 | 4.3 |
| Petrochemical Production | 1.2 | 1.5 | 1.6 | 1.6 | 1.6 | 1.7 | 1.7 | 1.5 |
| Field Burning of Agricultural Residues | 0.7 | 0.7 | 0.7 | 0.8 | 0.8 | 0.8 | 0.8 | 0.8 |
| Silicon Carbide Production | + | + | + | + | + | + | + | + |
| International Bunker Fuels* | 0.2 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 |
| Total* | 644.0 | 650.0 | 636.8 | 629.5 | 622.7 | 615.5 | 613.4 | 605.9 |

+ Does not exceed 0.05 Tg CO₂ Eq.

* Emissions from International Bunker Fuels are not included in totals.

Note: Totals may not sum due to independent rounding.

petroleum deposits, leakage from petroleum systems is also a source of emissions. Emissions vary greatly from facility to facility and are largely a function of operation and maintenance procedures and equipment conditions. In 2001, CH₄ emissions from U.S. natural gas systems were accounted for approximately 19 percent of U.S. CH₄ emissions.

Petroleum is often found in the same geological structures as natural gas, and the two are often retrieved together. Crude oil is saturated with many lighter hydrocarbons, including methane. When the oil is brought to the surface and processed, many of the dissolved lighter hydrocarbons (as well as water) are removed through a series of high-pressure and low-pressure separators. The remaining hydrocarbons in the oil are emitted at various points along the system. Methane emissions from the components of petroleum systems generally occur as a result of system leaks, disruptions, and routine maintenance. In 2001, emissions from petroleum systems were just under 4 percent of U.S. CH₄ emissions.

Coal Mining (60.7 Tg CO₂ Eq.)

Produced millions of years ago during the formation of coal, CH₄ trapped within coal seams and surrounding rock strata is released when the coal is mined. The quantity of CH₄ released to the atmosphere during coal mining operations depends primarily upon the type of coal and the method and rate of mining.

Methane from surface mines is emitted directly to the atmosphere as the rock strata overlying the coal seam are removed. Because CH₄ in underground mines is explosive at concentrations of 5 to 15 percent in air, most active underground mines are required to vent this methane, typically to the atmosphere. At some mines, CH₄-recovery systems may supplement these ventilation systems. Recovery of CH₄ in the United States has increased in recent years. During 2001, coal mining activities emitted 10 percent of U.S. CH₄ emissions. From 1990 to 2001, emissions from this source decreased by 30 percent due to increased use of the CH₄ collected by mine degasification systems and a general shift toward surface mining.

Stationary and Mobile Combustion (11.7 Tg CO₂ Eq.)

In 2001, stationary and mobile combustion were responsible for CH₄ emissions of 7.4 and 4.3 Tg CO₂ Eq., respectively. The majority of CH₄ emissions from stationary combustion resulted from the burning of wood in the residential end-use sector. The combustion of gasoline in highway vehicles was responsible for the majority of the CH₄ emitted from mobile combustion.

Petrochemical and Silicon Carbide Production (1.5 Tg CO₂ Eq.)

Small amounts of CH₄ are released during the production of five petrochemicals: carbon black, ethylene, ethylene dichloride, styrene, and methanol. These production processes resulted in emissions of 1.5 Tg CO₂ Eq. in 2001. Methane is also emitted from the production of silicon carbide, a material used as an industrial abrasive. In 2001, silicon carbide production resulted in emissions of less than 0.1 Tg CO₂ Eq.

Enteric Fermentation (114.8 Tg CO₂ Eq.)

During animal digestion, CH₄ is produced through the process of enteric fermentation, in which microbes residing in animal digestive systems break down food. Ruminants, which include cattle, buffalo, sheep, and goats, have the highest CH₄ emissions among all animal types because they have a rumen, or large fore-stomach, in which CH₄-producing fermentation occurs. Non-ruminant domestic animals, such as pigs and horses, have much lower CH₄ emissions. In 2001, enteric fermentation was the source of about 19 percent of U.S. CH₄ emissions, and more than 70 percent of the CH₄ emissions from agriculture. From 1990 to 2001, emissions from this source decreased by 3 percent. Emissions from enteric fermentation have been generally decreasing since 1995, primarily due to declining dairy cow and beef cattle populations.

Manure Management (38.9 Tg CO₂ Eq.)

The decomposition of organic animal waste in an anaerobic environment produces CH₄. The most important factor affecting the amount of CH₄ produced is how the manure is managed, because certain types of storage and treatment systems promote an oxygen-free environment. In particular, liquid systems tend to encourage anaerobic

conditions and produce significant quantities of CH₄, whereas solid waste management approaches produce little or no CH₄. Higher temperatures and moist climatic conditions also promote CH₄ production.

Emissions from manure management were about 6 percent of U.S. CH₄ emissions in 2001 and 24 percent of the CH₄ emissions from agriculture. From 1990 to 2001, emissions from this source increased by 24 percent. The bulk of this increase was from swine and dairy cow manure, and is attributed to the shift in the composition of the swine and dairy industries towards larger facilities. Larger swine and dairy farms tend to use liquid management systems.

Rice Cultivation (7.6 Tg CO₂ Eq.)

Most of the world's rice, and all of the rice in the United States, is grown on flooded fields. When fields are flooded, anaerobic conditions develop and the organic matter in the soil decomposes, releasing CH₄ to the atmosphere, primarily through the rice plants. In 2001, rice cultivation was the source of 1 percent of U.S. CH₄ emissions, and about 5 percent of U.S. CH₄ emissions from agriculture. Emission estimates from this source have increased about 7 percent since 1990 due to an increase in the area harvested.

Field Burning of Agricultural Residues (0.8 Tg CO₂ Eq.)

Burning crop residue releases a number of greenhouse gases, including CH₄. Because field burning is not a common debris clearing method used in the United States, it was responsible for only 0.1 percent of U.S. CH₄ emissions in 2001.

Wastewater Treatment (28.3 Tg CO₂ Eq.)

Wastewater from domestic sources (i.e., municipal sewage) and industrial sources is treated to remove soluble organic matter, suspended solids, pathogenic organisms and chemical contaminants. Soluble organic matter is generally removed using biological processes in which microorganisms consume the organic matter for maintenance and growth. Microorganisms can biodegrade soluble organic material in wastewater under aerobic or anaerobic conditions, with the latter condition producing CH₄. During collection and

treatment, wastewater may be accidentally or deliberately managed under anaerobic conditions. In addition, the sludge may be further biodegraded under aerobic or anaerobic conditions. Untreated wastewater may also produce CH₄ if contained under anaerobic conditions.

Nitrous Oxide Emissions

Nitrous oxide (N₂O) is produced naturally from biological sources in soil and water, and anthropogenically by a variety of agricultural, energy-related, industrial, and waste management activities. While total N₂O emissions are much lower than CO₂ emissions, N₂O is approximately 300 times more powerful than CO₂ at trapping heat in the atmosphere. Since 1750, the atmospheric concentration of N₂O has risen by approximately 16 percent (IPCC 2001). The main anthropogenic activities producing N₂O in the United States are agricultural soil management, fuel combustion in motor vehicles, manure management, nitric acid production, human sewage, and stationary fuel combustion (see Figure ES-15 and Table ES-11).

Figure ES-15




Table ES-11: U.S. Sources of Nitrous Oxide Emissions (Tg CO₂ Eq.)

| Source | 1990 | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 |
|--|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|
| Agricultural Soil Management | 267.5 | 284.1 | 293.2 | 298.2 | 299.2 | 297.0 | 294.6 | 294.3 |
| Mobile Sources | 50.6 | 60.9 | 60.7 | 60.3 | 59.7 | 58.8 | 57.5 | 54.8 |
| Manure Management | 16.2 | 16.6 | 17.0 | 17.3 | 17.3 | 17.4 | 17.9 | 18.0 |
| Nitric Acid | 17.8 | 19.9 | 20.7 | 21.2 | 20.9 | 20.1 | 19.1 | 17.6 |
| Human Sewage | 12.7 | 13.9 | 14.1 | 14.4 | 14.6 | 15.1 | 15.1 | 15.3 |
| Stationary Combustion | 12.5 | 13.2 | 13.8 | 13.7 | 13.7 | 13.7 | 14.3 | 14.2 |
| Adipic Acid | 15.2 | 17.2 | 17.0 | 10.3 | 6.0 | 5.5 | 6.0 | 4.9 |
| N ₂ O Product Usage | 4.3 | 4.5 | 4.5 | 4.8 | 4.8 | 4.8 | 4.8 | 4.8 |
| Field Burning of Agricultural Residues | 0.4 | 0.4 | 0.4 | 0.4 | 0.5 | 0.4 | 0.5 | 0.5 |
| Waste Combustion | 0.3 | 0.3 | 0.3 | 0.3 | 0.2 | 0.2 | 0.2 | 0.2 |
| International Bunker Fuels* | 1.0 | 0.9 | 0.9 | 1.0 | 1.0 | 0.9 | 0.9 | 0.9 |
| Total* | 397.6 | 430.9 | 441.7 | 440.9 | 436.8 | 433.0 | 429.9 | 424.6 |

* Emissions from International Bunker Fuels are not included in totals.
Note: Totals may not sum due to independent rounding.

Agricultural Soil Management (294.3 Tg CO₂ Eq.)

Nitrous oxide is produced naturally in soils through microbial processes of nitrification and denitrification. A number of anthropogenic activities add to the amount of nitrogen available to be emitted as N₂O by microbial processes. These activities may add nitrogen to soils either directly or indirectly. Direct additions occur through the application of synthetic and organic fertilizers; production of nitrogen-fixing crops and forages; the application of livestock manure, crop residues, and sewage sludge; cultivation of high-organic-content soils; and direct excretion by animals onto soil. Indirect additions result from volatilization and subsequent atmospheric deposition, and from leaching and surface run-off of some of the nitrogen applied to or deposited on soils as fertilizer, livestock manure, and sewage sludge.

In 2001, agricultural soil management accounted for 69 percent of U.S. N₂O emissions. From 1990 to 2001, emissions from this source increased by 10 percent as fertilizer consumption, manure production, and production of nitrogen-fixing and other crops rose.

Stationary and Mobile Combustion (68.9 Tg CO₂ Eq.)

Nitrous oxide is a product of the reaction that occurs between nitrogen and oxygen during fuel combustion. Both mobile and stationary combustion emit N₂O, and the quantity emitted varies according to the type of fuel, technology, and pollution control device used, as well as maintenance and operating practices. For example, some types of catalytic converters installed to reduce motor vehicle pollution can promote the formation of N₂O.

In 2001, N₂O emissions from mobile combustion were 13 percent of U.S. N₂O emissions, while stationary combustion accounted for 3 percent. From 1990 to 2001, combined N₂O emissions from stationary and mobile combustion increased by 9 percent, primarily due to increased rates of N₂O generation in highway vehicles.

Adipic Acid Production (4.9 Tg CO₂ Eq.)

Most adipic acid produced in the United States is used to manufacture nylon 6,6. Adipic acid is also used to produce some low-temperature lubricants and to add a “tangy” flavor to foods. Nitrous oxide is emitted as a by-product of the chemical synthesis of adipic acid.

In 2001, U.S. adipic acid plants emitted 1 percent of U.S. N₂O emissions. Even though adipic acid production has increased, by 1998 all three major adipic acid plants in the United States had voluntarily implemented N₂O abatement technology. As a result, emissions have decreased by 68 percent since 1990.

Nitric Acid Production (17.6 Tg CO₂ Eq.)

Nitric acid production is another industrial source of N₂O emissions. Used primarily to make synthetic commercial fertilizer, this raw material is also a major component in the production of adipic acid and explosives.

Virtually all of the nitric acid manufactured in the United States is produced by the oxidation of ammonia, during which N₂O is formed and emitted to the atmosphere. In 2001, N₂O emissions from nitric acid production accounted for 4 percent of U.S. N₂O emissions. From 1990 to 2001, emissions from this source category decreased by 1 percent with the trend in the time series closely tracking the changes in production.

N₂O from Product Usage (4.8 Tg CO₂ Eq.)

Nitrous oxide is used in carrier gases with oxygen to administer more potent inhalation anesthetics for general anesthesia and as an anesthetic in various dental and veterinary applications. As such, it is used to treat short-term pain, for sedation in minor elective surgeries and as an induction anesthetic. The second main use of N₂O is as a propellant in pressure and aerosol products, the largest application being pressure-packaged whipped cream.

Manure Management (18.0 Tg CO₂ Eq.)

Nitrous oxide is produced as part of microbial nitrification and denitrification processes in managed and unmanaged manure, the latter of which is addressed under agricultural soil management. Total N₂O emissions from managed manure systems in 2001 accounted for 4 percent of U.S. N₂O emissions. From 1990 to 2001, emissions from this source category increased by 11 percent, as poultry and swine populations have increased.

Field Burning of Agricultural Residues (0.5 Tg CO₂ Eq.)

Large quantities of agricultural crop residues are produced by farming activities, some of which is disposed by burning in the field. Field burning of crop residues is a source of N₂O, which is released during combustion. Because field burning is not a common method of agricultural residue disposal in the United States, emissions from this source are minor.

Human Sewage (Domestic Wastewater) (15.3 Tg CO₂ Eq.)

Domestic human sewage is usually mixed with other household wastewater, which includes shower drains, sink drains, washing machine effluent, etc., and transported by a collection system to either a direct discharge, an on-site or decentralized or centralized wastewater treatment system. After processing, treated effluent may be discharged to a receiving water environment (e.g., river, lake, estuary, etc.), applied to soils, or disposed of below the surface. Nitrous oxide may be generated during both nitrification and denitrification of the nitrogen present, usually in the form of urea, ammonia, and proteins. Emissions of N₂O from treated human sewage discharged into aquatic environments were estimated to be 15.3 Tg CO₂ Eq. in 2001.

Waste Combustion (0.2 Tg CO₂ Eq.)

Combustion is used to manage about 7 to 17 percent of the municipal solid wastes (MSW) generated in the United States. Almost all combustion of MSW in the United States occurs at waste-to-energy facilities where energy is recovered. Most of the organic materials in MSW are of biogenic origin (e.g., paper, yard trimmings), with some components, such as plastics, synthetic rubber, and synthetic fibers, of fossil origin, which together accounted for emissions of 0.2 Tg CO₂ Eq. in 2001.

HFC, PFC, and SF₆ Emissions

Hydrofluorocarbons (HFCs) and perfluorocarbons (PFCs) are families of synthetic chemicals that are being used as alternatives to the ozone depleting substances (ODSs), which are being phased out under the *Montreal Protocol* and Clean Air Act Amendments of 1990. HFCs and PFCs do not deplete the stratospheric ozone layer, and are therefore acceptable alternatives under the *Montreal Protocol*.

These compounds, however, along with sulfur hexafluoride (SF₆), are potent greenhouse gases. In addition to having high global warming potentials, SF₆ and PFCs have extremely long atmospheric lifetimes, resulting in their essentially irreversible accumulation in the atmosphere once emitted. Sulfur hexafluoride is the most potent greenhouse gas the IPCC has evaluated.

Other emissive sources of these gases include aluminum production, HCFC-22 production, semiconductor manufacturing, electrical transmission and distribution systems, and magnesium production and processing. Figure ES-16 and Table ES-12 present emission estimates for HFCs, PFCs, and SF₆, which totaled 111.0 Tg CO₂ Eq. in 2001.

Substitution of Ozone Depleting Substances (63.7 Tg CO₂ Eq.)

The use and subsequent emissions of HFCs and PFCs as substitutes for ozone depleting substances (ODSs) have increased from small amounts in 1990 to account for 57 percent of aggregate HFC, PFC, and SF₆ emissions. This increase was in large part the result of efforts to phase-out chlorofluorocarbons (CFCs) and other ODSs in the United States, especially the introduction of HFC-134a as a CFC substitute in refrigeration and air-conditioning applications.

Table ES-12: Emissions of HFCs, PFCs, and SF₆ (Tg CO₂ Eq.)

| Source | 1990 | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 |
|--|-------------|-------------|--------------|--------------|--------------|--------------|--------------|--------------|
| Substitution of Ozone Depleting Substances | 0.9 | 21.7 | 30.4 | 37.7 | 44.5 | 50.9 | 57.3 | 63.7 |
| HCFC-22 Production | 35.0 | 27.0 | 31.1 | 30.0 | 40.2 | 30.4 | 29.8 | 19.8 |
| Electrical Transmission and Distribution | 32.1 | 27.5 | 27.7 | 25.2 | 20.9 | 16.4 | 15.4 | 15.3 |
| Semiconductor Manufacture | 2.9 | 5.9 | 5.4 | 6.5 | 7.3 | 7.7 | 7.4 | 5.5 |
| Aluminum Production | 18.1 | 11.8 | 12.5 | 11.0 | 9.0 | 8.9 | 7.9 | 4.1 |
| Magnesium Production and Processing | 5.4 | 5.6 | 6.5 | 6.3 | 5.8 | 6.0 | 3.2 | 2.5 |
| Total | 94.4 | 99.5 | 113.6 | 116.8 | 127.6 | 120.3 | 121.0 | 111.0 |

Note: Totals may not sum due to independent rounding.

In the short term, this trend is expected to continue, and will likely accelerate in the next decade as hydrochlorofluoro-carbons (HCFCs), which are interim substitutes in many applications, are themselves phased-out under the provisions of the Copenhagen Amendments to the *Montreal Protocol*. Improvements in the technologies associated with the use of these gases and the introduction of alternative gases and technologies, however, may help to offset this anticipated increase in emissions.

Aluminum Production (4.1 Tg CO₂ Eq.)

During the production of primary aluminum CF₄ and C₂F₆ are emitted as intermittent by-products of the smelting process. These PFCs are formed when fluorine from the cryolite bath combines with carbon from the electrolyte anode. Emissions from aluminum production have decreased by 77 percent between 1990 and 2001 due to emission reduction efforts by the industry and falling domestic aluminum production.

HCFC-22 Production (19.8 Tg CO₂ Eq.)

HFC-23 is a by-product of the production of HCFC-22. Emissions from this source have decreased by 43 percent since 1990. The HFC-23 emission rate (i.e., the amount of HFC-23 emitted per kilogram of HCFC-22 manufactured) has declined significantly since 1990, although production has been increasing.

Semiconductor Manufacturing (5.5 Tg CO₂ Eq.)


The semiconductor industry uses combinations of HFCs, PFCs, SF₆, and other gases for plasma etching and to clean chemical vapor deposition tools. Emissions from this source category have increased with the growth in the semiconductor industry and the rising intricacy of chip

designs. However, the growth rate in emissions has slowed since 1997, and emissions actually declined between 1999 and 2001. This later reduction is due to the implementation of PFC emission reduction methods, such as process optimization.

Electrical Transmission and Distribution Systems (15.3 Tg CO₂ Eq.)

The primary use of SF₆ is as a dielectric in electrical transmission and distribution systems. Fugitive emissions of SF₆ occur from leaks in and servicing of substations and circuit breakers, especially from older equipment. Estimated emissions from this source decreased by 52 percent since 1990, primarily due to higher SF₆ prices and industrial efforts to reduce emissions.

Figure ES-16



Magnesium Production (2.5 Tg CO₂ Eq.)

Sulfur hexafluoride is also used as a protective cover gas for the casting of molten magnesium. Emissions from primary magnesium production and magnesium casting have

decreased by 53 percent since 1990. Emissions have decreased since 1999, due to a decrease in the quantity of magnesium die cast and the closure of a U.S. primary magnesium production facility.

Box ES-3: Emissions of Ozone Depleting Substances

Manmade halogenated compounds were first emitted into the atmosphere in significant quantities during the 20th century. This family of man-made compounds includes CFCs, halons, methyl chloroform, carbon tetrachloride, methyl bromide, and hydrochlorofluorocarbons (HCFCs). These substances have a variety of industrial applications, including refrigeration, air conditioning, foam blowing, solvent cleaning, sterilization, fire extinguishing, agricultural fumigation and sterilization, coatings, paints, and aerosols.

Because these compounds have been shown to deplete stratospheric ozone, they are typically referred to as ozone depleting substances (ODSs). They are also potent greenhouse gases.

Recognizing the harmful effects of these compounds on the ozone layer, in 1987 many countries ratified the *Montreal Protocol on Substances that Deplete the Ozone Layer* to limit the production and importation of a number of CFCs and other halogenated compounds. The United States furthered its commitment to phase out ODSs by signing and ratifying the Copenhagen Amendments to the *Montreal Protocol* in 1992. Under these amendments, the United States committed to ending the production and importation of halons by 1994, and CFCs by 1996.

The UNFCCC reporting guidelines do not include reporting instructions for estimating emissions of ODSs because they are tracked under the *Montreal Protocol*. Nevertheless, estimates for several Class I and Class II ODSs are provided in Table ES-13 for informational purposes. Compounds are grouped by class according to their ozone depleting potential. Class I compounds are the primary ODSs; Class II compounds include partially halogenated chlorine compounds (i.e., HCFCs), some of which were developed as interim replacements for CFCs. Because these HCFC compounds are only partially halogenated, their hydrogen-carbon bonds are more vulnerable to oxidation in the troposphere and, therefore, pose only one-tenth to one-hundredth the threat to stratospheric ozone compared to CFCs.

It should be noted that the effects of these compounds on radiative forcing are not provided. Although many ODSs have relatively high direct GWPs, their indirect effects from the destruction of ozone—also a greenhouse gas—are believed to have negative radiative forcing effects, and therefore could significantly reduce the overall magnitude of their radiative forcing effects. Given the uncertainties surrounding the net effect of these gases, emissions are reported on an unweighted basis.

Table ES-13: Emissions of Ozone Depleting Substances (Gg)

| Compound | 1990 | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 |
|----------------------|-------|------|------|------|------|------|------|------|
| Class I | | | | | | | | |
| CFC-11 | 53.5 | 36.2 | 26.6 | 25.1 | 24.9 | 24.0 | 22.8 | 22.8 |
| CFC-12 | 112.6 | 51.8 | 35.5 | 23.1 | 21.0 | 14.0 | 17.2 | 21.3 |
| CFC-113 | 52.7 | 17.1 | + | + | + | + | + | + |
| CFC-114 | 4.7 | 1.6 | + | + | + | + | + | + |
| CFC-115 | 4.2 | 3.0 | 3.2 | 2.9 | 2.7 | 2.6 | 2.3 | 1.5 |
| Carbon Tetrachloride | 32.3 | 4.7 | + | + | + | + | + | + |
| Methyl Chloroform | 316.6 | 92.8 | + | + | + | + | + | + |
| Halon-1211 | 1.0 | 1.1 | 1.1 | 1.1 | 1.1 | 1.1 | 1.1 | 1.1 |
| Halon-1301 | 1.8 | 1.4 | 1.4 | 1.3 | 1.3 | 1.3 | 1.3 | 1.2 |
| Class II | | | | | | | | |
| HCFC-22 | 34.0 | 39.3 | 41.0 | 42.4 | 43.8 | 74.1 | 79.1 | 80.5 |
| HCFC-123 | + | 0.6 | 0.7 | 0.8 | 0.9 | 1.0 | 1.1 | 1.2 |
| HCFC-124 | + | 5.6 | 5.9 | 6.2 | 6.4 | 6.5 | 6.5 | 6.5 |
| HCFC-141b | 1.3 | 9.9 | 9.9 | 8.8 | 9.7 | 10.9 | 10.9 | 10.7 |
| HCFC-142b | 0.8 | 3.6 | 4.0 | 4.3 | 4.7 | 5.0 | 5.4 | 5.8 |
| HCFC-225ca/cb | + | + | + | + | + | + | + | + |

Source: EPA, Office of Atmospheric Programs
+ Does not exceed 0.05 Gg

Ambient Air Pollutant Emissions

In the United States, carbon monoxide (CO), nitrogen oxides (NO_x), nonmethane volatile organic compounds (NMVOCs), and sulfur dioxide (SO₂) are referred to as “ambient air pollutants,” as termed in the Clean Air Act. These pollutants do not have a direct global warming effect, but indirectly affect terrestrial radiation absorption by influencing the formation and destruction of tropospheric and stratospheric ozone, or, in the case of SO₂, by affecting the absorptive characteristics of the atmosphere. Carbon monoxide is produced when carbon-containing fuels are

combusted incompletely. Nitrogen oxides (i.e., NO and NO₂) are created by lightning, fires, fossil fuel combustion, and in the stratosphere from nitrous oxide (N₂O). NMVOCs—which include hundreds of organic compounds that participate in atmospheric chemical reactions (i.e., propane, butane, xylene, toluene, ethane and many others)—are emitted primarily from transportation, industrial processes, and non-industrial consumption of organic solvents. In the United States, SO₂ is primarily emitted from coal combustion for electric power generation and the metals industry.

Table ES-14: Emissions of NO_x, CO, NMVOCs, and SO₂ (Gg)

| Gas/Activity | 1990 | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 |
|--|----------------|----------------|----------------|----------------|---------------|---------------|---------------|----------------|
| NO_x | 23,037 | 22,509 | 22,360 | 22,289 | 21,961 | 21,341 | 20,917 | 20,141 |
| Stationary Fossil Fuel Combustion | 9,884 | 9,822 | 9,540 | 9,578 | 9,419 | 8,716 | 8,226 | 7,826 |
| Mobile Fossil Fuel Combustion | 12,134 | 11,784 | 11,714 | 11,768 | 11,592 | 11,582 | 11,395 | 11,254 |
| Oil and Gas Activities | 139 | 100 | 126 | 130 | 130 | 113 | 115 | 117 |
| Waste Combustion | 82 | 88 | 135 | 140 | 145 | 142 | 149 | 149 |
| Industrial Processes | 769 | 682 | 808 | 634 | 635 | 748 | 992 | 755 |
| Solvent Use | 1 | 3 | 3 | 3 | 3 | 3 | 3 | 3 |
| Field Burning of Agricultural Residues | 28 | 29 | 32 | 34 | 35 | 34 | 35 | 35 |
| Waste | 0 | 1 | 3 | 3 | 3 | 3 | 3 | 3 |
| CO | 130,575 | 109,149 | 104,063 | 101,132 | 98,976 | 95,464 | 93,965 | 100,653 |
| Stationary Fossil Fuel Combustion | 4,999 | 5,383 | 3,935 | 3,927 | 3,927 | 4,941 | 4,163 | 4,169 |
| Mobile Fossil Fuel Combustion | 119,482 | 97,755 | 93,409 | 90,284 | 87,940 | 84,574 | 83,680 | 90,268 |
| Oil and Gas Activities | 302 | 316 | 321 | 333 | 332 | 152 | 152 | 153 |
| Waste Combustion | 978 | 1,073 | 2,628 | 2,668 | 2,826 | 2,833 | 2,914 | 2,916 |
| Industrial Processes | 4,124 | 3,958 | 3,016 | 3,153 | 3,163 | 2,145 | 2,214 | 2,327 |
| Solvent Use | 4 | 5 | 1 | 1 | 1 | 46 | 45 | 44 |
| Field Burning of Agricultural Residues | 685 | 656 | 747 | 761 | 781 | 760 | 784 | 762 |
| Waste | 1 | 2 | 5 | 5 | 5 | 14 | 14 | 14 |
| NMVOCs | 20,937 | 19,520 | 17,184 | 16,994 | 16,403 | 16,245 | 15,418 | 15,148 |
| Stationary Fossil Fuel Combustion | 912 | 973 | 1,018 | 1,016 | 1,016 | 1,312 | 1,088 | 1,087 |
| Mobile Fossil Fuel Combustion | 10,933 | 8,744 | 8,306 | 7,928 | 7,742 | 7,658 | 7,230 | 6,800 |
| Oil and Gas Activities | 555 | 582 | 433 | 442 | 440 | 376 | 348 | 357 |
| Waste Combustion | 222 | 237 | 304 | 313 | 326 | 326 | 332 | 333 |
| Industrial Processes | 2,426 | 2,643 | 1,997 | 2,038 | 2,047 | 1,890 | 1,845 | 1,829 |
| Solvent Use | 5,217 | 5,609 | 4,969 | 5,100 | 4,671 | 4,533 | 4,422 | 4,584 |
| Field Burning of Agricultural Residues | NA | NA | NA | NA | NA | NA | NA | NA |
| Waste | 673 | 731 | 158 | 157 | 161 | 151 | 153 | 158 |
| SO₂ | 20,936 | 16,892 | 16,682 | 17,091 | 17,189 | 16,013 | 14,802 | 14,324 |
| Stationary Fossil Fuel Combustion | 18,407 | 14,724 | 14,746 | 15,104 | 15,191 | 14,073 | 12,883 | 12,367 |
| Mobile Fossil Fuel Combustion | 793 | 673 | 649 | 659 | 665 | 701 | 632 | 636 |
| Oil and Gas Activities | 390 | 334 | 304 | 312 | 310 | 275 | 279 | 281 |
| Waste Combustion | 39 | 42 | 29 | 29 | 30 | 29 | 29 | 30 |
| Industrial Processes | 1,306 | 1,117 | 953 | 985 | 991 | 933 | 977 | 1,008 |
| Solvent Use | 0 | 1 | 1 | 1 | 1 | 1 | 1 | 1 |
| Field Burning of Agricultural Residues | NA | NA | NA | NA | NA | NA | NA | NA |
| Waste | 0 | 1 | 1 | 1 | 1 | 1 | 1 | 1 |

Source: (EPA 2003) except for estimates from field burning of agricultural residues.

+ Does not exceed 0.5 Gg

NA (Not Available)

Note: Totals may not sum due to independent rounding.

Box ES-4: Sources and Effects of Sulfur Dioxide

Sulfur dioxide (SO₂) emitted into the atmosphere through natural and anthropogenic processes affects the Earth's radiative budget through its photochemical transformation into sulfate aerosols that can (1) scatter radiation from the sun back to space, thereby reducing the radiation reaching the Earth's surface; (2) affect cloud formation; and (3) affect atmospheric chemical composition (e.g., by providing surfaces for heterogeneous chemical reactions). The indirect effect of sulfur-derived aerosols on radiative forcing can be considered in two parts. The first indirect effect is the aerosols' tendency to decrease water droplet size and increase water droplet concentration in the atmosphere. The second indirect effect is the tendency of the reduction in cloud droplet size to affect precipitation by increasing cloud lifetime and thickness. Although still highly uncertain, the radiative forcing estimates from both the first and the second indirect effect are believed to be negative, as is the combined radiative forcing of the two (IPCC 2001). However, because SO₂ is short-lived and unevenly distributed in the atmosphere, its radiative forcing impacts are highly uncertain.

Sulfur dioxide is also a major contributor to the formation of regional haze, which can cause significant increases in acute and chronic respiratory diseases. Once SO₂ is emitted, it is chemically transformed in the atmosphere and returns to the Earth as the primary source of acid rain. Because of these harmful effects, the United States has regulated SO₂ emissions in the Clean Air Act.

Electricity generation is the largest anthropogenic source of SO₂ emissions in the United States, accounting for 69 percent in 2001. Coal combustion contributes nearly all of those emissions (approximately 92 percent). Sulfur dioxide emissions have decreased in recent years, primarily as a result of electric power generators switching from high sulfur to low sulfur coal and installing flue gas desulfurization equipment.

Ambient air pollutants are regulated under the Clean Air Act in an effort to protect human health and the environment. These gases also indirectly affect the global climate by either acting as short-lived greenhouse gases or reacting with other chemical compounds in the atmosphere to form compounds that are greenhouse gases. Unlike the other ambient air pollutants, sulfur-containing compounds emitted into the atmosphere affect the Earth's radiative budget negatively; therefore, it is discussed separately.

One important indirect climate change effect of NMVOCs and NO_x is their role as precursors for tropospheric ozone formation. They can also alter the atmospheric lifetimes of other greenhouse gases. Another example of ambient air pollutant formation into greenhouse gases is carbon monoxide's interaction with the hydroxyl radical—the major atmospheric sink for methane emissions—to form CO₂. Therefore, increased atmospheric concentrations of CO limit the number of hydroxyl molecules (OH) available to destroy methane.

Since 1970, the United States has published estimates of annual emissions of ambient air pollutants (EPA 2003).¹³ Table ES-14 shows that fuel combustion accounts for the majority of emissions of these gases. Industrial processes—such as the manufacture of chemical and allied products, metals processing, and industrial uses of solvents—are also significant sources of CO, NO_x, and NMVOCs.

Uncertainty in and Limitations of Emission Estimates

While the current U.S. emissions inventory provides a solid foundation for the development of a more detailed and comprehensive national inventory, there are uncertainties associated with the emission estimates. Some of the current estimates, such as those for CO₂ emissions from energy-related activities and cement processing, are considered to be highly accurate. For some other categories of emissions, however, a lack of data or an incomplete understanding of how emissions are generated limits the scope or accuracy of the estimates presented. Despite these uncertainties, the UNFCCC reporting guidelines follow the recommendation of the Revised 1996 IPCC Guidelines for National Greenhouse Gas Inventories (IPCC/UNEP/OECD/IEA 1997) and require that countries provide single point estimates for each gas and emission or removal source category.

Currently, a qualitative discussion of uncertainty is presented for all source and sink categories. Within the discussion of each emission source, specific factors affecting the accuracy of the estimates are discussed. A limited number of sources also contain a quantitative uncertainty assessment. Beginning with the Inventory submission in 2004, the United States will provide quantitative estimates for all source and sink categories, in accordance with the new UNFCCC reporting guidelines.

¹³ NO_x and CO emission estimates from field burning of agricultural residues were estimated separately, and therefore not taken from EPA (2003).