



Health Benefits and Costs of Filtration

Wanyu Rengie Chan, Ph.D.

wrchan@lbl.gov https://iaqscience.lbl.gov/

Deputy Group Leader, Indoor Environment Group, Energy Technologies Area



EPA CLEAN AIR SPACES Indoor Air Filtration to Protect PublicHealth During Wildland Fire Smoke EpisodesJune 13, 2019

Outline

- 1. Cost effectiveness of filtration interventions in homes to reduce exposure to PM2.5 during wildfires
- 2. Filtration in classrooms to provide clean air at all times
- 3. Guidance to commercial building operators on how to response to wildfire smoke



INDOOR AIR *doi:10.1111/ina.12285*



Health benefits and costs of filtration interventions that reduce indoor exposure to PM2.5 during wildfires

Abstract Increases in hospital admissions and deaths are associated with increases in outdoor air particles during wildfires. This analysis estimates the health benefits expected if interventions had improved particle filtration in homes in Southern California during a 10-day period of wildfire smoke exposure. Economic benefits and intervention costs are also estimated. The six interventions implemented in all affected houses are projected to prevent 11% to 63% of the hospital admissions and 7% to 39% of the deaths attributable to wildfire particles. The fraction of the population with an admission attributable to wildfire smoke is small, thus, the costs of interventions in all homes far exceeds the economic benefits of reduced hospital admissions. However, the estimated economic value of the prevented deaths exceed or far exceed intervention costs for interventions that do not use portable air cleaners. For the interventions with portable air cleaner use, mortality-related economic benefits exceed intervention costs as long as the cost of the air cleaners, which have a multi-year life, are not attributed to the short wildfire period. Cost effectiveness is improved by intervening only in the homes of the elderly who experience most of the health effects of particles from wildfires.

W. J. Fisk, W. R. Chan

Indoor Environment Group, Lawrence Berkeley National Laboratory, Berkeley, CA, USA

Key words: Benefits; Costs; Health; Filtration; Wildfires; Homes.

W. J. Fisk Indoor Environment Group Lawrence Berkeley National Laboratory 1 Cyclotron Road Berkeley CA 94720 USA Tel.: + 510 486 5910 e-mail: wjfisk@lbl.gov

Received for review 5 November 2015. Accepted for publication 28 January 2016.



Work is sponsored by U.S. EPA, Office of Radiation & Indoor Air, Indoor Environments Division.

2003 Southern California Wildfires



- Impacts on six counties over a 10-day period
 - PM2.5 levels (Wu 2006)
 - Hospital admission rates (Delfino 2009)
 - Effects on mortality (Kochi 2012)

Q: Can filtration substantially reduce wildfirerelated hospitalizations and deaths? Are filtration interventions cost-effective?



Filtration Interventions

 Mass balance models to estimate PM2.5 in homes with and without filtration interventions





- 1. Upgrade to high efficiency air filters (MERV 12*)
- 2. Change to Fan ON mode to run system continuously
- 3. 1 + 2
- 4. Buy an air cleaner (HEPA filter, 1 AER)
- 5. Buy an air cleaner + 1 + 2

* Reduced to MERV 9 because of air leakage bypass.



- Assumed that health effects depend on the total inhalation intake of PM2.5: home, outside of home (work, school), vehicle, outdoor
- Because health outcomes have been associated to outdoor PM2.5, we estimated the health benefits filtration interventions in home by calculating an equivalent reductions in outdoor PM2.5
- Calculations were also performed for 22% of homes in the study area with age 65+ residents (assumed home at all times)



Filtration interventions in all 6.92 million homes exposed to 2003 southern California wildfires would have prevented 47 to 261 respiratory hospital admissions (11-63% reduction), and 9 to 52 premature deaths (7-39% reduction), related to PM2.5 exposures during the 10-day study period.

Fisk and Chan (2017). Health benefits and costs of filtration interventions that reduce indoor exposure to PM2.5 during wildfires. Indoor Air.



Intervention Costs of Filtration for 6.92m homes

- Operating forced air system continuously is expected to cost \$110m in electricity.
- Incremental cost of purchasing higher efficiency filters + electricity cost is \$133m.
- Energy costs of operating air cleaners is \$13 to 16m.
- Estimated cost of buying portable air cleaners range from \$1.8 to \$4.4b*.

*Though it is unlikely that anyone would make the purchase just for the 10-day period.








Summary

- Economic benefits from reduced hospitalization and premature deaths exceed intervention costs for all 6.92 million homes exposed to wildfire smoke, but are not sufficient to pay for portable air cleaners.
 - Estimated benefit-to-cost ratio range is about 2:1 using forced air system filtration interventions in all homes
 - For 22% of homes with age 65+ residents: benefit-to-cost = 16:1
- In 22% of homes with age 65+ residents, benefits from reduced hospitalization and premature deaths are sufficient to pay for purchase of portable air cleaners.



Wildfire Smoke Exposure in Classrooms

• Camp Fire, Nov 8-22, 2018





Classroom HVAC Retrofit Evaluation Study



FCAL





Phase II Field Study Goal


 Evaluate energy efficiency and IEQ performance of HVAC system retrofits in two California schools



Sacramento region high school (7 portable classrooms)

- Ventilation system types: CRV, ERV, ECON-DCV
- MERV 8 (Airguard DP Pleat 2-inch), MERV 13 (Airguard DP-G13EEN 2-inch)





IAQ Monitoring (week-long, 3x)


- **PM2.5**
- **Black carbon**
- Formaldehyde
- O₃ CO₂








PM2.5 Measured during Wildfire Smoke (Camp Fire)



California Air Resources Board, Air Quality and Meteorological Information System (AQMIS). https://www.arb.ca.gov/aqmis2/aqmis2.php



Daily Mean PM2.5 (ug/m³) Measured During School Hours





Summary

• PM2.5 substantially lower in classrooms with MERV 13 filters compared to MERV 8, both during wildfires <u>and</u> normal conditions

I/O Ratio	MERV 8	MERV 13
CRV	0.76–0.78	0.48–0.53
ERV	0.56-0.59	0.35–0.38
ECON-DCV	0.60–0.65	0.32–0.46
Normal conditions (DustTrak)	0.28-0.42	0.10-0.13

 Other important considerations: classrooms must provide adequate ventilation during normal conditions, proper filter installation & regular maintenance





About the Lab

Leadership/Organization

News Ce

Calendar

NEWS CENTER

Q&A: How to Protect Yourself and Your Family From Wildfire Smoke

Stay indoors, consider a mask, limit activities, use air filtration systems, or even build your own

Feature Story Laurel Kellner 510-486-5375 • NOVEMBER 14, 2018

f 6.5K	¥	6	G+	in	< 6.5K
--------	---	---	----	----	--------



Q. How worried should I be about the a pollution from a wildfire?


Q. How can I protect myself during wildfire air pollution events?

Q. How can I make my home safer?

Q. How do I use filtration systems in my home?

https://newscenter.lbl.gov/2018/11/14/qa-how-toprotect-yourself-and-your-family-from-wildfire-smoke/








Smoke Mitigation Efforts at LBNL

- At our main site, LBNL is operating ~2m sqft in 100+ buildings
- 60% of our largest, most populated buildings are 50+ years old
- HVAC smoke mitigation changes during 2018 wildfires:
 - 46 AHU in 21 buildings had HVAC control overwritten or manually controlled to recirculate conditioned air
 - 87% AHU were overridden to ~10% OA, remaining set to 100% recirculation


PM2.5 I/O ratio for buildings that had or did not have outdoor air reduced (data from LBNL EH&S)





Priority for Smoke Mitigation







HVAC: Improved Building Filtration System

Facilities, EH&S, HR, management working together on preparedness: smoke mitigation mode, better filter, monitoring (PM, CO2), response plan



Efficacy





https://iaqscience.lbl.gov/ Protecting Public from Wildfire Smoke

Brett Singer, Ph.D.bcsinger@lbl.govStaff Scientist510.486.4779Group Leader, Indoor Environment GroupEnergy Technologies AreaWanyu Rengie Chan, Ph.D.wrchan@lbl.gov510.486.6570



