

Technical Appendix C

Derivation of Model Exposure Parameters

Table of Contents

- 1 Introduction..... 3**
- 2 Exposure Parameters..... 3**
 - 2.1 Body Weights..... 4
 - 2.2 Inhalation 4
 - 2.3 Drinking Water Ingestion 6
 - 2.4 Fish Consumption 8
- 3 References 10**

1 Introduction

This appendix provides information on the data and methods used to derive subpopulation exposure parameters for use in the Indicators model. Three human exposure parameters (an inhalation rate, fish ingestion rate, and drinking water ingestion rate) and body weight are currently used in the RSEI model to generate sex-specific exposure factors for four age groups (ages #17, 18-44, 45-64, 65+). Earlier versions of the model used standard assumptions to represent intake for all individuals within the general population. However, because there are population-specific intake differences and because some populations may be more susceptible to certain chemicals than others, subpopulation intake and census data have been included in the model to estimate a more accurate surrogate dose.

2 Exposure Parameters

The most recent *Exposure Factors Handbook* (EPA, 2011) is the primary source of information used to generate exposure parameters for both inhalation and drinking water ingestion. Fish consumption data was obtained directly from EPA's Office of Water (EPA, 20002, based on an analysis of the 1994-1996 USDA Continuing Survey of Food Intake in Individuals (CSFII).

The exposure parameters were generated in such a way as to ensure as much consistency as possible among pathways, while basing the estimates on values recommended in the *Exposure Factors Handbook* (EPA, 2011), when available. The parameters generally reflect those recommended in EPA (2011), however, some estimates are derived from data in EPA (2011) that were not explicitly included in their summarized recommendations.

The exposure factors in EPA (2011) are available for a large number of discrete age groups, particularly for children. The RSEI age groups are generally broader than those in EPA (2011). To match the RSEI age groups, we calculate an average of exposure factors for all ages within the RSEI age group. The equation C-1 is shown below. Intake rates are then adjusted by body weight estimates from EPA (2011), when necessary. The sections below provide further detail on the calculation of pathway-specific exposure parameters.

Equation C-1:

$$\text{RSEI Exposure Factors} = \frac{\sum_i (IR_i \times n_i)}{N}$$

where:

IR is the intake rate for age group i ,
 n is the number of years in age group i , and
 N is the total number of years in the RSEI model age group for all age groups i that fall within the RSEI age group.

2.1 Body Weights

EPA (2011) provides updated estimates of mean body weight for boys and girls, for small increments up to one year, one-year age increments from one year up to 19 years of age and for several adult age groups. Generally, these estimates are higher than those provided in the previous EFH version.

Sex-specific body weights were averaged across the range of each RSEI age group and are presented in Table C-1.

Table C-1. Body Weights for Each RSEI Age Group from EPA (2011)

Model Age Group	Body Weight (kg)	
	Male	Female
0-17	38.4	36.5
18-44	85.8	73.2
45-64	89.7	77.2
65+	81.5	69.3

Source: EPA (2011), Tables 8-4 and 8-5, pp. 8-13 and 8-1.4

2.2 Inhalation

EPA (2011) recommended new studies as the basis for inhalation rates for both adults and children. For adults, EPA based their inhalation rates on three recent studies: Brochu et al. (2006a, as cited in EPA 2011), Stifelman et al. (2007, as cited in EPA 2011), and EPA (2009, as cited in EPA 2011). Additionally, for children EPA based their suggested inhalation rates on Arcus-Arth and Blaisdell (2007, as cited in EPA 2011). Data from these four studies were combined, where appropriate. If the data were combined from multiple studies, they were averaged by sex and grouped according to the age groups selected for use in the *Exposure Factors Handbook*. If age groups in the original reference did not match the EPA groupings in the Handbook, statistics were averaged from all age groupings in the original reference that overlapped with EPA's age groupings by more than one year, weighted by the number of observations contributed from each age group. EPA's final inhalation rate estimates are presented in Table C-2 along with the reference(s) used to derive them.

Table C-2. Inhalation Rates Recommended by EPA (2011)

Age	Mean (m ³ /day)	Reference
0 - <0.083 year (1 month)	3.6	1
0.083 - 0.25 year	3.5	2,3
0.25 - < 0.5 year	4.1	2,3
0.5 - <1 year	5.4	2,3
1 - <2 years	5.4	1,2,3,4
2 - <3 years	8.9	1,2,3,4
3 - <6 years	10.1	1,2,3,4
6 - <11 years	12	1,2,3,4
11 - <16 years	15.2	1,2,3,4
16 - <21 years	16.3	1,2,3,4
21 - <31 years	15.7	2,3,4
31 - <41 years	16	2,3,4
41 - <51 years	16	2,3,4
51 - <61 years	15.7	2,3,4
61 - <71 years	14.2	2,3,4
71 - <81 years	12.9	2,4
≥81 years	12.2	2,4

¹ Arcus-Arth and Blaisdell 2007 (as cited in EPA 2011)

² Brochu et al. 2006a (as cited in EPA 2011)

³ EPA 2009 (as cited in EPA 2011)

⁴ Stifelman 2007 (as cited in EPA 2011)

Source: Table 6-1, page 6-3 (incorrectly labeled 6-1) in EPA (2011)

The inhalation rates and body weights recommended by EPA (2011) were adjusted for the RSEI age groups using the weighted average approach explained in the previous section. Table C-2 below shows the adjusted inhalation rates. Male and female rates are the same because EPA (2011) only presented combined rates for both sexes.

Table C-3. Inhalation Rates, Based on EPA (2011), Adjusted for RSEI Age-Sex Groups

Age	Mean Inhalation Rate (m ³ /day)	
	Male	Female
0-17 years	12.1	12.1
18-44 years	15.9	15.9
45-64 years	15.5	15.5
65+ years	12.9	12.9

Source: Calculated from EPA (2011)

For adults and children, the age and sex-specific inhalation values were adjusted by body weight using estimates recommended by EPA and presented in the previous section. The final inhalation exposure factors used in the model are given in Table C-4.

Table C-4. Inhalation Exposure Factors Used in RSEI

Age	Exposure Factor (m ³ /kg-day)	
	Male	Female
0-17 years	0.315	0.332
18-44 years	0.185	0.217
45-64 years	0.173	0.201
65+ years	0.159	0.187

2.3 Drinking Water Ingestion

In the February 2019 update to Chapter 3 of the EFH, EPA's recommended values are taken from the Agency's own analysis of 2005-2010 NHANES data. These values are summarized in Table C-5 below.

Table C-5. EPA (2019) Recommended Tap Water Intake Estimates

Age range	Mean	
	Tap water intake (ml/day)	Tap water intake per kg of body weight (ml/kg-day)
0 to <1 month	184	42
1 to <3 months	145	25
3 to <6 months	187	27
6 months to <1 year	269	30

Age range	Mean	
	Tap water intake (ml/day)	Tap water intake per kg of body weight (ml/kg-day)
1 to <2 years	146	13
2 to <3 years	205	15
3 to <6 years	208	11
6 to <11 years	294	10
11 to <16 years	315	6
16 to <21 years	436	6
21 to <30 years	781	10
30 to <40 years	902	11
40 to <50 years	880	11
50 to <60 years	956	12
60 to <70 years	941	12
70 to <80 years	772	10
80+	784	11
21 to <50 years	858	11
50+ years	902	11
All ages	711	11

Source: These values are the per capita intake values from Table 3-1, page 3-3 in EPA (20191).

To convert these drinking water intake rates into exposure factors to fit the RSEI model they need be grouped according to the age categories used in RSEI. Specifically, RSEI's exposure factors are split into 4 age categories 0-17, 18-44, 45-64 and 65+ and are in units of L/kg-day. Therefore, to begin, mL/kg-day estimates were converted to L/kg-day for each age group. Then, to match the RSEI age groups, Abt calculated an average of EFs for all ages within the RSEI age group using the following equation:

$$EF = \frac{\sum(IR_i \times n_i)}{N}$$

Where:

(Equation 1)

EF= the exposure factor in L/kg-day

IR = the intake rate for age group I (L/kg-day)

n = the number of years in age group i

N = the total number of years in the RSEI model age group for all age groups i that fall within the RSEI age group.

The final drinking water ingestion rates used in the model are shown below in Table C-6.

Table C-6. EPA (2019) Recommended Tap Water Intake Estimates

Model Age Group	Exposure Factors (Male)	Exposure Factors (Female)
	<i>(L/kg-day)</i>	
0-17	0.0101	0.0101
18-44	0.0099	0.0099
45-64	0.0117	0.0117
>65	0.0108	0.0108

2.4 Fish Consumption

Data on fish consumption (g/day) by age group and gender were obtained directly from EPA’s Office of Water (EPA, 2002). The data is based on the 1994-1996 USDA Continuing Survey of Food Intake by Individuals (CSFII). Data on freshwater/estuarine fish consumption was available for three broad age groups: 14 and younger, 15-44 years old, and 45 and older. To estimate exposure parameters for recreational consumers, the 90th percentile of intake was used, while for subsistence consumers, the 99th percentile was chosen. Table C-7 shows the consumption values for recreational and subsistence consumers.

Table C-7. Fish Consumption Intake Data, CSFII 94-96¹

Age	Sex	Fish Consumption ¹ (g/day)	
		Recreational	Subsistence
<15	Male	0.00	79.03
	Female	0.00	58.83
15-44	Male	15.63	151.19
	Female	6.31	109.79
45+	Male	32.47	165.92
	Female	17.65	108.80

¹ Fish consumption data comes from EPA (2002, Section 5.1.1.1, Table 1, p. 5-3). Data is based on the 1994-96 USDA Continuing Survey of Food Intakes by Individuals (CSFII). The 90th percentile is used to represent recreational consumers and 99th percentile is used to represent subsistence consumers.

To estimate fish ingestion values for the RSEI age groups, average intake rates were calculated using Equation C-1. For example, in order to calculate fish ingestion rates for the RSEI #17 year

old age group, the intake rate for <15 year olds is multiplied by 15 and the intake rate for 15-44 year olds is multiplied by 3. These products are then summed and divided by the total number of years in the RSEI age group, 18. The fish ingestion intakes and body weights for each of the model age groups are presented in Table C-8. The corresponding fish ingestion exposure factors used in the RSEI model are shown in Table C-9.

Table C-8. Fish Ingestion Values and Body Weights for Each RSEI Age Group from EPA (2002) and EPA (2011)

Model Age Group	Recreational Fish Ingestion (g/day) ¹		Subsistence Fish Ingestion (g/day) ¹	
	Male	Female	Male	Female
0-17	2.61	1.05	91.1	67.3
18-44	15.6	6.31	151	110
45-64	32.5	17.7	166	109
65+	32.5	17.7	166	109

¹ See text for discussion of method used to calculate ingestion values.

Table C-9. Fish Ingestion Exposure Factors Used in RSEI Model

Model Age Group	Recreational (g/kg-day) ¹		Subsistence (g/kg-day) ¹	
	Male	Female	Male	Female
0-17	0.0756	0.0372	2.83	2.05
18-44	0.199	0.114	1.92	1.71
45-64	0.407	0.262	2.08	1.60
≥65	0.434	0.267	2.22	1.63

¹ Fish ingestion exposure factors are converted to kg/kg-day for the surrogate dose calculation in the RSEI model.

3 References

Canadian Ministry of National Health and Welfare. 1981. *Tapwater Consumption in Canada*. Doc # 82-EHD-80. Public Affairs Directorate, Department of National Health and Welfare, Ottawa, Canada.

Ershow AG and Cantor KP. 1989. *Total Water and Tapwater Intake in the United States: Population-Based Estimates of Quantities and Sources*. Life Sciences Research Office, Federation of American Societies for Experimental Biology.

National Center for Health Statistics (NCHS, 1987. *Anthropometric Reference Data and Prevalence of Overweight, United States, 1976-80*. Data from the National Health and Nutrition Examination Survey, Series 11, No. 238. U.S. Department of Health and Human Services, Public Health Service, National Center for Health Statistics. DHHS Publication No. (PHS) 87-1688.

U.S. Environmental Protection Agency (EPA, 2002). *Estimated Per Capita Fish Consumption in the United States*. Office of Water and Office of Science and Technology.
<https://nepis.epa.gov/Exe/ZyPURL.cgi?Dockey=901R0600.txt>.

U.S. Environmental Protection Agency (EPA, 2011). *Exposure Factors Handbook*. Office of Health and Environmental Assessment. EPA/600/R-09/052F.
<https://cfpub.epa.gov/ncea/risk/recordisplay.cfm?deid=236252>.

U.S. Environmental Protection Agency. (EPA, 2019). *Exposure Factors Handbook Chapter 3 (Update): Ingestion of Water and Other Select Liquids*. Office of Research and Development, Washington, DC, EPA/600/R-18/259F, 2019.
<https://cfpub.epa.gov/ncea/efp/recordisplay.cfm?deid=343661>.

[revised 12/22/2020]