


# Final Scope of the Risk Evaluation for 1,2-Dichloropropane

## CASRN 78-87-5



August 2020

## TABLE OF CONTENTS

ACKNOWLEDGEMENTS	6
ABBREVIATIONS AND ACRONYMS	7
EXECUTIVE SUMMARY	11
1 INTRODUCTION	14
2 SCOPE OF THE EVALUATION	
2.1 Reasonably Available Information	
2.1.1 Search of Gray Literature	
2.1.2 Search of Literature from Publicly Available Databases (Peer-Reviewed Literature)	
2.1.3 Search of TSCA Submissions	
2.2 Conditions of Use	
2.2.1 Categories and Subcategories of Conditions of Use Included in the Scope of the Risk	
Evaluation	27
2.2.2 Activities Excluded from the Scope of the Risk Evaluation	29
2.2.3 Production Volume	29
2.2.4 Overview of Conditions of Use and Lifecycle Diagram	30
2.3 Exposures	
2.3.1 Physical and Chemical Properties	
2.3.2 Environmental Fate and Transport	
2.3.3 Releases to the Environment	
2.3.4 Environmental Exposures	
2.3.5 Occupational Exposures	
2.3.6 Consumer Exposures	
2.3.7 General Population Exposures	
2.4 Hazards (Effects)	
<ul><li>2.4.1 Environmental Hazards</li><li>2.4.2 Human Health Hazards</li></ul>	
<ul><li>2.5 Potentially Exposed or Susceptible Subpopulations</li><li>2.6 Conceptual Models</li></ul>	
2.6.1 Conceptual Model for Industrial and Commercial Activities and Uses	
2.6.2 Conceptual Model for Endustrial and Commercial Activities and Uses: Potential Exposures and Hazard	
2.6.3 Conceptual Model for Environmental Releases and Wastes: Potential Exposures and	15 +5
Hazards (Regulatory Overlay)	
2.6.3.1 Exposure Pathways and Risks Addressed by Other EPA Administered Statutes	
2.6.3.2 Conceptual Model for Environmental Releases and Wastes: Potential Exposures a	
Hazards 54	
2.7 Analysis Plan	56
2.7.1 Physical and Chemical Properties and Environmental Fate	
2.7.2 Exposure	
2.7.2.1 Environmental Releases	57
2.7.2.2 Environmental Exposures	
2.7.2.3 Occupational Exposures	
2.7.2.4 Consumer Exposures	
2.7.2.5 General Population	63

2.7.3 Hazards (Effects)	
2.7.3.1 Environmental Hazards	
2.7.3.2 Human Health Hazards	
2.7.4 Summary of Risk Approaches for Characterization	
2.8 Peer Review	
REFERENCES	
APPENDICES	
Appendix A ABBREVIATED METHODS FOR SEARCHING AND SCREENING	. 75
A.1 Literature Search of Publicly Available Databases	75
A.1.1 Search Term Genesis and Chemical Verification	75
A.1.2 Publicly Available Database Searches	
A.1.2.1 Query Strings for the Publicly-Available Database Searches on 1,2-Dichloropropane	76
A.1.2.2 Data Prioritization for Environmental Hazard, Human Health Hazard, Fate and Physi	cal
Chemistry	81
A.1.2.3 Data Prioritization for Occupational Exposures and Environmental Releases and	
General Population, Consumer and Environmental Exposures	82
A.2 Peer-Reviewed Screening Process	82
A.2.1 Inclusion/Exclusion Criteria	83
A.2.1.1 PECO for Environmental and Human Health Hazards	83
A.2.1.2 PECO for Consumer, Environmental, and General Population Exposures	85
A.2.1.3 RESO for Occupational Exposure and Environmental Releases	86
A.2.1.4 PESO for Fate and Transport	88
A.2.1.5 Generation of Hazard Heat Maps	90
A.3 Gray Literature Search and Screening Strategies	91
A.3.1 Screening of Gray Literature	
A.3.2 Initial Screening of Sources using Decision Logic Tree	92
A.3.3 TSCA Submission Searching and Title Screening	
A.3.4 Gray Literature Search Results for 1,2-Dichloropropane	94
Appendix B PHYSICAL AND CHEMICAL PROPERTIES OF 1,2-DICHLOROPROPANE .	. 97
Appendix C ENVIRONMENTAL FATE AND TRANSPORT PROPERTIES OF 1,2-	
DICHLOROPROPANE	
Appendix D REGULATORY HISTORY	. 99
D.1 Federal Laws and Regulations	99
D.2 State Laws and Regulations	106
D.3 International Laws and Regulations	
Appendix E PROCESS, RELEASE AND OCCUPATIONAL EXPOSURE INFORMATION	
E.1 Process Information	110
E.1.1 Manufacture (Including Import)	
E.1.1.1 Manufacture	
E.1.1.2 Import	
E.1.2 Processing and Distribution	
E.1.2.1 Processing as a Reactant	
E.1.2.2 Incorporation into Formulation, Mixture, or Reaction Product	111

E.1.3 Us	es	111
E.1.3.1	Industrial Use – Non-Incorporative Activities	111
E.1.3.2	Commercial Laboratory Use	111
E.1.3.3	Commercial Cleaning and Furnishing Care Products	111
E.1.4 Dis	sposal	111
E.2 Prelin	ninary Occupational Exposure Data	111
Appendix F	SUPPORTING INFORMATION - CONCEPTUAL MODEL FOR IN AND COMMERCIAL ACTIVITIES AND USES	
11	SUPPORTING INFORMATION - CONCEPTUAL MODEL FOR C ACTIVITIES AND USES	
Appendix H	SUPPORTING INFORMATION – CONCEPTUAL MODEL FOR ENVIRONMENTAL RELEASES AND WASTES	118

### LIST OF TABLES

Table 2-1. Results of Title Screening of Submissions to EPA under Various Sections of TSCA <sup>a</sup>	26
Table 2-2. Categories and Subcategories of Conditions of Use Included in the Scope of the Risk	
Evaluation	27
Table 2-3. Physical and Chemical Properties of 1,2-Dichloropropane	32
Table 2-4. Summary of 1,2-Dichloropropane TRI Production-Related Waste Managed in 2018	35
Table 2-5. Summary of Releases of 1,2-Dichloropropane to the Environment During 2018	36
Table 2-6. Potential Categories and Sources of Environmental Release Data	57
Table 2-7. Potential Sources of Occupational Exposure Data	60

### LIST OF FIGURES

Figure 2-1. Gray Literature Tags by Discipline for 1,2-Dichloropropane
Figure 2-2. Peer-reviewed Literature Inventory Tree - Physical and chemical Properties Search Results
for 1,2-Dichloropropane17
Figure 2-3. Peer-reviewed Literature Inventory Tree – Fate and Transport Search Results for 1,2-
Dichloropropane
Figure 2-4. Peer-reviewed Literature Inventory Heat Map – Fate and Transport Search Results for 1,2-
Dichloropropane
Figure 2-5. Peer-reviewed Literature Inventory Tree - Engineering Search Results for 1,2-
Dichloropropane
Figure 2-6. Peer-reviewed Literature Inventory Heat Map - Engineering Search Results for 1,2-
Dichloropropane
Figure 2-7. Peer-Reviewed and Gray Literature Inventory Tree - Exposure Search Results for 1,2-
Dichloropropane
Figure 2-8. Peer-reviewed and Gray Literature Inventory Heat Map – Exposure Search Results for 1,2-
Dichloropropane
Figure 2-9. Peer-reviewed Literature Inventory Tree - Human Health and Environmental Hazards Search
Results for 1,2-Dichloropropane24
Figure 2-10. Peer-reviewed Literature Inventory Heat Map – Human Health and Environmental Hazards
Search Results for 1,2-Dichloropropane25
Figure 2-11. 1,2-Dichloropropane Life Cycle Diagram

Figure 2-12. Box and Whisker Plots of Reported Physical and Chemical Property Values	ł
Figure 2-13. 1,2-Dichloropropane Conceptual Model for Industrial and Commercial Activities and Uses	
Worker and Occupational Non-User Exposures and Hazards	2
Figure 2-14. 1,2-Dichloropropane Conceptual Model for Consumer Activities and Uses: Consumer	
Exposures and Hazards 44	ł
Figure 2-15. 1,2-Dichloropropane Conceptual Model for Environmental Releases and Wastes:	
Environmental and General Population Exposures and Hazards (Regulatory Overlay) 46	5
Figure 2-16. 1,2-Dichloropropane Conceptual Model for Environmental Releases and Wastes:	
Environmental Exposures and Hazards	5

## LIST OF APPENDIX TABLES

Table Any A.1. Sources of Verification for Chemical Names and Structures	_
Table_Apx A-1. Sources of Verification for Chemical Names and Structures       7:         Table_Apx A-2. Sources of Verification for Chemical Names and Structures       7:	2
Table_Apx A-2. Summary of Data Sources, Search Dates and Number of Peer-Reviewed Literature	_
Search Results for 1,2-Dichloropropane	
Table_Apx A-3. Hazards Title and Abstract and Full-text PECO Criteria for 1,2-Dichloropropane 8.	3
Table_Apx A-4. Major Categories of Potentially Relevant Supplemental Materials for 1,2-	
Dichloropropane	4
Table_Apx A-5. Generic Inclusion Criteria for the Data Sources Reporting Exposure Data on General	
Population, Consumers and Environmental Receptors	
Table_Apx A-6. Pathways Identified as Supplemental for 1,2-Dichloropropane <sup>a</sup>	
Table_Apx A-7. Inclusion Criteria for Data Sources Reporting Engineering and Occupational Exposure	
Data	
Table_Apx A-8. Engineering, Environmental Release and Occupational Data Necessary to Develop the	
Environmental Release and Occupational Exposure Assessments	7
Table_Apx A-9. Inclusion Criteria for Data or Information Sources Reporting Environmental Fate and	
Transport Data	9
Table_Apx A-10. Fate Endpoints and Associated Processes, Media and Exposure Pathways Considered	
in the Development of the Environmental Fate Assessment	9
Table_Apx A-11. Decision Logic Tree Overview	3
Table_Apx A-12. Gray Literature Sources that Yielded Results for 1,2-Dichloropropane	4
Table_Apx B-1. Summary Statistics for Reviewed Physical Properties97	7
Table_Apx C-1. Environmental Fate and Transport Properties of 1,2-Dichloropropane	8
Table_Apx D-1. Federal Laws and Regulations	9
Table_Apx D-2. State Laws and Regulations 100	б
Table_Apx D-3. Regulatory Actions by other Governments, Tribes, and International Agreements 108	8
Table_Apx E-1. Summary of Industry Sectors with 1,2-Dichloropropane Personal Monitoring Air	
Samples Obtained from OSHA Inspections Conducted since 1984	2
Table_Apx F-1. Worker and Occupational Non-User Exposure Conceptual Model Supporting Table 113	3
Table_Apx G-1. Consumer Exposure Conceptual Model Supporting Table 117	7
Table_Apx H-1. General Population and Environmental Exposure Conceptual Model Supporting Table	

### LIST OF APPENDIX FIGURES

Figure	Apx A-1	. Decision Logic Tree	Used to Screen Gra	y Literature Results	
$\mathcal{O}$	— I	$\mathcal{O}$			

### ACKNOWLEDGEMENTS

This report was developed by the United States Environmental Protection Agency (U.S. EPA), Office of Chemical Safety and Pollution Prevention (OCSPP), Office of Pollution Prevention and Toxics (OPPT).

#### Acknowledgements

The OPPT Assessment Team gratefully acknowledges participation or input from Intra-agency reviewers that included multiple offices within EPA, Inter-agency reviewers that included multiple Federal agencies, and assistance from EPA contractors Abt Associates (Contract No. EP-W-16-009), ERG (Contract No. EP-W-12-006), GDIT (Contract No. HHSN316201200013W), ICF (Contract No.68HERC19D0003), SRC (Contract No. 68HERH19F0213), and Versar (Contract No. EP-W-17-006). EPA also acknowledges the contributions of technical experts from EPA's Office of Research and Development.

#### Docket

Supporting information can be found in public docket: EPA-HQ-OPPT-2018-0428.

#### Disclaimer

Reference herein to any specific commercial products, process or service by trade name, trademark, manufacturer or otherwise does not constitute or imply its endorsement, recommendation or favoring by the United States Government.

## ABBREVIATIONS AND ACRONYMS

ACGIH	American Conference of Governmental Industrial Hygienists
ADME	Absorption, Distribution, Metabolism, and Excretion
AMTIC	EPA Ambient Monitoring Technology Information Center
AQS	Air Quality System
ATSDR	Agency for Toxic Substances and Disease Registry
AWQC	Ambient Water Quality Criteria
BAF	Bioaccumulation Factor
BCF	Bioconcentration Factor
BMF	Biomagnification factor
BOD	Biochemical Oxygen Demand
BP	Boiling Point
CAA	Clean Air Act
CalDPR	California Dept of Pesticide Regulation
CalEPA	California Environmental Protection Agency
CASRN	Chemical Abstracts Service Registry Number
CBI	Confidential Business Information
CDC	Centers for Diseases Control and Prevention
CDR	Chemical Data Reporting
CEHD	Chemical Exposure Health Data
CEM	Consumer Exposure Model
CEPA	Canadian Environmental Protection Act
CERCLA	Comprehensive Environmental Response, Compensation and Liability Act
CFR	Code of Federal Regulations
ChemSTEER	Chemical Screening Tool for Occupational Exposures and Releases
CHRIP	Chemical Risk Information Platform
CITI	Chemicals Inspection Testing Institute Japan
COC	Concentration of Concern
CPCat	Chemical and Product Categories
CPDat	Consumer Product Database
CSCL	Chemical Substances Control Law
CWA	Clean Water Act
DCP	1,2-Dichloropropane
DMR	Discharge Monitoring Report
EC	Engineering Control(s)
ECHA	European Chemicals Agency
EPA	Environmental Protection Agency
EPCRA	Emergency Planning and Community Right-to-Know Act
GACT	Generally Available Control Technology
ERG	Eastern Research Group
ESD	Emission Scenario Document
EU	European Union
FIFRA	Federal Insecticide, Fungicide, and Rodenticide Act
FR	Federal Register
FYI	For your information
GC	Gas Chromatography
GDIT	General Dynamics Information Technology

CECTIC	Substance Database contains information for the safe handling of harondous substances
GESTIS	Substance Database contains information for the safe handling of hazardous substances
00	and other chemical substances at work
GS	Generic Scenario
HAP	Hazardous Air Pollutant
HERO	Health and Environmental Research Online
HLC	Henry's Law Constant
HMTA	Federal Hazardous Materials Transportation Act
HPLC	High Performance Liquid Chromatography
HSDB	Hazardous Substances Data Bank
IARC	International Agency for Research on Cancer
IBCs	Intermediate Bulk Containers
ICF	ICF is a global consulting services company
IMAP	Inventory Multi-Tiered Assessment and Prioritisation (Australia)
ISHA	Industrial Safety and Health Act
Koc	Organic Carbon: Water Partition Coefficient
KOECT	Kirk-Othmer Encyclopedia of Chemical Technology
Kow	Octanol: Water Partition Coefficient
LC <sub>x</sub>	Lethal Concentration
LOAELs	Lowest Observed Adverse Effect Level
LOEC	Lowest Observed Effect Concentration
MACT	Maximum Achievable Control Technology
MCL	Maximum Contaminant Level
MCLG	Maximum Contaminant Level Goal
MITI	Ministry of International Trade and Industry
MOA	Mode of Action
MP	Melting Point
MSW	Municipal Solid Waste
NAICS	North American Industry Classification System
NATA	National-scale Air Toxics Assessment
NEI	National Emissions Inventory
NESHAP	National Emission Standards for Hazardous Air Pollutants
NICNAS	National Industrial Chemicals Notification and Assessment Scheme (Australia)
NIOSH	National Institute for Occupational Safety and Health
NITE	National Institute of Technology and Evaluation
NLM	National Library of Medicine
NOAELs	No Observed Adverse Effect Level
NOEC	No Observed Effect Concentration
NPDES	National Pollutant Discharge Elimination System
NPDWR	National Primary Drinking Water Regulation
NPL	National Priorities List
NPRI	National Pollutant Release Inventory
NSPS	New Source Performance Standards
NTP	National Toxicology Program
OCSPP	Office of Chemical Safety and Pollution Prevention
OECD	Organisation for Economic Co-operation and Development
OEHHA	Office of Environmental Health Hazard Assessment (California)
OELs	
OLLS	Occupational Exposure Limits

ONU	Occupational Non-User
OPPT	Office of Pollution Prevention and Toxics
OSHA	Occupational Safety and Health Administration
OW	EPA's Office of Water
PBPK	Physiologically Based Pharmacokinetic
PBT	Persistent, Bioaccumulative, and Toxic
PECO	Population, Exposure, Comparator, Outcome
PEL	Permissible Exposure Limit
PESO	Pathways and Processes, Exposure, Setting or Scenario and Outcomes
PESS	Potentially Exposed or Susceptible Subpopulation
PODs	Points of Departure
POTW	Publicly Owned Treatment Works
PPE	Personal Protective Equipment
QC	Quality Control
RCRA	Resource Conservation and Recovery Act
REACH	Registration, Evaluation, Authorisation and Restriction of Chemicals (European Union)
REL	Recommended Exposure Limit
RESO	Receptors, Exposure, Setting or Scenario, and Outcomes
RIVM	Dutch National Institute for Public Health and the Environment
RQs	Risk Quotients
SARA	Superfund Amendments and Reauthorization Act
SDS	Safety Data Sheet
SDWA	Safe Drinking Water Act
SIDS	Screening Information Dataset
SPIN	Substances in Preparations in Nordic Countries
SRC	SRC Inc., formerly Syracuse Research Corporation
STEL	Short-term Exposure Limit
STORET	Storage and Retrieval for Water Quality Data; EPA's repository of water quality
	monitoring data
SYKE	The Finnish Environment Institute
TBD	To be determined
TERA	Toxicology Excellence for Risk Assessment
TG	Test Guideline
TIAB	Title and Abstract
ТК	Toxicokinetics
TLV	Threshold Limit Value
TMF	Trophic Magnification Factors
TRI	Toxics Release Inventory
TSCA	Toxic Substances Control Act
TTO	Total Toxic Organics
TURA	Toxic Use Reduction Act
TWA	Time-weighted average
UIC	Underground Injection Control
UCMR	Unregulated Contaminants Monitoring Rule
U.S. DOC	U.S. Department of Commerce
USGS	United States Geological Survey
VOC	Volatile Organic Compound

VP	Vapor Pressure
WHO	World Health Organization
WQX	Water Quality Exchange
WS	Water Solubility
WWT	Wastewater Treatment

### **EXECUTIVE SUMMARY**

In December 2019, EPA designated 1,2-dichloropropane (CASRN 78-87-5) as a high priority substance for risk evaluation following the prioritization process as required by Section 6(b) of the Toxic Substances Control Act (TSCA) and implementing regulations (40 CFR Part 702) (Docket ID: <u>EPA-HQ-OPPT-2019-0131</u>). The first step of the risk evaluation process is the development of the draft scope document. EPA published *Draft Scope of the Risk Evaluation for 1,2-Dichloropropane CASRN 78-87-5* (EPA Document No. EPA-740-D-20-006) (U.S. EPA, 2020c) and provided a 45-day comment period on the draft scope per 40 CFR 702.41(c)(7). EPA has considered comments received (Docket ID: <u>EPA-HQ-OPPT-2018-0428</u>) during the public comment period to inform the development of this final scope document, and public comments received will continue to inform the development of the risk evaluation for 1,2-dichloropropane. This document fulfills the TSCA requirement to issue a final scope document per TSCA Section 6(b)(4)(D) and as described in 40 CFR 702.41(c)(8). The scope for 1,2dichloropropane includes the following information: the conditions of use, potentially exposed or susceptible subpopulations (PESS), hazards, and exposures that EPA plans to consider in the risk evaluation, along with a description of the reasonably available information, conceptual model, analysis plan and science approaches, and plan for peer review for this chemical substance.

*General Information.* 1,2-Dichloropropane (CASRN 78-87-5) is a colorless, flammable liquid with a characteristic (chloroform-like) odor. It is moderately soluble in water and readily evaporates into air. It does not occur naturally in the environment.

**Reasonably Available Information.** EPA leveraged the data and information sources already described in the *Proposed Designation of 1,2-Dichloropropane (CASRN 78-87-5) as a High-Priority Substance for Risk Evaluation* (U.S. EPA, 2019e) to inform the development of this scope document. Furthermore, EPA conducted a comprehensive search to identify and screen multiple evidence streams (*i.e.*, chemistry, fate, release and engineering, exposure, hazard) and the search and screening results to date are provided in Section 2.1. EPA used the systematic review process described in Appendix A to search for and screen reasonably available information, including information already in EPA's possession, for inclusion in the risk evaluation. This information for 1,2-dichloropropane. EPA has focused on the data collection phase (consisting of data search, data screening, and data extraction) during the preparation of the scope document, whereas the data evaluation and integration stages will occur during the development of the risk evaluation and thus are not part of the scoping activities described in this document. EPA plans to consider additional information identified following publication of this scope document, as appropriate, in developing the risk evaluation, including the Chemical Data Reporting (CDR) information that the Agency will receive by the end of November 2020.

*Conditions of Use.* EPA plans to evaluate manufacturing (including importing), processing, distribution in commerce, industrial, commercial and consumer uses, and disposal of 1,2-dichloropropane in the risk evaluation. 1,2-Dichloropropane is manufactured (including imported) in the United States. The chemical is processed as a reactant, incorporated into formulation, mixture, or reaction products, and used industrially in non-incorporative activities. Commercial and consumer uses were identified in laboratory chemicals and cleaning and furnishing care products. EPA identified these conditions of use from information reported to EPA through CDR and Toxics Release Inventory (TRI) reporting, published literature, public comments, and consultation with stakeholders for both uses currently in production and uses whose production may have ceased. EPA did not revise any conditions of use in the final scope document for 1,2-dichloropane based on public comments received (Docket ID: <u>EPA-HQ-</u>

<u>OPPT-2018-0428</u>) on the draft scope. EPA is aware of information reporting use of 1,2-dichloropropane in pesticides; however, they are not conditions of use for the chemical substance as defined in TSCA § 3(2) and (4). Section 2.2.1 provides details about the conditions of use within the scope of the risk evaluation.

*Conceptual Models.* The conceptual models for 1,2-dichloropropane are presented in Section 2.6. Conceptual models are graphical depictions of the actual or predicted relationships of conditions of use, exposure pathways (*e.g.*, media), exposure routes (*e.g.*, inhalation, dermal, oral), hazards and receptors throughout the life cycle of the chemical substance. EPA considered reasonably available information as well as public comments received on the draft scope document for 1,2-dichloropropane in finalizing the exposure pathways, exposure routes, and hazards EPA plans to evaluate in the risk evaluation. As a result, EPA plans to focus the risk evaluation for 1,2-dichloropropane on the following exposures, hazards and receptors:

• *Exposures (Pathways and Routes), Receptors and PESS.* EPA plans to evaluate releases to the environment as well as human and environmental exposures resulting from the conditions of use of 1,2-dichloropropane that EPA plans to consider in risk evaluation. Exposures for 1,2-dichloropropane are discussed in Section 2.3. Additional information identified through the results of systematic review searches will also inform expected exposures.

EPA's plan for evaluating environmental exposure pathways in the scope of the risk evaluation considers whether and how other EPA-administered statutes and regulatory programs cover 1,2-dichloropropane in media pathways falling under the jurisdiction of those authorities. Section 2.6.3.1 discusses pathways under the jurisdiction of other EPA-administered laws. In Section 2.6.3.2, EPA presents the conceptual model describing the identified exposures (pathways and routes), receptors and hazards associated with the conditions of use of 1,2-dichloropropane within the scope of the risk evaluation. Based on pathways under the jurisdiction of other EPA-administered laws, EPA does not plan to evaluate the general population as a receptor in the scope of the risk evaluation (see Figure 2-16).

EPA considered reasonably available information and comments received on the draft scope for 1,2-dichloropropane in determining the human and environmental exposure pathways, routes, receptors and PESS for inclusion in the final scope. EPA plans to evaluate the following human and environmental exposure pathways, routes, receptors and PESS in the scope of the risk evaluation:

- *Occupational exposure:* EPA plans to evaluate exposures to workers and occupational non-users (ONUs) via the inhalation route and exposures to workers via the dermal route associated with manufacturing, processing, use or disposal of 1,2-dichloropropane.
- Consumer and bystander exposure: EPA plans to evaluate inhalation exposure to 1,2dichloropropane for consumers and bystanders, and dermal exposure to 1,2dichloropropane for consumers during the use of cleaning and furniture care products.
- *PESS:* EPA plans to include children, women of reproductive age (*e.g.*, pregnant women), workers and consumers as PESS in the risk evaluation.
- *Environmental exposure:* EPA plans to evaluate exposure to 1,2-dichloropropane for aquatic receptors.
- *Hazards*. Hazards for 1,2-dichloropropane are discussed in Section 2.4. EPA completed preliminary reviews of information (*e.g.*, federal and international government chemical

assessments) to identify potential environmental and human health hazards for 1,2dichloropropane as part of the prioritization (<u>U.S. EPA, 2019e</u>) and scoping process (<u>U.S. EPA, 2020c</u>). EPA also considered reasonably available information collected through systematic review methods as outlined in Appendix A and public comments received on the draft scope for 1,2-dichloropropane in determining the broad categories of environmental and human health hazard effects to be evaluated in the risk evaluation. EPA plans to use systematic review methods to evaluate the epidemiological and toxicological literature for 1,2-dichloropropane.

EPA plans to evaluate all potential environmental and human health hazard effects identified for 1,2-dichloropropane in Sections 2.4.1 and 2.4.2, respectively. Identified through the data screening phase of systematic review, the potential environmental hazard effects and related information that EPA plans to consider for the risk evaluation include: ADME, cancer, developmental, endocrine, gastrointestinal, hepatic, mortality, musculoskeletal, neurological, nutritional and metabolic and reproductive for 1,2-dichloropropane. Similarly, the potential human health hazard effects and related information identified through prioritization and the data screening phase of systematic review for 1,2-dichloropropane that EPA plans to consider for the risk evaluation include: ADME, PBPK, cancer, cardiovascular, developmental, endocrine, gastrointestinal, hematological and immune, hepatic, mortality, musculoskeletal, neurological, nutritional and metabolic, ocular and sensory, renal, reproductive, respiratory and skin and connective tissue.

*Analysis Plan.* The analysis plan for 1,2-dichloropropane is presented in Section 2.7. The analysis plan outlines the general science approaches that EPA plans to use for the various evidence streams (*i.e.*, chemistry, fate, release and engineering, exposure, hazard) supporting the risk evaluation. The analysis plan is based on EPA's knowledge of 1,2-dichloropropane to date which includes a review of information as described in Section 2.1. Should additional data or approaches become reasonably available, EPA may consider them for the risk evaluation.

*Peer Review.* The draft risk evaluation for 1,2-dichloropropane will be peer reviewed. Peer review will be conducted in accordance with relevant and applicable methods for chemical risk evaluations, including using EPA's Peer Review Handbook (U.S. EPA, 2015b) and other methods consistent with Section 26 of TSCA (see 40 CFR 702.45).

## **1 INTRODUCTION**

This document presents the scope of the risk evaluation to be conducted for 1,2-dichloropropane under the Frank R. Lautenberg Chemical Safety for the 21st Century Act. The Frank R. Lautenberg Chemical Safety for the 21st Century Act amended the TSCA on June 22, 2016. The new law includes statutory requirements and deadlines for actions related to conducting risk evaluations of existing chemicals.

Under TSCA § 6(b), the Environmental Protection Agency (EPA) must designate chemical substances as high-priority substances for risk evaluation or low-priority substances for which risk evaluations are not warranted at the time, and upon designating a chemical substance as a high-priority substance, initiate a risk evaluation on the substance. TSCA § 6(b)(4) directs EPA to conduct risk evaluations for existing chemicals, to "*determine whether a chemical substance presents an unreasonable risk of injury to health or the environment, without consideration of costs or other nonrisk factors, including an unreasonable risk to a potentially exposed or susceptible subpopulation identified as relevant to the risk evaluation by the Administrator under the conditions of use.*"

TSCA § 6(b)(4)(D) and implementing regulations require that EPA publish the scope of the risk evaluation to be conducted, including the hazards, exposures, conditions of use and PESS that the Administrator expects to consider, within 6 months after the initiation of a risk evaluation. In addition, a draft scope is to be published pursuant to 40 CFR 702.41. In December 2019, EPA published a list of 20 chemical substances that have been designated high priority substances for risk evaluations (Docket ID: EPA-HQ-OPPT-2018-0428) (84 FR 71924, December 30, 2019), as required by TSCA § 6(b)(2)(B), which initiated the risk evaluation process for those chemical substances. 1,2-Dichloropropane is one of the chemicals designated as a high priority substance for risk evaluation. On April 9, 2020, EPA published the *Draft Scope of the Risk Evaluation for 1,2-Dichloropropane* (EPA Document No. 740-D-20-006) (85 FR 19941, April 9, 2020) (U.S. EPA, 2020c) for a 45-day public comment period. After reviewing and considering the public comments received (Docket ID: EPA-HQ-OPPT-2018-0428) on the draft scope document, EPA is now publishing this final scope document pursuant to 40 CFR 702.41(c)(8).

### **2** SCOPE OF THE EVALUATION

### 2.1 Reasonably Available Information

EPA conducted a comprehensive search for reasonably available information<sup>1</sup> to support the development of this scope document for 1,2-dichloropropane. EPA leveraged the data and information sources already collected in the documents supporting the high-priority substance designations. In addition, EPA searched for additional data and information on physical and chemical properties, environmental fate, engineering, exposure, environmental and human health hazards that could be obtained from in the following general categories of sources:

1. Databases containing publicly available, peer-reviewed literature;

<sup>&</sup>lt;sup>1</sup>*Reasonably available information* means information that EPA possesses or can reasonably generate, obtain, and synthesize for use in risk evaluations, considering the deadlines specified in TSCA Section 6(b)(4)(G) for completing such evaluation. Information that meets the terms of the preceding sentence is reasonably available information whether or not the information is confidential business information, that is protected from public disclosure under TSCA Section 14 (40 CFR 702.33).

- 2. Gray literature, which is defined as the broad category of data/information sources not found in standard, peer-reviewed literature databases;
- 3. Data and information submitted under TSCA Sections 4, 5, 8(e), and 8(d), as well as "for your information" (FYI) submissions.

Following the comprehensive search, EPA performed a title and abstract screening to identify information potentially relevant for the risk evaluation process. This step also classified the references into useful categories or tags to facilitate the sorting of information through the systematic review process.

Search terms were used to search each of the literature streams and gather 1,2-dichloropropane studies. These terms and the methods used to develop them are listed in Appendix A. The studies resulting from the search process were loaded into the EPA Health and Environmental Research Online (HERO) database and then prioritized to screen first the literature likely relevant for each of the disciplines: fate, physical and chemical properties, engineering, exposure and hazard. The tools and methods used to manage the screening process are also outlined in Appendix A. The studies resulting from the search underwent a title/abstract screening process, which tagged them by topic or category. Following this, a determination was made to move studies forward into full-text screening. The criteria used in the screening process for each discipline are found in the population, exposure, comparator, outcome (PECO) statements listed in Appendix A. The screening process was conducted based on EPA's planning, execution and assessment activities outlined in Appendix A.


EPA has focused on the data collection phase (consisting of data search, data screening, and data extraction) during the preparation of the scope document, whereas the data evaluation and integration stages will occur during the development of the risk evaluation and thus are not part of the scoping activities described in this document.

The subsequent sections summarize the data collection activities completed to date for the general categories of sources and topic areas (or disciplines) using systematic review methods.

### 2.1.1 Search of Gray Literature

EPA surveyed the gray literature<sup>2</sup> and identified 77 search results relevant to EPA's risk evaluation needs for 1,2-dichloropropane. Appendix A.3.4 lists the gray literature sources that yielded 77 discrete data or information sources relevant to 1,2-dichloropropane. EPA further categorized the data and information into the various topic areas (or disciplines) supporting the risk evaluation (*e.g.*, physical and chemical properties, environmental fate, ecological hazard, human health hazard, exposure, engineering), and the breakdown is shown in Figure 2-1. EPA plans to consider additional reasonably available information from gray literature if it becomes available during the risk evaluation phase.

<sup>&</sup>lt;sup>2</sup> Gray literature is defined as the broad category of data/information sources not found in standard, peer-reviewed literature databases (e.g., PubMed and Web of Science). Gray literature includes data/information sources such as white papers, conference proceedings, technical reports, reference books, dissertations, information on various stakeholder websites, and other databases.



### Figure 2-1. Gray Literature Tags by Discipline for 1,2-Dichloropropane


The percentages across disciplines do not add up to 100%, as each source may provide data or information for various topic areas (or disciplines).

## 2.1.2 Search of Literature from Publicly Available Databases (Peer-Reviewed Literature)

EPA has begun the systematic review process and has conducted searching and screening of the reasonably available literature using the process outlined in Appendix A. This includes performing a comprehensive search of the reasonably available peer review literature on physical and chemical properties, environmental fate and transport, engineering (environmental release and occupational exposure), exposure (environmental, general population and consumer) and environmental and human health hazards of 1,2-dichloropropane. Eligibility criteria were applied in the form of PECO statements (see Appendix A). Included references met the PECO criteria, whereas excluded references did not meet the criteria (*i.e.*, not relevant), and supplemental material was considered as potentially relevant (see Appendix A.2). EPA plans to evaluate the reasonably available information identified for each discipline during the development of the risk evaluation.


EPA created literature inventory trees to graphically illustrate the flow of data and information sources following full-text screening (see Figure 2-2, Figure 2-3, Figure 2-5, Figure 2-7, Figure 2-9). EPA used the Health Assessment Workplace Collaborative (HAWC) tool to develop web-based literature inventory trees illustrating, through interactive links, studies that were included or excluded. These literature inventory trees enhance the transparency of the decisions resulting from the screening process described in Appendix A. For each of the corresponding disciplines, the literature was tagged to be included for evaluation during the risk evaluation. Literature inventory trees for physical and chemical properties are provided as static diagrams (Figure 2-2). For all other disciplines, static screen captures are provided in addition to links to the interactive trees. The links show individual studies that were tagged as included, excluded or supplemental. Supplemental studies did not meet all inclusion criteria but may be considered during risk evaluation as supporting information (see Appendix A). These studies can be accessed through the hyperlink provided in the associated caption below each figure. In some figures, the sum of the numbers for the various sub-categories may be larger than the broader category because some studies may be included under multiple sub-categories. In other cases, the sum of the various sub-categories may be smaller than the main category because some studies may not be depicted in the sub-categories if their relevance to the risk evaluation was unclear.

In addition, EPA tabulated the number and characteristics of the data and information sources included in the full-text screening process in the form of literature heat maps for the fate, engineering, exposure and hazard information (see Figure 2-4, Figure 2-6, Figure 2-8, and Figure 2-10). For each of these four disciplines, a static image of the literature inventory heat map is provided and a link to interactive version presented in HAWC is included in the caption below each diagram.




## Figure 2-2. Peer-reviewed Literature Inventory Tree - Physical and chemical Properties Search Results for 1,2-Dichloropropane

Data in this static figure represent references obtained from the publicly available databases search (see Appendix A.1.2) that were included during full-text screening as of June 2, 2020. TIAB refers to "title and abstract" screening.



## Figure 2-3. Peer-reviewed Literature Inventory Tree – Fate and Transport Search Results for 1,2-Dichloropropane

Click <u>here</u> to view the interactive literature inventory tree. Data in this figure represent references obtained from the publicly available databases search (see Appendix A.1.2) that were included during full-text screening as of June 2, 2020. Additional data may be added to the interactive version as they become available.



## Figure 2-4. Peer-reviewed Literature Inventory Heat Map – Fate and Transport Search Results for 1,2-Dichloropropane

Click <u>here</u> to view the interactive version for additional study details. The column totals, row totals, and grand totals indicate total numbers of unique references, as some references may be included in multiple cells. The various shades of color visually represent the number of relevant references identified by media or endpoint. The darker the color, the more references are available for a given media or endpoint. Data in this figure represents references obtained from the publicly available databases search (see Appendix A.1.2) that were included during full-text screening as of June 2, 2020. Additional data may be added to the interactive version as they become available.





Figure 2-5. Peer-reviewed Literature Inventory Tree - Engineering Search Results for 1,2-Dichloropropane

Click <u>here</u> to view the interactive literature inventory tree. Data in this figure represent references obtained from the publicly available databases search (see Appendix A.1.2) that were included during full-text screening as of August 5, 2020. Additional data may be added to the interactive version as they become available.

Data Type 💈	Evidence Tags	
	Description of release source	14
	Release frequency	1
Environmental	Release or emission factors	11
Releases	Release quantity	14
	Waste treatment methods and pollution control	7
	Total	25
	Chemical concentration	6
	Life cycle description	5
General	No evidence tag	5
Engineering	Number of sites	8
Assessment	Process description	8
Assessment	Production, import, or use volume	14
	Throughput	5
	Total	29
	Area sampling data	17
	Dermal exposure data	4
	Engineering control	3
	Exposure duration	9
	Exposure frequency	4
	Exposure route	19
Occupational	No evidence tag	5
Exposures	Number of workers	11
	Particle size characterization	
	Personal protective equipment	4
	Personal sampling data	12
	Physical form	10
	Worker activity description	18
	Total	37
Grand Total		54

#### Figure 2-6. Peer-reviewed Literature Inventory Heat Map - Engineering Search Results for 1,2-Dichloropropane

Click <u>here</u> to view the interactive version for additional study details. Data in this figure represent references obtained from the publicly available databases search (see Appendix A.1.2) that were included during full-text screening as of August 5, 2020. Additional data may be added to the interactive version as they become available.




#### Figure 2-7. Peer-Reviewed and Gray Literature Inventory Tree - Exposure Search Results for 1,2-Dichloropropane

Click <u>here</u> to view the interactive literature inventory tree. Data in this figure represent all references obtained from the publicly available databases search (see Appendix A.1.2), and gray literature references search (see Appendix A.3) that were included during full-text screening as of July 31, 2020. Additional data may be added to the interactive version as they become available.

				Data	а Туре			
Media (group)	Monitoring Study	Modeling Study	Completed Assessment	Experimental Study	Epidemiological Study	Database	Survey	Grand Tota
Ambient Air								
Biosolids/Sludge		1	1					1
Drinking Water								
Groundwater								
Land Disposal/ Landfill								
Sediment		1	2					2
Soil	3	1	2					5
Surface Water	1	1	2			1		5 3 1
Wastewater			1					1
Aquatic Species		1	2					2
Terrestrial Species								
Consumer	4	2	3	3				8
Dietary		1	1	1				2
Dust								
Exposure Factors	1	1	1	1				2
Exposure Pathway	2	2	4					2 5 3
Human Biomonitoring	3			1	1	1		3
Indoor Air	30	3	11	4	2		2	34
Isomers	2	1	1	1	1			34 2 4
Use Information	1	2	4					4
No Evidence Type	2	1	1		1	1	1	2
Grand Total	37	7	15	5	4	3	3	44

## Figure 2-8. Peer-reviewed and Gray Literature Inventory Heat Map –Exposure Search Results for 1,2-Dichloropropane

Click <u>here</u> to view the interactive version for additional study details. The column totals, row totals, and grand totals indicate total numbers of unique references only, as some references may be included in multiple cells. The various shades of color visually represent the number of relevant references identified by exposure media or data type. The darker the color, the more references are available for a given exposure media or data type. Data in this figure represent all references obtained from the publicly available databases search (see Appendix A.1.2), and gray literature references search (see Appendix A.3) that were included during full-text screening as of July 31, 2020. Additional data may be added to the interactive version as they become available.



## Figure 2-9. Peer-reviewed Literature Inventory Tree - Human Health and Environmental Hazards Search Results for 1,2-Dichloropropane

Click <u>here</u> to view the interactive literature inventory tree. Data in this figure represent references obtained from the publicly available databases search (see Appendix A.1.2.) that were included during full-text screening as of June 2, 2020. Additional data may be added to the interactive version as they become available.

Health Outcomes Human		Animal - Human Health Model	Animal - Environmental Model	Plant	Grand Total
ADME	6	12	1		18
Cancer	7	14	1		21
Cardiovascular	1	1			2
Developmental	2	11	3		15
Endocrine	2	7	1		9
Gastrointestinal	3	5	1		8
Hematological and Immune	1	11			12
Hepatic	2	18	1		20
Mortality	1	5	2		8
Musculoskeletal	1	6	1		8
Neurological	1	3	1		4
Nutritional and Metabolic	1	5	1		6
Ocular and Sensory	1	5			6
РВРК	1				1
Renal		7			7
Reproductive	2	9	1		12
Respiratory	3	10			13
Skin and Connective Tissue	3	1			4
No Tag	1	7	7		14
Grand Total	10	38	12		57

#### Evidence Type

## Figure 2-10. Peer-reviewed Literature Inventory Heat Map – Human Health and Environmental Hazards Search Results for 1,2-Dichloropropane

Click <u>here</u> to view the interactive version for additional study details. The numbers indicate the number of studies with TIAB keywords related to a particular health outcome, not the number of studies that observed an association with 1,2-dichloropropane. Evidence types were manually extracted, and Health Systems were determined via machine learning. Therefore, the studies examining multiple Health Outcomes and Evidence types, connections between health outcome, and evidence type may not be accurately represented. If a study evaluated multiple health outcomes or included multiple populations or study designs, it is shown here multiple times. Data in this figure represents references obtained from the publicly available databases search (see Appendix A.1.2.) that were included during full-text screening as of June 2, 2020. Additional data may be added to the interactive version as they become available.

### 2.1.3 Search of TSCA Submissions

Table 2-1 presents the results of screening the titles of data sources and reports submitted to EPA under various sections of TSCA. EPA screened a total of 127 submissions using PECO or similar statements that identify inclusion/exclusion criteria specific to individual disciplines (see Table 2-1 for the list of disciplines). The details about the criteria are presented in Appendix A.2.1. EPA identified 105 submissions that met the inclusion criteria in these statements

and identified 12 submissions with supplemental data.<sup>3</sup> EPA excluded 10 submissions because the reports were identified as one of the following:

- Published report that would be identified via other peer or gray literature searches
- Preliminary or draft report of a final available submitted report
- Ecosystem modeling study
- Memo regarding meeting
- Environmental impact statement for proposed equipment
- Ranking of chemicals for proposed evaluation

EPA plans to conduct additional deduplication at later stages of the systematic review process (*e.g.*, full text screening), when more information regarding the reports is available.

ntal <sup>b</sup>

Table 2-1. Results of Title Screening of Submissions to EPA under Various Sections of TSCA<sup>a</sup>

<sup>a</sup>Individual submissions may be relevant to multiple disciplines.

<sup>b</sup>Included submissions may contain supplemental data for other disciplines, which will be identified at full-text review.

### 2.2 Conditions of Use

As described in the *Proposed Designation of* 1,2-*Dichloropropane (CASRN* 78-87-5) *as a High-Priority Substance for Risk Evaluation* (U.S. EPA, 2019e), EPA assembled information from the CDR and TRI programs to determine conditions of use<sup>4</sup> or significant changes in conditions of use of the chemical substance. Once the 2020 CDR reporting period ends in November 2020, EPA plans to utilize the most recent CDR information. EPA also consulted a variety of other sources to identify uses of 1,2-dichloropropane, including: published literature, company websites, and government and commercial trade databases and publications. To identify formulated products containing 1,2-dichloropropane, EPA searched for safety data sheets (SDS) using internet searches, EPA Chemical and Product Categories (CPCat) (U.S. EPA, 2019d) data, and other resources in which SDSs could be found. SDSs were cross-checked with company websites to make sure that each product SDS was current. In addition, EPA incorporated communications with companies, industry groups, and public comments to supplement the use information.

EPA identified and described the categories and subcategories of conditions of use that EPA plans to consider in the risk evaluation (Section 2.2.1; Table 2-2). The conditions of use included in the scope of the risk evaluation are those reflected in the life cycle diagrams and conceptual models.

<sup>&</sup>lt;sup>3</sup> EPA may further consider some supplemental or excluded references depending on the reasons for tagging as supplemental or excluded.

<sup>&</sup>lt;sup>4</sup> *Conditions of use* means the circumstances, as determined by the Administrator, under which a chemical substance is intended, known, or reasonably foreseen to be manufactured, processed, distributed in commerce, used, or disposed of (TSCA § 3(4)).

After gathering reasonably available information related to the manufacture, processing, distribution in commerce, use, and disposal of 1,2-dichloropropane, EPA identified those activities for 1,2-dichloropropane the Agency determined not to be conditions of use or are otherwise excluded from the scope of the risk evaluation. These excluded activities are described in Section 2.2.2.

## 2.2.1 Categories and Subcategories of Conditions of Use Included in the Scope of the Risk Evaluation

Table 2-2 lists the conditions of use that are included in the scope of the risk evaluation.

 Table 2-2. Categories and Subcategories of Conditions of Use Included in the Scope of the Risk

 Evaluation

Life-Cycle Stage <sup>a</sup>	Category <sup>b</sup>	Subcategory <sup>c</sup>	References
Manufacturing	Domestic Manufacturing	Domestic Manufacturing	<u>U.S. EPA (2019b)</u>
	Import	Import	<u>U.S. EPA (2019b)</u>
Processing	As a reactant	Intermediate in all other basic organic chemical manufacturing	<u>U.S. EPA (2019b)</u>
	Incorporation into formulation, mixture, or reaction product	Intermediate in all other chemical product and preparation manufacturing	<u>U.S. EPA (2019b)</u>
Distribution in Commerce	Distribution in Commerce	Distribution in Commerce	
Industrial Use	Non-incorporative activities	Processing aids, not otherwise listed	EPA-HQ-OPPT-2018- 0428-0015
Commercial use	Other use	Laboratory chemicals	Chem Service (2018)
	Cleaning and furnishing care product	All-purpose liquid spray cleaner	<u>Tenax Spa (2015c)</u>
	Cleaning and furnishing care product	All-purpose waxes and polishes	<u>Tenax (2019); Tenax Spa</u> (2015a, 2015b)
	Cleaning and furnishing care product	All-purpose liquid cleaner/polish	<u>Axson USA (2015)</u>
Consumer use	Cleaning and furnishing care product	All-purpose liquid spray cleaner	<u>Tenax Spa (2015c)</u>

Life-Cycle Stage <sup>a</sup>	Category <sup>b</sup>	Subcategory <sup>c</sup>	References
	Cleaning and furnishing care product	All-purpose waxes and polishes	<u>Tenax (2019); Tenax Spa</u> (2015a, 2015b)
	Cleaning and furnishing care product	All-purpose liquid cleaner/polish	<u>Axson USA (2015)</u>
Disposal	Disposal	Disposal	

a. Life Cycle Stage Use Definitions (40 CFR § 711.3)

- "Industrial use" means use at a site at which one or more chemicals or mixtures are manufactured (including imported) or processed.
- "Commercial use" means the use of a chemical or a mixture containing a chemical (including as part of an article) in a commercial enterprise providing saleable goods or services.
- "Consumer use" means the use of a chemical or a mixture containing a chemical (including as part of an article, such as furniture or clothing) when sold to or made available to consumers for their use.
- Although EPA has identified both industrial and commercial uses here for purposes of distinguishing scenarios in this document, the Agency interprets the authority over "any manner or method of commercial use" under TSCA Section 6(a)(5) to reach both.
- b. These categories of conditions of use appear in the Life Cycle Diagram, reflect CDR codes, and broadly represent conditions of use of 1,2-dichloropropane in industrial and/or commercial settings.
- c. These subcategories reflect more specific conditions of use of 1,2-dichloropropane.

In the final scope EPA made the following changes to conditions of use:

 In the draft scope document, there are conditions of use for 1,2-dichloropropane for commercial and consumer use –cleaning and furniture care products – All-purpose liquid spray cleaner; waxes and polishes; liquid cleaner/polish. After checking the reporting categories under CDR, EPA has determined that the category for these conditions of use should be cleaning and furnishing care products.

#### 2.2.2 Activities Excluded from the Scope of the Risk Evaluation

As explained in the final rule, *Procedures for Chemical Risk Evaluation Under the Amended Toxic Substances Control Act* (82 FR 33726, July 20, 2017), TSCA Section 6(b)(4)(D) requires EPA to identify the hazards, exposures, conditions of use, and the PESS the Administrator expects to consider in a risk evaluation, suggesting that EPA may exclude certain activities that it determines to be conditions of use on a case-by-case basis (82 FR 33726, 33729; July 20, 2017). TSCA Section 3(4) also grants EPA discretion to determine the circumstances that are appropriately considered to be conditions of use for a particular chemical substance<sup>5</sup>. As a result, EPA does not plan to include in this scope or in the risk evaluation the activities described below that the Agency does not consider to be conditions of use or for which EPA is exercising discretionary authority provided by TSCA Section 6(b)(4)(D).

TSCA Section 3(2) also excludes from the definition of "chemical substance" "any food, food additive, drug, cosmetic, or device (as such terms are defined in Section 201 of the Federal Food, Drug, and Cosmetic Act [21 U.S.C. 321]) when manufactured, processed, or distributed in commerce for use as a food, food additive, drug, cosmetic, or device" as well as "any pesticide (as defined in the Federal Insecticide, Fungicide, and Rodenticide Act [7 U.S.C. 136 et seq.]) when manufactured, processed, or distributed in commerce for use as a pesticide." EPA has determined that the following uses of 1,2-dichloropropane are non-TSCA uses:

The National Library of Medicine's (NLM) hazardous substances data bank (HSDB) identifies use of 1,2-dichloropropane in insecticidal fumigant mixtures and former use as a soil fumigant for nematodes. NLM also identifies use of this chemical in dichloropropane-dichloropropene (D-D) mixture an obsolete soil fumigant mixture of three chemicals (a mixture of 1,2-dichloropropane, 1,3-dichloropropene and other chlorinated hydrocarbons) (NLM, 2001). CalDPR lists 16 inactive registrants (companies) for use of 1,2-dichloropropane in pesticides (CDPR, 2013). Ullmann's identifies use of 1,2-dichloropropane as an insecticidal and nematocidal fumigant, and in mixtures with other pesticides (Hartwig, 2008). EPA has determined that this use falls outside TSCA's definition of "chemical substance." Under TSCA § 3(2)(B)(vi), the definition of "chemical substance" does not include any pesticide (as the term is defined by EPA's Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), 7 U.S.C. § 136 et seq. (1996)). Activities and releases associated with such pesticidal uses are therefore not "conditions of use," and EPA does not plan to evaluate them during risk evaluation.

### 2.2.3 Production Volume

As reported to EPA during the 2016 CDR reporting period and described here as a range to protect production volumes that were claimed as confidential business information (CBI), total production volume of 1,2-dichloropropane in 2015 was between 100 million and 1 billion pounds (U.S. EPA,

<sup>&</sup>lt;sup>5</sup> *Chemical substance* means any organic or inorganic substance of a particular molecular identity, including any combination of such substances occurring in whole or in part as a result of a chemical reaction or occurring in nature, and any element or uncombined radical. Chemical substance does not include (1) any mixture; (2) any pesticide (as defined in the Federal Insecticide, Fungicide, and Rodenticide Act) when manufactured, processed, or distributed in commerce for use as a pesticide; (3) tobacco or any tobacco product; (4) any source material, special nuclear material, or byproduct material (as such terms are defined in the Atomic Energy Act of 1954 and regulations issued under such Act); (5) any article the sale of which is subject to the tax imposed by Section 4181 of the Internal Revenue Code of 1954 (determined without regard to any exemptions from such tax provided by Section 4182 or 4221 or any other provision of such Code), and; (6) any food, food additive, drug, cosmetic, or device (as such terms are defined in Section 201 of the Federal Food, Drug, and Cosmetic Act) when manufactured, processed, or distributed in commerce for use as a food, food additive, drug, cosmetic, or device (TSCA § 3(2)).

2020a). EPA also uses pre-2015 CDR production volume information, as detailed in the *Proposed Designation of 1,2-Dichloropropane (CASRN 78-87-5) as a High-Priority Substance for Risk Evaluation* (U.S. EPA, 2019e) and will include more recent production volume information from the 2020 CDR reporting period in the risk evaluation to support the exposure assessment.

### 2.2.4 Overview of Conditions of Use and Lifecycle Diagram

Figure 2-11 provides the lifecycle diagram for 1,2-dichloropropane. The life cycle diagram is a graphical representation of the various life stages of the industrial, commercial and consumer use categories included within the scope of the risk evaluation. The information in the life cycle diagram is grouped according to the CDR processing codes and use categories (including functional use codes for industrial uses and product categories for industrial, commercial and consumer uses). Appendix E contains additional descriptions (*e.g.*, process descriptions, worker activities, process flow diagrams) for each manufacture, processing, distribution in commerce, use and disposal category.




Figure 2-11. 1,2-Dichloropropane Life Cycle Diagram

### 2.3 Exposures

For TSCA exposure assessments, EPA plans to analyze human and environmental exposures and releases to the environment resulting from the conditions of use within the scope of the risk evaluation for 1,2-dichloropropane. In this section, the physical and chemical properties, environmental fate and transport properties and releases to the environment are described in addition to potential human and environmental exposures from TSCA conditions of use and from other possible or known sources. Release pathways and routes will be described in Section 2.6 to characterize the relationship or connection between the conditions of use of the chemical and the exposure to human receptors, including PESS, and environmental receptors. EPA plans to consider, where relevant, the duration, intensity (concentration), frequency, and number of exposures in characterizing exposures to 1,2-dichloropropane.

#### 2.3.1 Physical and Chemical Properties

Consideration of physical and chemical properties is essential for a thorough understanding or prediction of environmental fate (*i.e.*, transport and transformation) and the eventual environmental concentrations. It can also inform the hazard assessment. Table 2-3 summarizes the physical and chemical property values preliminarily selected for use in the risk evaluation from among the range of reported values collected as of June 2020. This information differs from that presented in the *Proposed Designation of 1,2-Dichloropropane (CASRN 78-87-5) as a High-Priority Substance for Risk Evaluation* (U.S. EPA, 2019e) and may be updated as EPA continues to evaluate and integrate additional information through systematic review methods. Figure 2-12 summarizes the distribution of reported values for eight physical and chemical properties routinely used in existing chemical risk evaluations. Appendix A presents summary statistics for reported physical and chemical property values. All physical and chemical property values that were extracted and evaluated as of June 2020 are presented in the supplemental file *Data Extraction and Data Evaluation Tables for Physical and Chemical Property Studies* (EPA-HQ-OPPT-2018-0428).

Property or Endpoint	Value <sup>a</sup>	Reference	Data Quality Rating
Molecular formula	C <sub>3</sub> H <sub>6</sub> Cl <sub>2</sub>	NA	NA
Molecular weight	112.99 g/mol	NA	NA
Physical state	Liquid	<u>RSC (2019)</u>	High
Physical properties	Colorless liquid, chloroform-like odor	<u>NLM (2019)</u>	High
Melting point	-100.53°C	<u>Rumble (2018)</u>	High
Boiling point	96.4°C	<u>Rumble (2018)</u>	High
Density	1.159 g/cm <sup>3</sup> at 25°C	<u>NLM (2019)</u>	High
Vapor pressure	40.00 at 20.5°C (99.95 mol%)	<u>Steele et al. (1997)</u>	High
Vapor density	3.9 (air = 1)	<u>NLM (2019)</u>	High

Table 2-3. Physical and Chemical Properties of 1,2-Dichloropropane

Property or Endpoint	Value <sup>a</sup>	Reference	Data Quality Rating
Water solubility	2800 mg/L at 25°C	<u>NLM (2019)</u>	High
Octanol/water partition coefficient (log Kow)	1.99 at 25°C	Elsevier (2019)	High
Henry's Law constant	0.00282 atm·m³/mol at 25°C	<u>U.S. EPA (2019c)</u>	High
Flash point	21°C (ASTM open cup)	<u>O'Neil (2013)</u>	High
Auto flammability	Not available		
Viscosity	Not available		
Refractive index	1.43656 at 25°C	Sen and Kim (2009)	High
Dielectric constant	8.92	Elsevier (2019)	High

<sup>a</sup> Measured unless otherwise noted.

NA = Not applicable

Figure 2-12 displays a summary of the data collected as of June 2020 for eight physical and chemical values routinely used in TSCA existing chemical risk evaluations. The box and whisker plots for each endpoint illustrate the mean (average, indicated by the blue diamond) and the 10<sup>th</sup>, 25<sup>th</sup>, 50<sup>th</sup> (median), 75<sup>th</sup>, and 90<sup>th</sup> percentiles. All individual data points are indicated by black squares, and value preliminarily selected for use in the risk evaluation is overlaid (indicated by the orange circle) to provide context for where it lies within the distribution of the dataset. The number of unique primary data sources is indicated below each box and whisker plot. If multiple sources presented equivalent values and cited the same primary source, only one of those was included in the statistical calculations. As a result, the number of sources listed in Figure 2-12 may differ from the total number of data sources presented in Figure 2-2. Where no data could be identified through systematic review, text appears to clearly demonstrate the gap for the endpoint.




Figure 2-12. Box and Whisker Plots of Reported Physical and Chemical Property Values

### 2.3.2 Environmental Fate and Transport

Understanding of environmental fate and transport processes assists in the determination of the specific exposure pathways and potential human and environmental receptors that need to be assessed in the risk evaluation for 1,2-dichloropropane. EPA plans to use the environmental fate characteristics described in Appendix C to support the development of the risk evaluation for 1,2dichloropropane. The values for the environmental fate properties may be updated as EPA evaluates and integrates additional information into the risk evaluation through systematic review methods.

### 2.3.3 Releases to the Environment

Releases to the environment from conditions of use are a component of potential exposure and may be derived from reported data that are obtained through direct measurement, calculations based on empirical data and/or assumptions and models.

A source of information that EPA plans to consider in the risk evaluation in evaluating exposure is data reported under the TRI program. EPA's TRI database contains information on chemical waste management activities that are disclosed by industrial and federal facilities, including quantities released

into the environment (*i.e.*, to air, water, and disposed of to land), treated, burned for energy, recycled, or transferred off-site to other facilities for these purposes.

Under the Emergency Planning and Community Right-to-Know Act (EPCRA) Section 313, 1,2dichloropropane is a TRI-reportable substance effective January 1, 1987 (40 CFR 372.65). For TRI reporting<sup>6</sup>, facilities in covered sectors in the United States are required to disclose releases and other waste management activity quantities of 1,2-dichloropropane under the CASRN 78-87-5 if they manufacture (including import) or process more than 25,000 pounds or otherwise use more than 10,000 pounds of the chemical in a given year by July 1 of the following year.

Table 2-4 provides production-related waste management data for 1,2-dichloropropane reported by facilities to the TRI program for reporting year 2018.<sup>7</sup> As shown in the table, 13 facilities reported a total of approximately 19.4 million pounds of 1,2-dichloropropane waste managed. Of this total, over 10 million pounds were burned for energy recovery, over 8.5 million pounds were recycled, nearly 755,000 pounds were treated, and nearly 18,000 pounds were released to the environment. Nearly all 1,2-dichloropropane burned for energy recovery was burned on site; only 3,100 pounds were sent off site for energy recovery. Similarly, most recycling and treatment occurred on site, with about 8 million pounds recycled on site and approximately 700,000 pounds treated on site. Overall, 96% of 1,2-dichloropropane waste managed was managed on site.

Year	Number of Facilities	Recycled (lbs)	Recovered for Energy (lbs)	Treated (lbs)	Released <sup>a,b,c</sup> (lbs)	Total Production Related Waste (lbs)
2018	13	8,546,388	10,034,130	754,885	17,973	19,353,377

Table 2-4. Summary of 1,2-Dichloropropane TRI Production-Related Waste Managed in 2018

Data source: U.S. EPA (2019f) (Updated November 2019)

<sup>a</sup> Terminology used in these columns may not match the more detailed data element names used in the TRI public data and analysis access points. <sup>b</sup> Does not include releases due to one-time event not associated with production such as remedial actions or earthquakes.

<sup>c</sup> Counts all releases including release quantities transferred and release quantities disposed of by a receiving facility reporting to TRI.

Table 2-5 provides a summary of 1,2-dichloropropane TRI releases to the environment for the same reporting year as Table 2-4.<sup>7</sup> Nearly 18,000 pounds of 1,2-dichloropropane were released to the environment in 2018; the 16,725 pounds released to the air accounted for 93% of all releases. Point source and fugitive air releases contributed roughly equally to air emissions with about 8,100 and 8,600 pounds, respectively. All other contributions to releases were small, with roughly 300 pounds discharged to surface waters on site, and 861 pounds disposed of to land (481 pounds disposed on-site to landfills, and 380 pounds disposed off-site to landfills). Overall, 97% of 1,2-dichloropropane releases occurred on site.

<sup>&</sup>lt;sup>6</sup> For TRI reporting criteria see <u>https://www.epa.gov/toxics-release-inventory-tri-program/basics-tri-reporting</u>.

<sup>&</sup>lt;sup>7</sup> Reporting year 2018 is the most recent TRI data available. Data presented in Table 2-4 were queried using TRI Explorer and uses the 2018 National Analysis data set (released to the public in November 2019). This dataset includes revisions for the years 1988 to 2018 processed by EPA.

		Air Re	eleases			Land Dispos	al		
	Number of Facilities	Stack Air Releases (lbs)	Fugitive Air Releases (lbs)	Water Releases (lbs)	Class I Under- ground Injection (lbs)	RCRA Subtitle C Landfills (lbs)	All other Land Disposal <sup>a</sup> (lbs)	Other Releases <sup>a</sup> (lbs)	Total Releases <sup>b, c</sup> (lbs)
Totals 2018	13	8,113	8,612	303.878	0	0	861	83.16	17,973
		16,	725			861			

Table 2-5. Summary of Releases of 1,2-Dichloropropane to the Environment During 2018

Data source: U.S. EPA (2019f) (Updated November 2019)

<sup>a</sup> Terminology used in these columns may not match the more detailed data element names used in the TRI public data and analysis access points.

<sup>b</sup> These release quantities do include releases due to one-time events not associated with production such as remedial actions or earthquakes.

Counts release quantities once at final disposition, accounting for transfers to other TRI reporting facilities that ultimately dispose of the chemical waste.

While production-related waste managed shown in Table 2-4 excludes any quantities reported as catastrophic or one-time releases (TRI Section 8 data), release quantities shown in Table 2-5 include both production-related and non-production-related quantities. For 1,2-dichloropropane the total release quantities shown in the two tables are the same, but for other TRI chemicals the total release quantities may differ slightly and may further reflect differences in TRI calculation methods for reported release range estimates (U.S. EPA, 2017).

EPA plans to review these data in conducting the exposure assessment component of the risk evaluation for 1,2-dichloropropane.

### 2.3.4 Environmental Exposures

The manufacturing, processing, distribution, use and disposal of 1,2-dichloropropane can result in releases to the environment and exposure to aquatic and terrestrial receptors (biota). Environmental exposures to biota are informed by releases into the environment, overall persistence, degradation, and bioaccumulation, and partitioning across different media. Concentrations of chemical substances in biota provide evidence of exposure.

EPA plans to review reasonably available environmental monitoring data for 1,2-dichloropropane. EPA's Ambient Monitoring Technology Information Center Air Toxics database has identified 1,2dichloropropane in air (U.S. EPA, 1990). In addition, EPA's Unregulated Contaminant Monitoring Rule has identified 1,2-dichloropropane in drinking water (U.S. EPA, 1996). USGS's Monitoring Data – National Water Quality Monitoring Council has identified 1,2-dichloropropane in ground water, sediment, soil, surface water and ecological tissue (*e.g.*, fish tissue concentrations) (USGS, 1991a, b, c, d, e, f, g). Existing assessments reported that 1,2-dichloropropane appears to be stable and present in the air, soil, surface water, and groundwater (RIVM, 2007; OECD, 2006; CalEPA, 1999).

In addition, 1,2-dichloropropane can be introduced to the environment as an impurity of 1,3dichloropropane. Until the early 1980s, 1,3-dichloropropane was used as a fumigant in soil and in grain crops, where most of the chemical was released to the air or groundwater where breakdown is slow (U.S. EPA, 2016b; ATSDR, 1989).

### 2.3.5 Occupational Exposures

EPA plans to evaluate worker activities where there is a potential for exposure under the various conditions of use (manufacturing, processing, industrial/commercial uses, and disposal) described in

Section 2.2. In addition, EPA plans to evaluate exposure to occupational non-users (ONUs), *i.e.*, workers who do not directly handle the chemical but perform work in an area where the chemical is present. EPA also plans to consider the effect(s) that engineering controls (EC) and/or personal protective equipment (PPE) have on occupational exposure levels as part of the risk evaluation.

Examples of worker activities associated with the conditions of use within the scope of the risk evaluation for 1,2-dichloropropane that EPA may analyze include, but are not limited to:

- Unloading and transferring 1,2-dichloropropane to and from storage containers and process vessels;
- Handling and disposing of waste containing 1,2-dichloropropane;
- Cleaning and maintaining equipment;
- Sampling chemicals, formulations, or products containing 1,2-dichloropropane for quality control;
- Repackaging chemicals, formulations, or products containing 1,2-dichloropropane;
- Performing other work activities in or near areas where 1,2-dichloropropane is used.

1,2-Dichloropropane is a liquid with vapor pressure of 53.3 mmHg at 25°C (see Appendix A). Based on the chemical's high volatility, EPA plans to analyze inhalation exposure to vapor for workers and occupational non-users (ONUs). EPA also plans to analyze inhalation exposure to mists for workers and ONUs where products containing 1,2-dichloropropane may be spray-applied. EPA plans to evaluate dermal exposures for workers, who are expected to have skin contact with 1,2-dichloropropane. Occupational non-users do not directly handle 1,2-dichloropropane; therefore, skin contact with 1,2-dichloropropane is not expected for occupational non-users.

EPA generally does not evaluate occupational exposures through the oral route. Workers and ONUs may inadvertently ingest inhaled particles that deposit in the upper respiratory tract. In addition, workers may transfer chemicals from their hands to their mouths. The frequency and significance of this exposure route are dependent on several factors including the physical and chemical properties of the substance during worker activities, the visibility of the chemicals on the hands while working, workplace training and practices, and personal hygiene that is difficult to predict (Cherrie et al., 2006). EPA plans to consider the relevance of this exposure route on a case-by-case basis, taking into consideration the aforementioned factors and any reasonably available information, and may assess oral exposure for workers for certain COUs and worker activities where warranted.

The United States has several regulatory and non-regulatory exposure limits of 1,2-dichloropropane: the Occupational Safety and Health Administration (OSHA) Permissible Exposure Limit (PEL) of 75 ppm or 350 mg/m<sup>3</sup> over an 8-hour workday, time weighted average (TWA). This chemical also has an American Conference of Governmental Industrial Hygienists (ACGIH) sets the Threshold Limit Value (TLV) at 10 ppm TWA (OSHA, 2009). NIOSH considers 1,2-dichloropropane to be a potential occupational carcinogen with an Immediately Dangerous to Life or Health (IDLH) value of 400 ppm (NIOSH, 2020, 2016).

#### 2.3.6 Consumer Exposures

No consumer conditions of use information for 1,2-dichloropropane was found in the 2012 or 2016 CDR. Material safety data sheets indicated that the 1,2-dichloropropane is found in cleaning and furniture care products, including all-purpose liquid spray cleaner, all-purposes waxes and polishes and all-purpose liquid cleaners and polishes. In addition, the Consumer Product Database indicated that 1,2-dichloropropane in used as a consumer product in cleaners (U.S. EPA, 2019a).

Consumers using or disposing of cleaning and furniture care products may be exposed to 1,2dichloropropane through direct liquid contact which may lead to a dermal exposure, or through vapor emissions which may lead to inhalation exposure, given its high volatility at room temperature. Bystanders present during the consumer use of cleaning and furniture care products, or disposal of 1,2dichloropropane may also be exposed to vapor emissions leading to an inhalation exposure.

Based on these potential sources and pathways of exposure, EPA plans to analyze inhalation and dermal routes of exposures to consumers that may result from the conditions of use of 1,2-dichloropropane. EPA does not plan to evaluate consumer exposures to 1,2-dichloropropane via the oral route as the exposure is not expected.

#### 2.3.7 General Population Exposures

Releases of 1,2-dichloropropane from certain conditions of use, such as manufacturing and processing activities, may result in general population exposure through breathing contaminated ambient air or consuming contaminated drinking water (<u>CalEPA, 1999</u>). Populations living near industrial wastewater treatment or incineration facilities may have higher exposure to 1,2-dichloropropane (<u>CalEPA, 1999</u>).

The OECD monitoring database has identified human biomonitoring data for 1,2-dichloropropane (OECD, 2018). However, blood concentrations of 1,2-dichloropropane were below the limit of detection in the 2,740 individuals who participated in the National Health and Nutrition Examination Survey (NHANES) 2011–2012 subsample of the U.S. population (CDC, 2018). The general population pathways in the scope of this evaluation are described in Sections 2.6.3 and 2.7.2.5.

### 2.4 Hazards (Effects)

#### 2.4.1 Environmental Hazards

EPA considered reasonably available information (*e.g.*, federal and international government chemical assessments) on 1,2-dichloropropane as well as public comments received on the *Proposed Designation of 1,2-Dichloropropane (CASRN 78-87-5) as a High-Priority Substance for Risk Evaluation* (U.S. EPA, 2019e) and draft scope for 1,2-dichloropropane (U.S. EPA, 2020c) to identify potential environmental hazards. During prioritization, EPA identified environmental hazard effects for aquatic and terrestrial organisms.

Since prioritization, EPA applied automated techniques during the data screening phase of systematic review to identify the following potential environmental hazards and related information that may be considered for the risk evaluation (as explained in Appendix A): ADME, cancer, developmental, endocrine, gastrointestinal, hepatic, mortality, musculoskeletal, neurological, nutritional and metabolic

and reproductive (

			Endence Type		
Health Outcomes	Human	Animal - Human Health Model	Animal - Environmental Model	Plant	Grand Total
ADME	6	12	1		18
Cancer	7	14	1		21
Cardiovascular	1	1			2
Developmental	2	11	3		15
Endocrine	2	7	1		9
Gastrointestinal	3	5	1		8
Hematological and Immune	1	11			12
Hepatic	2	18	1		20
Mortality	1	5	2		8
Musculoskeletal	1	6	1		8
Neurological	1	3	1		4
Nutritional and Metabolic	1	5	1		6
Ocular and Sensory	1	5			6
РВРК	1				1
Renal		7			7
Reproductive	2	9	1		12
Respiratory	3	10			13
Skin and Connective Tissue	3	1			4
No Tag	1	7	7		14
Grand Total	10	38	12		57

Evidence Type

Figure 2-10). A summary of references identified during the screening step of systematic review is included in the interactive literature inventory trees (Figure 2-9). As EPA continues to evaluate reasonably available and relevant hazard information identified through systematic review, EPA may update the list of potential hazard effects to be analyzed in the risk evaluation.

#### 2.4.2 Human Health Hazards

EPA considered reasonably available information (*e.g.*, federal and international government chemical assessments) on 1,2-dichloropropane as well as public comments on the *Proposed Designation of 1,2-Dichloropropane (CASRN 78-87-5) as a High-Priority Substance for Risk Evaluation* (U.S. EPA, 2019e) and draft scope for 1,2-dichloropropane (U.S. EPA, 2020c) to identify potential human health hazards. During prioritization, EPA identified the following potential human health hazards and related information: acute, repeat dose, genetic, reproductive, toxicokinetic, developmental, irritation/corrosion, dermal sensitization, cancer and neurological effects.

Since prioritization, EPA applied automated techniques during the data screening phase of systematic review to identify the following additional potential human health hazards and related information that may be considered for the risk evaluation (as explained in Appendix A): ADME, PBPK, cardiovascular, endocrine, gastrointestinal, hematological and immune, hepatic, mortality, musculoskeletal, nutritional

and metabolic, ocular and sensory, renal, respiratory and skin and connective tissue (

Health Outcomes	Human	Animal - Human Health Model	Animal - Environmental Model	Plant	Grand Total
ADME	6	12	1		18
Cancer	7	14	1		21
Cardiovascular	1	1			2
Developmental	2	11	3		15
Endocrine	2	7	1		9
Gastrointestinal	3	5	1		8
Hematological and Immune	1	11			12
Hepatic	2	18	1		20
Mortality	1	5	2		8
Musculoskeletal	1	6	1		8
Neurological	1	3	1		4
Nutritional and Metabolic	1	5	1		6
Ocular and Sensory	1	5			6
РВРК	1				1
Renal		7			7
Reproductive	2	9	1		12
Respiratory	3	10			13
Skin and Connective Tissue	3	1			4
No Tag	1	7	7		14
Grand Total	10	38	12		57

Evidence Type

Figure 2-10). A summary of the references identified during the screening step of systematic review is included in the interactive literature inventory trees (Figure 2-9). As EPA continues to evaluate reasonably available and relevant hazard information identified through systematic review, EPA may update the list of potential hazard effects to be analyzed in the risk evaluation.

### 2.5 Potentially Exposed or Susceptible Subpopulations

TSCA § 6(b)(4) requires EPA to determine whether a chemical substance presents an unreasonable risk to "a potentially exposed or susceptible subpopulation identified as relevant to the risk evaluation." TSCA §3(12) states that "the term 'potentially exposed or susceptible subpopulation' means a group of individuals within the general population identified by the Administrator who, due to either greater susceptibility or greater exposure, may be at greater risk than the general population of adverse health effects from exposure to a chemical substance or mixture, such as infants, children, pregnant women, workers, or the elderly." General population is "the total of individuals inhabiting an area or making up a whole group" and refers here to the U.S. general population (U.S. EPA, 2011a).

EPA identified the following PESS based on CDR information and studies reporting developmental and reproductive effects: children, women of reproductive age (*e.g.*, pregnant women), workers, including ONUs and users, and consumers, including users and bystanders (U.S. EPA, 2019e). EPA plans to evaluate these PESS in the risk evaluation. Following further evaluation of the reasonably available information, EPA may evaluate PESS in the general population as they relate to fence line communities.

In developing exposure scenarios, EPA plans to analyze reasonably available data to ascertain whether some human receptor groups may be exposed via exposure pathways that may be distinct to a particular subpopulation or life stage (*e.g.*, children's crawling, mouthing or hand-to-mouth behaviors) and whether some human receptor groups may have higher exposure via identified pathways of exposure due to unique characteristics (*e.g.*, activities, duration or location of exposure) when compared with the general population (U.S. EPA, 2006b). Likewise, EPA plans to evaluate available human health hazard information to ascertain whether some human receptor groups may have greater susceptibility than the general population to the chemical's hazard(s). Based on these analyses, EPA may update the list of PESS in the risk evaluation.


### 2.6 Conceptual Models

In this section, EPA presents the conceptual models describing the identified exposures (pathways and routes), receptors and hazards associated with the conditions of use of 1,2-dichloropropane. Pathways and routes of exposure associated with workers and occupational non-users are described in Section 2.6.1, and pathways and routes of exposure associated with environmental releases and wastes, including those pathways that are under the jurisdiction of other EPA-administered laws, are discussed and depicted in the conceptual model shown in Section 2.6.3.1. Pathways and routes of exposure associated with environmental releases and wastes, excluding those pathways that are under the jurisdiction of other EPA-administered laws, are discussed with environmental releases and wastes, excluding those pathways that are under the jurisdiction of other EPA-administered laws, are discussed and depicted in the conceptual model shown in Section 2.6.3.1. Pathways that are under the jurisdiction of other EPA-administered laws, are discussed and depicted in the conceptual model shown in Section 2.6.3.1. Pathways that are under the jurisdiction of other EPA-administered laws, are discussed and depicted in the conceptual model shown in Section 2.6.3.2.

#### 2.6.1 Conceptual Model for Industrial and Commercial Activities and Uses

Figure 2-13 illustrates the conceptual model for the pathways of exposure from industrial and commercial activities and uses of 1,2-dichloropropane that EPA plans to include in the risk evaluation. There is potential for exposures to workers and ONUs via inhalation routes and exposures to workers via dermal routes. It is expected that inhalation exposure to vapors is the most likely exposure route. In addition, workers at waste management facilities may be exposed via inhalation or dermal routes from wastewater treatment, incineration or via other disposal methods. EPA plans to evaluate activities resulting in exposures associated with distribution in commerce (*e.g.*, loading, unloading) throughout the various lifecycle stages and conditions of use (*e.g.*, manufacturing, processing, industrial use, commercial use, and disposal) rather than a single distribution scenario.

For each condition of use identified in Table 2-2, a determination was made as to whether or not EPA plans to evaluate each combination of exposure pathway, route, and receptor in the risk evaluation. The results of that analysis along with the supporting rationale are presented in Appendix F.





### Figure 2-13. 1,2-Dichloropropane Conceptual Model for Industrial and Commercial Activities and Uses: Worker and Occupational Non-User Exposures and Hazards

The conceptual model presents the exposure pathways, exposure routes and hazards to human receptors from industrial and commercial activities and uses of 1,2-dichloropropane.

### 2.6.2 Conceptual Model for Consumer Activities and Uses: Potential Exposures and Hazards

The conceptual model in Figure 2-14 presents the exposure pathways, exposure routes and hazards to human receptors from consumer activities and uses of 1,2-dichloropropane. EPA expects inhalation to be the primary route of exposure and plans to analyze inhalation exposures to 1,2-dichloropropane for consumers and bystanders. There is potential for dermal exposures to 1,2-dichloropropane via direct contact with liquid during consumer uses. Bystanders are not expected to have direct dermal contact to 1,2-dichloropropane. EPA plans to evaluate direct dermal contact with liquid 1,2-dichloropropane for consumers using cleaning and furniture care products. The supporting rationale for consumer pathways considered for 1,2-dichloropropane are included in Appendix G.






The conceptual model presents the exposure pathways, exposure routes and hazards to human receptors from consumer activities and uses of 1,2-dichloropropane. Note:

a) Receptors include PESS (see Section 2.5).

### 2.6.3 Conceptual Model for Environmental Releases and Wastes: Potential Exposures and Hazards (Regulatory Overlay)

In this section, EPA presents the conceptual models describing the identified exposures (pathways and routes from environmental releases and wastes) and hazards to general population and environmental receptors associated with the conditions of use of 1,2-dichloropropane within the scope of the risk evaluation. This section also discusses those pathways that may be addressed pursuant to other EPA-administered laws.

The conceptual model in Figure 2-15 presents the potential exposure pathways, exposure routes and hazards to general population and environmental receptors from releases and waste streams associated with industrial, commercial and consumer uses of 1,2-dichloropropane. The conceptual model shows the overlays, labeled and shaded to depict the regulatory programs under EPA-administered statutes and associated pathways that EPA considered for the scope of the risk evaluation. The regulatory programs that cover these environmental release and waste pathways are further described in Section 2.6.3.1.



# Figure 2-15. 1,2-Dichloropropane Conceptual Model for Environmental Releases and Wastes: Environmental and General Population Exposures and Hazards (Regulatory Overlay)

The conceptual model presents the exposure pathways, exposure routes and hazards to human and environmental receptors from releases and wastes from industrial, commercial, and consumer uses of 1,2-dichloropropane including the environmental statutes covering those pathways.

- a) Industrial wastewater or liquid wastes may be treated on-site and then released to surface water (direct discharge), or pre-treated and released to Publicly Owned Treatment Works (POTW) (indirect discharge). For consumer uses, such wastes may be released directly to POTW. Drinking water will undergo further treatment in drinking water treatment plant. Ground water may also be a source of drinking water. Inhalation from drinking water may occur via showering
- b) Receptors include PESS (see Section 2.5).

# 2.6.3.1 Exposure Pathways and Risks Addressed by Other EPA Administered Statutes

In its TSCA Section 6(b) risk evaluations, EPA is coordinating action on certain exposure pathways and risks falling under the jurisdiction of other EPA-administered statutes or regulatory programs. More specifically, EPA is exercising its TSCA authorities to tailor the scope of its risk evaluations, rather than focusing on environmental exposure pathways addressed under other EPA-administered statutes or regulatory programs or risks that could be eliminated or reduced to a sufficient extent by actions taken under other EPA-administered laws. EPA considers this approach to be a reasonable exercise of the Agency's TSCA authorities, which include:

- TSCA Section 6(b)(4)(D): "The Administrator shall, not later than 6 months after the initiation of a risk evaluation, publish the scope of the risk evaluation to be conducted, including the hazards, exposures, conditions of use, and the potentially exposed or susceptible subpopulations the Administrator expects to consider..."
- TSCA Section 9(b)(1): "The Administrator shall coordinate actions taken under this chapter with actions taken under other Federal laws administered in whole or in part by the Administrator. If the Administrator determines that a risk to health or the environment associated with a chemical substance or mixture could be eliminated or reduced to a sufficient extent by actions taken under the authorities contained in such other Federal laws, the Administrator shall use such authorities to protect against such risk unless the Administrator determines, in the Administrator's discretion, that it is in the public interest to protect against such risk by actions taken under this chapter."
- TSCA Section 9(e): "...[I]f the Administrator obtains information related to exposures or releases of a chemical substance or mixture that may be prevented or reduced under another Federal law, including a law not administered by the Administrator, the Administrator shall make such information available to the relevant Federal agency or office of the Environmental Protection Agency."
- TSCA Section 2(c): "It is the intent of Congress that the Administrator shall carry out this chapter in a reasonable and prudent manner, and that the Administrator shall consider the environmental, economic, and social impact of any action the Administrator takes or proposes as provided under this chapter."
- TSCA Section 18(d)(1): "Nothing in this chapter, nor any amendment made by the Frank R. Lautenberg Chemical Safety for the 21st Century Act, nor any rule, standard of performance, risk evaluation, or scientific assessment implemented pursuant to this chapter, shall affect the right of a State or a political subdivision of a State to adopt or enforce any rule, standard of performance, risk evaluation, scientific assessment, or any other protection for public health or the environment that- (i) is adopted or authorized under the authority of any other Federal law or adopted to satisfy or obtain authorization or approval under any other Federal law..."

These TSCA authorities supporting tailored risk evaluations and intra-agency referrals are described in more detail below:

#### TSCA Section 6(b)(4)(D)

TSCA Section 6(b)(4)(D) requires EPA, in developing the scope of a risk evaluation, to identify the hazards, exposures, conditions of use, and potentially exposed or susceptible subpopulations the Agency

"expects to consider" in a risk evaluation. This language suggests that EPA is not required to consider all conditions of use, hazards, or exposure pathways in risk evaluations. As EPA explained in the "Procedures for Chemical Risk Evaluation Under the Amended Toxic Substances Control Act" ("Risk Evaluation Rule"), "EPA may, on a case-by-case basis, tailor the scope of the risk evaluation in order to focus its analytical efforts on those exposures that are likely to present the greatest concern, and consequently merit an unreasonable risk determination." 82 FR 33726, 33729 (July 20, 2017).

In the problem formulation documents for many of the first 10 chemicals undergoing risk evaluation, EPA applied the same authority and rationale to certain exposure pathways, explaining that "EPA is planning to exercise its discretion under TSCA 6(b)(4)(D) to focus its analytical efforts on exposures that are likely to present the greatest concern and consequently merit a risk evaluation under TSCA, by excluding, on a case-by-case basis, certain exposure pathways that fall under the jurisdiction of other EPA-administered statutes." This is informed by the legislative history of the amended TSCA, which supports the Agency's exercise of discretion to focus the risk evaluation on areas that raise the greatest potential for risk. See June 7, 2016 Cong. Rec., S3519-S3520. Consistent with the approach articulated in the problem formulation documents, and as described in more detail below, EPA is exercising its authority under TSCA to tailor the scope of exposures evaluated in TSCA risk evaluations, rather than focusing on environmental exposure pathways addressed under other EPA-administered, media-specific statutes and regulatory programs.

#### TSCA Section 9(b)(1)

In addition to TSCA Section 6(b)(4)(D), the Agency also has discretionary authority under the first sentence of TSCA Section 9(b)(1) to "coordinate actions taken under [TSCA] with actions taken under other Federal laws administered in whole or in part by the Administrator." This broad, freestanding authority provides for intra-agency coordination and cooperation on a range of "actions." In EPA's view, the phrase "actions taken under [TSCA]" in the first sentence of Section 9(b)(1) is reasonably read to encompass more than just risk management actions, and to include actions taken during risk evaluation as well. More specifically, the authority to coordinate intra-agency actions exists regardless of whether the Administrator has first made a definitive finding of risk, formally determined that such risk could be eliminated or reduced to a sufficient extent by actions taken under authorities in other EPA-administered Federal laws, and/or made any associated finding as to whether it is in the public interest to protect against such risk by actions taken under TSCA. TSCA Section 9(b)(1) therefore provides EPA authority to coordinate actions with other EPA offices without ever making a risk finding, or following an identification of risk. This includes coordination on tailoring the scope of TSCA risk evaluations to focus on areas of greatest concern rather than exposure pathways addressed by other EPA-administered statutes and regulatory programs, which does not involve a risk determination or public interest finding under TSCA Section 9(b)(2).

In a narrower application of the broad authority provided by the first sentence of TSCA Section 9(b)(1), the remaining provisions of Section 9(b)(1) provide EPA authority to identify risks and refer certain of those risks for action by other EPA offices. Under the second sentence of Section 9(b)(1), "[i]f the Administrator determines that a risk to health or the environment associated with a chemical substance or mixture could be eliminated or reduced to a sufficient extent by actions taken under the authorities contained in such other Federal laws, the Administrator shall use such authorities to protect against such risk unless the Administrator determines, in the Administrator's discretion, that it is in the public interest to protect against such risk by actions taken under [TSCA]." Coordination of intra-agency action on risks under TSCA Section 9(b)(1) therefore entails both an identification of risk, and a referral of any

risk that could be eliminated or reduced to a sufficient extent under other EPA-administered laws to the EPA office(s) responsible for implementing those laws (absent a finding that it is in the public interest to protect against the risk by actions taken under TSCA).

Risk may be identified by OPPT or another EPA office, and the form of the identification may vary. For instance, OPPT may find that one or more conditions of use for a chemical substance present(s) a risk to human or ecological receptors through specific exposure routes and/or pathways. This could involve a quantitative or qualitative assessment of risk based on reasonably available information (which might include, *e.g.*, findings or statements by other EPA offices or other federal agencies). Alternatively, risk could be identified by another EPA office. For example, another EPA office administering non-TSCA authorities may have sufficient monitoring or modeling data to indicate that a particular condition of use presents risk to certain human or ecological receptors, based on expected hazards and exposures. This risk finding could be informed by information made available to the relevant office under TSCA Section 9(e), which supports cooperative actions through coordinated information-sharing.

Following an identification of risk, EPA would determine if that risk could be eliminated or reduced to a sufficient extent by actions taken under authorities in other EPA-administered laws. If so, TSCA requires EPA to "use such authorities to protect against such risk," unless EPA determines that it is in the public interest to protect against that risk by actions taken under TSCA. In some instances, EPA may find that a risk could be sufficiently reduced or eliminated by future action taken under non-TSCA authority. This might include, *e.g.*, action taken under the authority of the Safe Drinking Water Act (SDWA) to address risk to the general population from a chemical substance in drinking water, particularly if the Office of Water has taken preliminary steps such as listing the subject chemical substance on the Contaminant Candidate List (CCL). This sort of risk finding and referral could occur during the risk evaluation process, thereby enabling EPA to use more a relevant and appropriate authority administered by another EPA office to protect against hazards or exposures to affected receptors.

Legislative history on TSCA Section 9(b)(1) supports both broad coordination on current intra-agency actions, and narrower coordination when risk is identified and referred to another EPA office for action. A Conference Report from the time of TSCA's passage explained that Section 9 is intended "to assure that overlapping or duplicative regulation is avoided while attempting to provide for the greatest possible measure of protection to health and the environment." S. Rep. No. 94-1302 at 84. See also H. Rep. No. 114-176 at 28 (stating that the 2016 TSCA amendments "reinforce TSCA's original purpose of filling gaps in Federal law," and citing new language in Section 9(b)(2) intended "to focus the Administrator's exercise of discretion regarding which statute to apply and to encourage decisions that avoid confusion, complication, and duplication"). Exercising TSCA Section 9(b)(1) authority to coordinate on tailoring TSCA risk evaluations is consistent with this expression of Congressional intent.

Legislative history also supports a reading of Section 9(b)(1) under which EPA coordinates intra-agency action, including information-sharing under TSCA Section 9(e), and the appropriately positioned EPA office is responsible for the identification of risk and actions to protect against such risks. See, *e.g.*, Senate Report 114-67, 2016 Cong. Rec. S3522 (under TSCA Section 9, "if the Administrator finds that disposal of a chemical substance may pose risks that could be prevented or reduced under the Solid Waste Disposal Act, the Administrator should ensure that the relevant office of the EPA receives that information"); H. Rep. No. 114-176 at 28, 2016 Cong. Rec. S3522 (under Section 9, "if the Administrator should ensure that a risk to health or the environment associated with disposal of a chemical substance or reduced to a sufficient extent under the Solid Waste Disposal Act, the

Administrator should use those authorities to protect against the risk"). Legislative history on Section 9(b)(1) therefore supports coordination with and referral of action to other EPA offices, especially when statutes and associated regulatory programs administered by those offices could address exposure pathways or risks associated with conditions of use, hazards, and/or exposure pathways that may otherwise be within the scope of TSCA risk evaluations.

#### TSCA Sections 2(c) and 18(d)

Finally, TSCA Section 2(c) supports coordinated action on exposure pathways and risks addressed by other EPA-administered statutes and regulatory programs. Section 2(c) directs EPA to carry out TSCA in a "reasonable and prudent manner" and to consider "the environmental, economic, and social impact" of its actions under TSCA. Legislative history from around the time of TSCA's passage indicates that Congress intended EPA to consider the context and take into account the impacts of each action under TSCA. S. Rep. No. 94-698 at 14 ("the intent of Congress as stated in this subsection should guide each action the Administrator takes under other sections of the bill").

Section 18(d)(1) specifies that state actions adopted or authorized under any Federal law are not preempted by an order of no unreasonable risk issued pursuant to TSCA Section 6(i)(1) or a rule to address unreasonable risk issued under TSCA Section 6(a). Thus, even if a risk evaluation were to address exposures or risks that are otherwise addressed by other federal laws and, for example, implemented by states, the state laws implementing those federal requirements would not be preempted. In such a case, both the other federal and state laws, as well as any TSCA Section 6(i)(1) order or TSCA Section 6(a) rule, would apply to the same issue area. See also TSCA Section 18(d)(1)(A)(iii). In legislative history on amended TSCA pertaining to Section 18(d), Congress opined that "[t]his approach is appropriate for the considerable body of law regulating chemical releases to the environment, such as air and water quality, where the states have traditionally had a significant regulatory role and often have a uniquely local concern." Sen. Rep. 114-67 at 26.

EPA's careful consideration of whether other EPA-administered authorities are available, and more appropriate, for addressing certain exposures and risks is consistent with Congress' intent to maintain existing federal requirements and the state actions adopted to locally and more specifically implement those federal requirements, and to carry out TSCA in a reasonable and prudent manner. EPA believes it is both reasonable and prudent to tailor TSCA risk evaluations rather than attempt to evaluate and regulate potential exposures and risks from those media under TSCA. This approach furthers Congressional direction and EPA aims to efficiently use Agency resources, avoid duplicating efforts taken pursuant to other Agency programs, and meet the statutory deadline for completing risk evaluations.

EPA-administered statutes and regulatory programs that address specific exposure pathways and/or risks are listed as follows:

#### Ambient Air Pathway

The Clean Air Act (CAA) contains a list of hazardous air pollutants (HAP) and provides EPA with the authority to add to that list pollutants that present, or may present, a threat of adverse human health effects or adverse environmental effects. For stationary source categories emitting HAP, the CAA requires issuance of technology-based standards and, if necessary, additions or revisions to address developments in practices, processes, and control technologies, to ensure the standards adequately protect public health and the environment. The CAA thereby provides EPA with comprehensive

authority to regulate emissions to ambient air of any hazardous air pollutant. 1,2-Dichloropropane is a HAP. See 42 U.S.C. 7412. EPA has issued a number of technology-based standards for source categories that emit 1,2-dichloropropane to ambient air and, as appropriate, has reviewed, or is in the process of reviewing remaining risks. See 40 CFR part 63.

Emission pathways to ambient air from commercial and industrial stationary sources and associated inhalation exposure of the general population or terrestrial species in this TSCA evaluation from stationary source releases of 1,2-dichloropropane to ambient air are covered under the jurisdiction of the CAA. EPA's Office of Air and Radiation and Office of Pollution Prevention and Toxics will continue to work together to exchange information related to toxicity and occurrence data on chemicals undergoing risk evaluation under TSCA. As such, EPA does not plan to evaluate exposures to the general population from ambient air in the risk evaluation under TSCA. This regulatory coverage is represented by the red shading in Figure 2-15.

#### Drinking Water Pathway

EPA has regular analytical processes to identify and evaluate unregulated drinking water contaminants of potential regulatory concern for public water systems under the SDWA. In addition, the SDWA requires EPA to review and revise "as appropriate" existing drinking water regulations every 6 years.

EPA has promulgated National Primary Drinking Water Regulations (NPDWRs) under the SDWA for 1,2-dichloropropane. EPA has set an enforceable Maximum Contaminant Level (MCL) as close as feasible to a health based, non-enforceable Maximum Contaminant Level Goal (MCLG). Public water systems are required to monitor for the regulated chemical based on a standardized monitoring schedule to ensure compliance with the MCL. The MCL for 1,2-dichloropropane in water is 0.005 mg/L and the MCLG is 0 mg/L.

The drinking water exposure pathway for 1,2-dichloropropane is currently addressed in the NPDWR. As such, EPA does not plan to evaluate exposures to the general population from drinking water exposure in the risk evaluation. This regulatory coverage is represented by the dark blue shading in Figure 2-15. EPA's Office of Water and Office of Pollution Prevention and Toxics will continue to work together providing understanding and analysis of the SDWA regulatory analytical processes and to exchange information related to toxicity and occurrence data on chemicals undergoing risk evaluation under TSCA.

#### Ambient Water Pathway

EPA has developed Clean Water Act (CWA) Section 304(a) recommended human health criteria for 122 chemicals and aquatic life criteria for 47 chemicals. A subset of these chemicals is identified as "priority pollutants" (103 human health and 27 aquatic life), including 1,2-dichloropropane. The CWA requires that states adopt numeric criteria for priority pollutants for which EPA has published recommended criteria under Section 304(a), the discharge or presence of which in the affected waters could reasonably be expected to interfere with designated uses adopted the state. For pollutants with recommended human health criteria, EPA regulations require that state criteria contain sufficient parameters and constituents to protect designated uses. Once states adopt criteria as water quality standards, the CWA requires that National Pollutant Discharge Elimination System (NPDES) discharge permits include effluent limits as stringent as necessary to meet standards CWA Section 301(b)(1)(C). This permit issuance process accounts for risk in accordance with the applicable ambient water exposure pathway (human health or aquatic life as applicable) for the designated water use. The human health criteria for the consumption of water and organism is 0.90 ug/L and for the consumption of organism only is 31 ug/L.

EPA develops recommended water quality criteria under Section 304(a) of the Clean Water Act (CWA) for pollutants in surface water that are protective of aquatic life or human health designated uses. EPA has developed recommended water quality criteria for protection of human health for 1,2-dichloropropane which are available for possible adoption into state water quality standards and are available for possible use by NPDES permitting authorities in deriving effluent limits to meet state narrative criteria. See, *e.g.*, 40 CFR part 423, Appendix D; 40 CFR 131.11(b)(1); 40 CFR 122.44(d)(1)(vi). As such, EPA does not plan to evaluate exposures to the general population from surface water in the risk evaluation under TSCA. This regulatory coverage is represented by the light blue shading in Figure 2-15. EPA's OW and OPPT will continue to work together to exchange information related to toxicity of chemicals undergoing risk evaluation under TSCA. EPA may update its CWA Section 304(a) water quality criteria for 1,2-dichloropropane in the future under the CWA.

EPA has not developed CWA Section 304(a) recommended water quality criteria for the protection of aquatic life for 1,2-dichloropropane, so there are no national recommended criteria for this use available for adoption into state water quality standards and available for use in NPDES permits. As a result, EPA is evaluating exposures to aquatic species from surface water in the risk evaluation under TSCA. The Office of Water may issue CWA Section 304(a) aquatic life criteria for 1,2-dichloropropane in the future.

#### Onsite Releases to Land Pathway

The Comprehensive Environmental Response, Compensation, and Liability Act, otherwise known as CERCLA, provides broad authority under the statute (generally referred to as Superfund) to clean up uncontrolled or abandoned hazardous-waste sites as well as accidents, spills, and other releases of hazardous substances, pollutants and contaminants into the environment. Through CERCLA, EPA was given authority to seek out those parties potentially responsible for the release of hazardous substances and either have them clean up the release or compensate the Federal government for undertaking the response action.

CERCLA Section 101(14) defines "hazardous substance" by referencing other environmental statutes, including toxic pollutants listed under CWA Section 307(a); hazardous substances designated pursuant to CWA Section 311(b)(2)(A); hazardous air pollutants listed under CAA Section 112; imminently hazardous substances with respect to which EPA has taken action pursuant to TSCA Section 7; and hazardous wastes having characteristics identified under or listed pursuant to RCRA Section 3001. See 40 CFR 302.4. CERCLA Section 102(a) also authorizes EPA to promulgate regulations designating as hazardous substances those substances which, when released into the environment, may present substantial danger to the public health or welfare or the environment. EPA must also promulgate regulations establishing the quantity of any hazardous substance the release of which must be reported under Section 103. Section 103 requires persons in charge of vessels or facilities to report to the National Response Center if they have knowledge of a release of a hazardous substance above the reportable quantity threshold.

1,2-Dichloropropane is a hazardous substance under CERCLA. Releases of 1,2-dichloropropane in excess of 1000 pounds within a 24-hour period must be reported (40 CFR 302.4. 302.6). The scope of this EPA TSCA risk evaluation does not include on-site releases to the environment of 1,2-dichloropropane at Superfund sites and subsequent exposure of the general population or non-human species.

#### **Disposal and Soil Pathways**

1,2-Dichloropropane is included on the list of hazardous wastes pursuant to the Resource Conservation and Recovery Act (RCRA) Section 3001 (40 CFR § 261.33) as a listed waste on the U list (U083). The general standard in RCRA Section 3004(a) for the technical criteria that govern the management (treatment, storage, and disposal) of hazardous waste are those "*necessary to protect human health and the environment*," RCRA Section 3004(a). The regulatory criteria for identifying "characteristic" hazardous wastes and for "listing" a waste as hazardous also relate solely to the potential risks to human health or the environment (40 CFR §§ 261.11, 261.21-261.24). RCRA statutory criteria for identifying hazardous wastes require EPA to "tak[e] into account toxicity, persistence, and degradability in nature, potential for accumulation in tissue, and other related factors such as flammability, corrosiveness, and other hazardous wastes that are incinerated (subject to joint control under RCRA Subtitle C and the CAA hazardous waste combustion Maximum Achievable Control Technology (MACT)) or injected into Underground Injection Control (UIC) Class I hazardous waste wells (subject to joint control under Subtitle C and the SDWA)<sup>8</sup>.

Emissions to ambient air from municipal and industrial waste incineration and energy recovery units that form combustion by-products from incineration treatment of 1,2 dichloropropane wastes may be subject to regulations, as would 1,2-dichloropropane burned for energy recovery.

EPA does not plan to evaluate on-site releases to land that go to underground injection or associated exposures to the general population or terrestrial species in its risk evaluation. TRI reporting in 2018 indicated 1,2-dichloropropane was not released to underground injection to Class I hazardous waste wells. However, if environmental disposal of 1,2-dichloropropane were injected into Class I hazardous waste well, these well types fall under the jurisdiction of RCRA and SDWA and disposal of 1,2 - dichloropropane via underground injection would not likely to result in environmental and general population exposures. See 40 CFR part 144.

Based on 2018 TRI reporting, there were no releases reported for 1,2-dichloropropane to RCRA Subtitle C hazardous waste landfills. Design standards for Subtitle C landfills require double liner, double leachate collection and removal systems, leak detection system, run on, runoff, and wind dispersal controls, and a construction quality assurance program. They are also subject to closure and post-closure care requirements including installing and maintaining a final cover, continuing operation of the leachate collection and removal system until leachate is no longer detected, maintaining and monitoring the leak detection and groundwater monitoring system. Bulk liquids may not be disposed in Subtitle C landfills. Subtitle C landfill operators are required to implement an analysis and testing program to ensure adequate knowledge of waste being managed, and to train personnel on routine and emergency operations at the facility. Hazardous waste being disposed in Subtitle C landfills must also meet RCRA waste treatment standards before disposal. See 40 CFR part 264. Based on 2018 reporting, the majority of TRI land disposal is transferred to "other landfills" both on-site and off-site (861 pounds reported in 2018). As a result, EPA does not plan to evaluate on-site releases to land from RCRA Subtitle C hazardous waste landfills or exposures of the general population or terrestrial species from such releases in the TSCA evaluation. This regulatory coverage is represented by the pink shading in Figure 2-15.

<sup>&</sup>lt;sup>8</sup> This is not an exclusive list of Subtitle C authority, as it also covers, for example, disposal to surface impoundments, waste piles, and land treatment.


1,2-Dichloropropane is present in commercial and consumer products that may be disposed of in Municipal Solid Waste (MSW) landfills. Based on 2018 reporting, the majority of TRI land disposal is transferred to "all other land disposal" both on-site and off-site. On-site RCRA Subtitle D municipal solid waste landfill releases leading to exposures of the general population (including susceptible populations) or terrestrial species may be minimal based on current TRI releases (*i.e.*, 861 lb in 2018) for 1.2-dichloropropane. While permitted and managed by the individual states, municipal solid waste (MSW) landfills are required by federal regulations to implement some of the same requirements as Subtitle C landfills. MSW landfills generally must have a liner system with leachate collection and conduct groundwater monitoring and corrective action when releases are detected. MSW landfills are also subject to closure and post-closure care requirements and must have financial assurance for funding of any needed corrective actions. MSW landfills have also been designed to allow for the small amounts of hazardous waste generated by households and very small quantity waste generators (less than 220 lb per month). Bulk liquids, such as free solvent, may not be disposed of at MSW landfills. See 40 CFR part 258. As a result, EPA does not plan to evaluate on-site releases to land from RCRA Subtitle D municipal solid waste (MSW) landfills or exposures of the general population or terrestrial species from such releases in the TSCA evaluation. This regulatory coverage is represented by the pink shading in Figure 2-15.

On-site releases to land may occur from industrial non-hazardous and construction/demolition waste landfills. Industrial non-hazardous and construction/demolition waste landfills are primarily regulated under authorized state regulatory programs. States must also implement limited federal regulatory requirements for siting, groundwater monitoring, and corrective action, and a prohibition on open dumping and disposal of bulk liquids. States may also establish additional requirements such as for liners, post-closure and financial assurance, but are not required to do so. See *e.g.*, RCRA Section 3004(c), 4007; 40 CFR part 257. As a result, EPA does not plan to evaluate on-site releases to land from industrial non-hazardous waste and construction/demolition waste landfills or associated exposures to the general population. This regulatory coverage is represented by the pink shading in Figure 2-15.

### 2.6.3.2 Conceptual Model for Environmental Releases and Wastes: Potential Exposures and Hazards

As described in Section 2.6.3.1, some pathways in the conceptual models are covered under the jurisdiction of other environmental statutes administered by EPA. The conceptual model depicted in Figure 2-16 presents the exposure pathways, exposure routes and hazards to environmental receptors from releases and wastes from industrial, commercial, and consumer uses of 1,2-dichloropropane that EPA plans to evaluate.

The diagram shown in Figure 2-16 includes releases from industrial, commercial and/or consumer uses to water/sediment; biosolids and soil, via direct and indirect discharges to water, that may lead to exposure to aquatic receptors. The supporting basis for environmental pathways considered for 1,2-dichloropropane are included in Appendix H.



### Figure 2-16. 1,2-Dichloropropane Conceptual Model for Environmental Releases and Wastes: Environmental Exposures and Hazards

The conceptual model presents the exposure pathways, exposure routes and hazards to and environmental receptors from releases and wastes from industrial, commercial, and consumer uses of 1,2-dichloropropane that EPA plans to consider in the risk evaluation.

a) Industrial wastewater or liquid wastes may be treated on-site and then released to surface water (direct discharge), or pre-treated and released to POTW (indirect discharge). For consumer uses, such wastes may be released directly to POTW.

### 2.7 Analysis Plan

The analysis plan is based on EPA's knowledge of 1,2-dichloropropane resulting from the full-text screening of reasonably available information as described in Section 2.1. EPA encourages submission of additional existing data, such as full study reports or workplace monitoring from industry sources, that may be relevant to EPA's evaluation of conditions of use, exposures, hazards and PESS during risk evaluation. As discussed in the *Application of Systematic Review in TSCA Risk Evaluations* document (U.S. EPA, 2018), targeted supplemental searches during the analysis phase may be necessary to identify additional information (*e.g.*, commercial mixtures) for the risk evaluation of 1,2-dichloropropane. For any additional data needs identified during the risk evaluation, EPA may use the Agency's TSCA authorities under Sections 4, 8 or 11, as appropriate.

#### 2.7.1 Physical and Chemical Properties and Environmental Fate

EPA plans to analyze the physical and chemical properties and environmental fate and transport of 1,2dichloropropane as follows:

- Review reasonably available measured or estimated physical and chemical properties and environmental fate endpoint data collected using systematic review procedures and, where reasonably available, environmental assessments conducted by other regulatory agencies. EPA plans to evaluate data and information collected through the systematic review methods and public comments about the physical and chemical properties (Appendix A) and fate endpoints (Appendix C), some of which appeared in the *Proposed Designation of 1,2-Dichloropropane* (*CASRN 78-87-5*) as a High-Priority Substance for Risk Evaluation (U.S. EPA, 2019e). All sources cited in EPA's analysis will be evaluated according to the procedures and metrics described in the *Application of Systematic Review in TSCA Risk Evaluations* (U.S. EPA, 2018). Where the systematic review process does not identify experimentally measured chemical property values of sufficiently high quality, testing will be requested under the TSCA Section 4 authority, or values will be estimated using chemical parameter estimation models as appropriate. Model-estimated fate properties will be reviewed for applicability and quality.
- 2) Using measured data and/or modeling, determine the influence of physical and chemical properties and environmental fate endpoints (*e.g.*, persistence, bioaccumulation, partitioning, transport) on exposure pathways and routes of exposure to human and environmental receptors.

Measured data and, where necessary, model predictions of physical and chemical properties and environmental fate endpoints will be used to characterize the persistence and movement of 1,2dichloropropane within and across environmental media. The fate endpoints of interest include volatilization, sorption to organic matter in sediments, water solubility, aqueous and atmospheric photolysis rates, aerobic and anaerobic biodegradation rates, and potential bioconcentration and bioaccumulation. These endpoints will be used in exposure calculations. **3)** Conduct a weight of the scientific evidence evaluation of physical and chemical properties and environmental fate data, including qualitative and quantitative sources of information. During risk evaluation, EPA plans to evaluate and integrate the physical and chemical properties and environmental fate evidence identified in the literature inventory using the methods described in the *Application of Systematic Review in TSCA Risk Evaluations* (U.S. EPA, 2018).

#### 2.7.2 Exposure

EPA plans to analyze exposure levels in indoor air, surface water, sediment and aquatic biota associated with exposure to 1,2-dichloropropane. Based on its physical and chemical properties, expected sources, and transport and transformation within the outdoor and indoor environment, 1,2-dichloropropane is more likely to be present in some of these media and less likely to be present in others. EPA has not yet determined the exposure levels in these media. Exposure level(s) can be characterized through a combination of reasonably available monitoring data and estimated exposure levels from modeling approaches. Exposure scenarios are combinations of sources (uses), exposure pathways, and exposed receptors. Draft exposure scenarios corresponding to various conditions of use for 1,2-dichloropropane are presented in Appendix F, Appendix G, and Appendix H. EPA plans to analyze scenario-specific exposures.

#### 2.7.2.1 Environmental Releases

EPA plans to analyze releases to environmental media as follows:

1) Review reasonably available published literature and other reasonably available information on processes and activities associated with the conditions of use to analyze the types of releases and wastes generated.

EPA has reviewed some sources containing information on processes and activities resulting in releases, and the information found is described in Appendix E. EPA plans to review additional data sources identified. Potential sources of environmental release data are summarized in Table 2-6 below:

#### Table 2-6. Potential Categories and Sources of Environmental Release Data

U.S. EPA TRI Data	
U.S. EPA Generic Scenarios	
OECD Emission Scenario Documents	
Discharge Monitoring Report (DMR) surface water discharge data for 1,2-dichloropropane	
from NPDES-permitted facilities	

2) Review reasonably available chemical-specific release data, including measured or estimated release data (e.g., data from risk assessments by other environmental agencies). EPA has reviewed key release data sources including the TRI, and the data from this source is summarized in Section 2.3.3. EPA plans to continue to consider additional reasonably available information and will evaluate it during development of the risk evaluation. EPA plans to match identified data to applicable conditions of use and identify data gaps where no data are found for particular conditions of use. EPA plans to address data gaps identified as described in steps 3 and 4 below by considering potential surrogate data and models.

Additionally, for conditions of use where no measured data on releases are reasonably available, EPA may use a variety of methods including release estimation approaches and assumptions in

the Chemical Screening Tool for Exposures and Environmental Releases (ChemSTEER) (<u>U.S.</u> <u>EPA, 2015a</u>).

**3**) Review reasonably available measured or estimated release data for surrogate chemicals that have similar uses and physical properties.

EPA plans to review literature sources identified and if surrogate data are found, these data will be matched to applicable conditions of use for potentially filling data gaps.

4) Review reasonably available data that may be used in developing, adapting or applying exposure models to the particular risk evaluation.

This item will be performed after completion of #2 and #3 above. EPA plans to evaluate relevant data to determine whether the data can be used to develop, adapt or apply models for specific conditions of use (and corresponding release scenarios). EPA has identified information from various EPA statutes and sources (including, for example, regulatory limits, reporting thresholds or disposal requirements) that may be relevant to consider for release estimation and environmental exposures. EPA plans to consider relevant regulatory requirements in estimating releases during risk evaluation.

5) Review and determine applicability of OECD Emission Scenario Documents (ESDs) and EPA Generic Scenarios to estimation of environmental releases.

EPA has identified potentially relevant OECD Emission Scenario Documents (ESDs) and EPA Generic Scenarios (GS) that correspond to some conditions of use; for example, the <u>2011 ESD</u> <u>on the Chemical Industry</u> (<u>OECD</u>, <u>2011</u>) may be useful. EPA plans to critically review these generic scenarios and ESDs to determine their applicability to the conditions of use assessed.

EPA Generic Scenarios are available at the following: <u>https://www.epa.gov/tsca-screening-tools/chemsteer-chemical-screening-tool-exposures-and-environmental-releases.</u>

OECD Emission Scenario Documents are available at the following: <u>http://www.oecd.org/chemicalsafety/risk-assessment/emissionscenariodocuments.htm</u>

If ESDs and GSs are not available, other methods may be considered. EPA may also perform supplemental targeted searches of peer-reviewed or gray literature for applicable models and associated parameters that EPA may use to estimate releases for certain conditions of use. Additionally, for conditions of use where no measured data on releases are available, EPA may use a variety of methods including the application of default assumptions such as standard loss fractions associated with drum cleaning (3%) or single process vessel cleanout (1%).

#### 6) Map or group each condition of use to a release assessment scenario(s).

EPA has identified release scenarios and mapped (*i.e.*, grouped) them to relevant conditions of use as shown in Appendix F. EPA plans to refine the mapping/grouping of release scenarios based on factors (*e.g.*, process equipment and handling, magnitude of production volume used, and exposure/release sources) corresponding to conditions of use using reasonably available information identified. EPA may perform supplemental targeted searches of peer-reviewed or gray literature to better understand certain conditions of use to further develop release scenarios.

#### 7) Evaluate the weight of the scientific evidence of environmental release data.

During risk evaluation, EPA plans to evaluate and integrate the exposure evidence identified in the literature inventory using the methods described in the *Application of Systematic Review in TSCA Risk Evaluation* (U.S. EPA, 2018). EPA plans to integrate the data using systematic review methods to assemble the relevant data, evaluate the data for quality and relevance, including strengths and limitations, followed by synthesis and integration of the evidence.

#### 2.7.2.2 Environmental Exposures

EPA plans to analyze the following in developing its environmental exposure assessment of 1,2-dichloropropane:

1) Review available environmental and biological monitoring data for all media relevant to environmental exposure.

For 1,2-dichloropropane, environmental media which will be analyzed are sediment, and surface water.

- 2) Review reasonably available information on releases to determine how modeled estimates of concentrations near industrial point sources compare with available monitoring data. EPA plans to analyze and consider reasonably available environmental exposure models that meet the scientific standards under TSCA Section 26(h) and that estimate surface water, sediment concentrations will be analyzed and considered alongside available surface water, sediment monitoring data to characterize environmental exposures. Modeling approaches to estimate surface water concentrations, sediment concentrations may generally include the following inputs: direct release into surface water, or sediment, and indirect release into surface water, sediment, fate and transport (partitioning within media) and characteristics of the environment (*e.g.*, river flow, volume of lake, meteorological data).
- 3) Determine applicability of existing additional contextualizing information for any monitored data or modeled estimates during risk evaluation.

Review and characterize monitoring data or modeled estimates to determine how use patterns have changed over recent years and will determine how representative they are of applicable use patterns.

Any studies which relate levels of 1,2-dichloropropane in the environment or biota with specific sources or groups of sources will be evaluated.

4) Group each condition(s) of use to environmental assessment scenario(s).

Refine and finalize exposure scenarios for environmental receptors by considering combinations of sources (use descriptors), exposure pathways including routes, and populations exposed. For 1,2-dichloropropane, the following are noteworthy considerations in constructing exposure scenarios for environmental receptors:

- Estimates of surface water concentrations, sediment concentrations near industrial point sources based on reasonably available monitoring data.
- Consider the following modeling inputs: release into the media of interest, fate and transport and characteristics of the environment.
- Reasonably available biomonitoring data. Monitoring data could be used to compare with species or taxa-specific toxicological benchmarks.

- Applicability of existing additional contextualizing information for any monitored data or modeled estimates during risk evaluation. Review and characterize the spatial and temporal variability, to the extent that data are reasonably available, and characterize exposed aquatic populations.
- Weight of the scientific evidence of environmental occurrence data and modeled estimates
- 5) Evaluate the weight of the scientific evidence of environmental occurrence data and modeled estimates.

During risk evaluation, EPA plans to evaluate and integrate the exposure evidence identified in the literature inventory using the methods described in the *Application of Systematic Review in TSCA Risk Evaluations* (U.S. EPA, 2018).

#### 2.7.2.3 Occupational Exposures

EPA plans to analyze both worker and occupational non-user exposures as follows:

1) Review reasonably available exposure monitoring data for specific condition(s) of use. EPA plans to review exposure data including workplace monitoring data collected by government agencies such as the OSHA and the NIOSH, and monitoring data found in published literature. These workplace monitoring data include personal exposure monitoring data (direct exposures) and area monitoring data (indirect exposures).

EPA plans to consider the influence of applicable regulatory limits and recommended exposure guidelines on occupational exposures in the occupational exposure assessment. The following are some data sources identified thus far:

#### Table 2-7. Potential Sources of Occupational Exposure Data

2019 ATSDR Toxicological Profile for 1,2-dichloropropane		
U.S. OSHA Chemical Exposure Health Data (CEHD) program data		

2) Review reasonably available exposure data for surrogate chemicals that have uses, volatility and physical and chemical properties similar to 1,2-dichloropropane.

EPA plans to review literature sources identified and if surrogate data are found, these data will be matched to applicable conditions of use for potentially filling data gaps. For several conditions of use, EPA believes data for other chlorinated solvents may serve as surrogate for 1,2-dichloropropane.

3) For conditions of use where data are limited or not reasonably available, review existing exposure models that may be applicable in estimating exposure levels.

EPA has identified potentially relevant OECD emission scenario documents (ESDs) and EPA generic scenarios (GSs) corresponding to some conditions of use. For example, the <u>April 2015</u> <u>ESD on Industrial Use of Industrial Cleaners</u> (OECD, 2015) may be used to estimate occupational exposures. EPA plans to critically review these generic scenarios and ESDs to determine their applicability to the conditions of use assessed. EPA may conduct industry outreach or perform supplemental targeted searches of peer-reviewed or gray literature to understand those conditions of use, which may inform identification of exposure scenarios. EPA may also need to perform targeted supplemental searches to identify applicable models that EPA may use to estimate exposures for certain conditions of use.

4) Review reasonably available data that may be used in developing, adapting or applying exposure models to a particular risk evaluation scenario.

This will be performed after #2 and #3 are completed, and based on information developed from #2 and #3, EPA plans to evaluate relevant data to determine whether the data can be used to develop, adapt, or apply models for specific conditions of use (and corresponding exposure scenarios). EPA may utilize existing, peer-reviewed exposure models developed by EPA, other government agencies, or reasonably available in the scientific literature, or EPA may elect to develop additional models to assess specific condition(s) of use. Inhalation exposure models may be simple box models or two-zone (near-field/far-field) models. In two-zone models, the near-field exposure represents potential inhalation exposures to Workers, and the far-field exposure represents potential inhalation exposures to ONUs.

#### 5) Consider and incorporate applicable EC and/or PPE into exposure scenarios.

EPA plans to review potentially relevant data sources on EC and PPE to determine their applicability and incorporation into exposure scenarios during risk evaluation. OSHA recommends employers utilize the hierarchy of controls to address hazardous exposures in the workplace. The hierarchy of controls strategy outlines, in descending order of priority, the use of elimination, substitution, engineering controls, administrative controls, and lastly personal protective equipment (PPE). EPA plans to assess worker exposure pre- and post-implementation of EC, using reasonably available information on control technologies and control effectiveness. For example, EPA may assess worker exposure in industrial use scenarios before and after implementation of local exhaust ventilation.

#### 6) Map or group each condition of use to occupational exposure assessment scenario(s).

EPA has identified occupational exposure scenarios and mapped them to relevant conditions of use (see Appendix F). As presented in the fourth column in Table\_Apx F-1, EPA has completed an initial mapping of exposure scenarios to conditions of use. EPA plans to refine mapping or grouping of occupational exposure scenarios based on factors (*e.g.*, process equipment and handling, magnitude of production volume used, and exposure/release sources) corresponding to conditions of use as additional information is reviewed during risk evaluation. EPA may perform supplemental targeted searches of peer-reviewed or gray literature to better understand certain conditions of use to further develop exposure scenarios.

# 7) Evaluate the weight of the scientific evidence of occupational exposure data, which may include qualitative and quantitative sources of information.

During risk evaluation, EPA plans to evaluate and integrate the exposure evidence identified in the literature inventory using the methods described in the *Application of Systematic Review in TSCA Risk Evaluations* (U.S. EPA, 2018). EPA plans to rely on the weight of the scientific evidence when evaluating and integrating occupational data. EPA plans to integrate the data using systematic review methods to assemble the relevant data, evaluate the data for quality and relevance, including strengths and limitations, followed by synthesis and integration of the evidence.

#### 2.7.2.4 Consumer Exposures

EPA plans to analyze both consumers using a consumer product and bystanders associated with the consumer using the product as follows:

#### 1) Group each condition of use to consumer exposure assessment scenario(s).

Refine and finalize exposure scenarios for consumers by considering combinations of sources (ongoing consumer uses), exposure pathways including routes, and exposed populations.

For 1,2-dichloropropane, the following are noteworthy considerations in constructing consumer exposure scenarios:

- Conditions of use
- Duration, frequency and magnitude of exposure
- Weight fraction of chemical in products
- Amount of chemical used

# 2) Evaluate the relative potential of indoor exposure pathways based on reasonably available data.

Based on physical and chemical properties of 1,2-dichloropropane and the consumer uses identified, inhalation of vapors and mists is expected to be an important indoor exposure pathway for consumers. EPA plans to review all reasonably available information in developing the consumer exposure scenarios and evaluating the exposure pathways in indoor environments.

# **3**) Review existing indoor exposure models that may be applicable in estimating indoor air exposures.

Indoor exposure models that estimate emissions from use of consumer products are available. These models generally consider p-chem properties (*e.g.*, vapor pressure, molecular weight), product specific properties (*e.g.*, weight fraction of the chemical in the product), use patterns (*e.g.*, duration and frequency of use), user environment (*e.g.*, room of use, ventilation rates), and receptor characteristics (*e.g.*, exposure factors, activity patterns). The OPPT's Consumer Exposure Model (CEM) and other similar models can be used to estimate indoor air exposures from consumer products.

4) Review reasonably available empirical data that may be used in developing, adapting or applying exposure models to a particular risk evaluation scenario. For example, existing models developed for a chemical assessment may be applicable to another chemical assessment if model parameter data are reasonably available.

To the extent other organizations have already modeled a 1,2-dichloropropane consumer exposure scenario that is relevant to the OPPT's assessment, EPA plans to evaluate those modeled estimates. In addition, if other chemicals similar to 1,2-dichloropropane have been modeled for similar uses, those modeled estimates will also be evaluated. The underlying parameters and assumptions of the models will also be evaluated.

5) Review reasonably available consumer product-specific sources to determine how those exposure estimates compare with each other and with indoor monitoring data reporting 1,2-dichloropropane in specific media (*e.g.* indoor air).

The availability of 1,2-dichloropropane concentration for various conditions of use will be evaluated. This data provides the source term for any subsequent indoor modeling.

# 6) Review reasonably available population- or subpopulation-specific exposure factors and activity patterns to determine if potentially exposed or susceptible subpopulations need to be further refined.

For 1,2-dichloropropane, EPA plans to consider age-specific behaviors, activity patterns and exposure factors unique to those subpopulations. For some exposure scenarios related to consumer uses, EPA plans to consider whether exposures for adults may differ those of children due to different activities (*e.g.*, children may mouth certain articles) or exposure factors (*e.g.*, inhalation rates).

# 7) Evaluate the weight of the scientific evidence of consumer exposure estimates based on different approaches.

EPA plans to rely on the weight of the scientific evidence when evaluating and integrating data related to consumer exposure. The weight of the scientific evidence may include qualitative and quantitative sources of information. EPA plans to integrate the data using systematic review methods to assemble the relevant data, evaluate the data for quality and relevance, including strengths and limitations, followed by synthesis and integration of the evidence.

#### 2.7.2.5 General Population

EPA does not plan to analyze general population exposures, based on a review of exposure pathways as described in Section 2.6.3. EPA does not plan to include in the risk evaluation pathways under programs of other environmental statutes, administered by EPA, for which long-standing regulatory and analytical processes already exist.

#### 2.7.3 Hazards (Effects)

#### 2.7.3.1 Environmental Hazards

EPA plans to conduct an environmental hazard assessment of 1,2-dichloropropane as follows:

1) Review reasonably available environmental hazard data, including data from alternative test methods (*e.g.*, computational toxicology and bioinformatics; high-throughput screening methods; data on categories and read-across; *in vitro* studies).

EPA plans to analyze the hazards of 1,2-dichloropropane to aquatic organisms, including plants, invertebrates (*e.g.*, insects, arachnids, mollusks, crustaceans), and vertebrates (*e.g.*, mammals, birds, amphibians, fish, reptiles) across exposure durations and conditions if potential environmental hazards are identified through systematic review results and public comments. Additional types of environmental hazard information will also be considered (*e.g.*, analogue and read-across data) when characterizing the potential hazards of 1,2-dichloropropane to aquatic organisms.

EPA plans to evaluate environmental hazard data using the evaluation strategies laid out in the *Application of Systematic Review in TSCA Risk Evaluation* (U.S. EPA, 2018). The study evaluation results will be documented in the risk evaluation phase and data from acceptable studies will be extracted and integrated in the risk evaluation process.

Mechanistic data may include analyses of alternative test data such as novel *in vitro* test methods and high throughput screening. The association between acute and chronic exposure scenarios to the agent and each health outcome will also be integrated. Study results will be extracted and presented in evidence tables or another appropriate format by organ/system.

#### 2) Derive hazard thresholds for aquatic organisms.

Depending on the robustness of the evaluated data for a particular organism or taxa (*e.g.*, aquatic invertebrates), environmental hazard values (*e.g.*, EC<sub>x</sub>. LC<sub>x</sub>, NOEC, LOEC) may be derived and used to further understand the hazard characteristics of 1,2-dichloropropane to aquatic species. Identified environmental hazard thresholds may be used to derive concentrations of concern (COC), based on endpoints that may affect populations of organisms or taxa analyzed.

#### 3) Evaluate the weight of the scientific evidence of environmental hazard data.

During risk evaluation, EPA plans to evaluate and integrate the environmental hazard evidence identified in the literature inventory using the methods described in the *Application of Systematic Review in TSCA Risk Evaluations* (U.S. EPA, 2018).

- 4) Consider the route(s) of exposure, based on reasonably available monitoring and modeling data and other available approaches to integrate exposure and hazard assessments. EPA plans to consider aquatic (*e.g.*, water and sediment exposures) pathways in the 1,2-dichloropropane conceptual model. These organisms may be exposed to 1,2-dichloropropane via a number of environmental pathways (*e.g.*, surface water, sediment, diet).
- 5) Consider a persistent, bioaccumulative, and toxic (PBT) assessment of 1,2dichloropropane.

EPA plans to consider the persistence, bioaccumulation, and toxic (PBT) potential of 1,2dichloropropane after reviewing relevant physical and chemical properties and exposure pathways. EPA plans to assess the reasonably available studies collected from the systematic review process relating to bioaccumulation and bioconcentration (*e.g.*, BAF, BCF) of 1,2dichloropropane. In addition, EPA plans to integrate traditional environmental hazard endpoint values (*e.g.*, LC<sub>50</sub>, LOEC) and exposure concentrations (*e.g.*, surface water concentrations, tissue concentrations) for 1,2-dichloropropane with the fate parameters (*e.g.*, BAF, BCF, BMF, TMF).

#### 6) Conduct an environmental risk estimation and characterization of 1,2-dichloropropane.

EPA plans to conduct a risk estimation and characterization of 1,2-dichloropropane to identify if there are risks to the aquatic environments from the measured and/or predicted concentrations of 1,2-dichloropropane in environmental media (*e.g.*, water, sediment). Risk quotients (RQs) may be derived by the application of hazard and exposure benchmarks to characterize environmental risk (U.S. EPA, 1998; Barnthouse et al., 1982). Analysis of risk for characterization includes a confidence statement in risk estimation which qualitative judgment describing the certainty of the risk estimate considering the strength the evidence scores for hazard and exposure and the limitations, and relevance.

#### 2.7.3.2 Human Health Hazards

EPA plans to analyze human health hazards as follows:

Review reasonably available human health hazard data, including data from alternative test methods (*e.g.*, computational toxicology and bioinformatics; high-throughput screening methods; data on categories and read-across; *in vitro* studies; systems biology).
 EPA plans to evaluate human health studies using the evaluation strategies laid out in the *Application of Systematic Review in TSCA Risk Evaluation* (U.S. EPA, 2018) and updates to the epidemiological data quality criteria released with the first ten risk evaluations. The study

evaluation results will be documented in the risk evaluation phase and data from acceptable studies will be extracted and integrated in the risk evaluation process.

Mechanistic data may include analyses of alternative test data such as novel *in vitro* test methods and high throughput screening. The association between acute and chronic exposure scenarios to the agent and each health outcome will also be integrated. Study results will be extracted and presented in evidence tables or another appropriate format by organ/system.

2) In evaluating reasonably available data, determine whether particular human receptor groups may have greater susceptibility to the chemical's hazard(s) than the general population.

Reasonably available human health hazard data will be evaluated to ascertain whether some human receptor groups may have greater susceptibility than the general population to 1,2-dichloropropane hazard(s). Susceptibility of particular human receptor groups to 1,2-dichloropropane will be determined by evaluating information on factors that influence susceptibility.

EPA has reviewed some sources containing hazard information associated with susceptible populations and lifestages such as pregnant women and infants. Pregnancy (*i.e.*, gestation) and childhood are potential susceptible lifestages for 1,2-dichloropropane exposure. EPA may quantify these differences in the risk evaluation following further evaluation of the reasonably available data and information.

3) Conduct hazard identification (the qualitative process of identifying non-cancer and cancer endpoints) and dose-response assessment (the quantitative relationship between hazard and exposure) for identified human health hazard endpoints.

Human health hazards from acute and chronic exposures will be identified by evaluating the human and animal data that meet the systematic review data quality criteria described in the systematic review documentation that EPA plans to publish prior to finalizing the scope document. Hazards identified by studies meeting data quality criteria will be grouped by routes of exposure relevant to humans (*e.g.*, oral, dermal, inhalation) and by cancer and noncancer endpoints.

Dose-response assessment will be performed in accordance with EPA guidance (U.S. EPA, 2012a, 2011b, 1994) developing points of departure (POD) for either margins of exposure (MOEs), cancer slope factors (CSFs), oral slope factors (OSFs), and/or inhalation unit risks (IURs). Dose-response analyses may be used if the data meet data quality criteria and if additional information on the identified hazard endpoints are not reasonably available or would not alter the analysis.

The cancer mode of action (MOA) analyses determine the relevancy of animal data to human risk and how data can be quantitatively evaluated. If cancer hazard is determined to be applicable to 1,2-dichloropropane, EPA plans to evaluate information on genotoxicity and the MOA for all cancer endpoints to determine the appropriate approach for quantitative cancer assessment in accordance with the U.S. EPA Guidelines for Carcinogen Risk Assessment (U.S. EPA, 2005a). In accordance with EPA's Supplemental Guidance for Assessing Susceptibility from Early-life Exposures to Carcinogens (U.S. EPA, 2005b), EPA plans to determine whether age-dependent

adjustment factors (ADAFs) are appropriate for 1,2-dichloropropane for specific conditions of use based upon potential exposures to children.

4) Derive points of departure (PODs) where appropriate; conduct benchmark dose modeling depending on the reasonably available data. Adjust the PODs as appropriate to conform (*e.g.*, adjust for duration of exposure) to the specific exposure scenarios evaluated. Hazard data will be evaluated to determine the type of dose-response modeling that is applicable. Where modeling is feasible, a set of dose-response models that are consistent with a variety of potentially underlying biological processes will be applied to empirically model the dose-response relationships in the range of the observed data consistent with EPA's *Benchmark Dose Technical Guidance Document* (U.S. EPA, 2012a). Where dose-response modeling is not feasible, NOAELs or LOAELs will be identified. Non-quantitative data will also be evaluated for contribution to weight of the scientific evidence or for evaluation of qualitative endpoints that are not appropriate for dose-response assessment.

EPA plans to evaluate whether the reasonably available PBPK and empirical kinetic models are adequate for route-to-route and interspecies extrapolation of the POD, or for extrapolation of the POD to standard exposure durations (*e.g.*, lifetime continuous exposure). If application of the PBPK model is not possible, oral PODs may be adjusted by  $BW^{3/4}$  scaling in accordance with U.S. EPA (2011b), and inhalation PODs may be adjusted by exposure duration and chemical properties in accordance with U.S. EPA (1994).

- 5) Evaluate the weight of the scientific evidence of human health hazard data. During risk evaluation, EPA plans to evaluate and integrate the human health hazard evidence identified in the literature inventory under acute and chronic exposure conditions using the methods described in the *Application of Systematic Review in TSCA Risk Evaluations* (U.S. EPA, 2018).
- 6) Consider the route(s) of exposure (*e.g.*, oral, inhalation, dermal), reasonably available route-to-route extrapolation approaches; biomonitoring data; and approaches to correlate internal and external exposures to integrate exposure and hazard assessment. At this stage of review, EPA believes there will be sufficient reasonably available data to conduct a dose-response analysis and/or benchmark dose modeling for the oral route of exposure. EPA plans to also evaluate any potential human health hazards following dermal and inhalation exposure to 1,2-dichloropropane, which could be important for worker, consumer and general population risk analysis. Reasonably available data will be assessed to determine whether or not a point of departure can be identified for the dermal and inhalation routes.

If sufficient reasonably available toxicity studies are not identified through the systematic review process to assess risks from inhalation or dermal exposure, then a route-to-route extrapolation may be needed. The preferred approach is to use a PBPK model (U.S. EPA, 2006a). Without an adequate PBPK model, considerations regarding the adequacy of data for route-to-route extrapolation are described in *Methods for Derivation of Inhalation Reference Concentrations and Application of Inhalation Dosimetry* (U.S. EPA, 1994). EPA may use these considerations when determining whether to extrapolate from the oral to the inhalation route of exposure. Similar approaches for oral-to-dermal route extrapolation are described in EPA guidance

document Risk Assessment Guidance for Superfund Volume I: Human Health Evaluation Manual (Part E, Supplemental Guidance for Dermal Risk Assessment) (U.S. EPA, 2004).

If there are acceptable inhalation data after completion of systematic review, EPA may also consider extrapolating from the inhalation to the dermal route if first-pass metabolism through the liver via the oral route is expected because in that case, use of data from the oral route is not recommended (U.S. EPA, 1994). EPA may also consider inhalation-to-dermal route extrapolation if an inhalation toxicity study with a sensitive hazard endpoint is used to evaluate risks. Based on these considerations, EPA extrapolated from the inhalation to the dermal route for several of the first ten risk evaluations under amended TSCA, including methylene chloride (U.S. EPA, 2020d) and carbon tetrachloride (U.S. EPA, 2020b).

7) Conduct a human health risk estimation and characterization of 1,2-dichloropropane. Analysis of risk for characterization includes a confidence statement in risk estimation. This confidence statement is based on qualitative judgment describing the certainty of the risk estimate considering the strength of the evidence scores for hazard and exposure along with their limitations and relevance. The lowest confidence evaluation for either hazard or exposure will drive the overall confidence estimate.

#### 2.7.4 Summary of Risk Approaches for Characterization

Risk characterization is an integral component of the risk assessment process for both environmental and human health risks. EPA plans to derive the risk characterization in accordance with EPA's *Risk Characterization Handbook* (U.S. EPA, 2000). As defined in EPA's Risk Characterization Policy, "the risk characterization integrates information from the preceding components of the risk evaluation and synthesizes an overall conclusion about risk that is complete, informative and useful for decision makers" (U.S. EPA, 2000). Risk characterization is considered to be a conscious and deliberate process to bring all important considerations about risk, not only the likelihood of the risk but also the strengths and limitations of the assessment, and a description of how others have assessed the risk into an integrated picture.

The level of information contained in each risk characterization varies according to the type of assessment for which the characterization is written. Regardless of the level of complexity or information, the risk characterization for TSCA risk evaluations will be prepared in a manner that is transparent, clear, consistent, and reasonable (U.S. EPA, 2000) and consistent with the requirements of the *Procedures for Chemical Risk Evaluation Under the Amended Toxic Substances Control Act* (82 FR 33726, July 20, 2017). As discussed in 40 CFR 702.43, risk characterization has a number of considerations. This is the step where EPA integrates the hazard and exposure assessments into risk estimates for the identified populations (including any PESS) and ecological characteristics and weighs the scientific evidence for the identified hazards and exposures. The risk characterization does not consider costs or other nonrisk factors, and takes into account, "where relevant, the likely duration, intensity, frequency, and number of exposures under the condition(s) of use ...." The risk characterization also summarizes the following considerations: (1) uncertainty and variability in each step of the risk evaluation; (2) data quality, and any applicable assumptions used; (3) alternative interpretations of data and analyses, where appropriate; and (4) any considerations for environmental risk evaluations, if necessary (*e.g.*, related to nature and magnitude of effects).

EPA plans to also be guided by EPA's Information Quality Guidelines (U.S. EPA, 2002) as it provides guidance for presenting risk information. Consistent with those guidelines, in the risk characterization, EPA plans to also identify: (1) each population addressed by an estimate of applicable risk effects; (2) the expected risk or central estimate of risk for the PESS affected; (3) each appropriate upper-bound or lower-bound estimate of risk; (4) each significant uncertainty identified in the process of the assessment of risk effects and the studies that would assist in resolving the uncertainty; and (5) peer reviewed studies known to the Agency that support, are directly relevant to, or fail to support any estimate of risk effects and the methodology used to reconcile inconsistencies in the scientific information.

### 2.8 Peer Review

Peer review will be conducted in accordance with EPA's regulatory procedures for chemical risk evaluations, including using EPA's Peer Review Handbook (U.S. EPA, 2015b) and other methods consistent with Section 26 of TSCA (see 40 CFR 702.45). As explained in the Risk Evaluation Rule, the purpose of peer review is for the independent review of the science underlying the risk assessment. Peer review will therefore address aspects of the underlying science as outlined in the charge to the peer review panel such as hazard assessment, assessment of dose-response, exposure assessment, and risk characterization. The draft risk evaluation for 1.2-dichloropropane will be peer reviewed.

### REFERENCES

- ATSDR (Agency for Toxic Substances and Disease Registry). (1989). Toxicological profile for 1,2dichloropropane. Atlanta, GA: U.S. Department of Health and Human Services, U.S. Public Health Service. <u>https://www.atsdr.cdc.gov/ToxProfiles/tp134.pdf</u>
- <u>Axson USA.</u> (2015). Safety data sheet: W&S Ultra Premium Sealer. <u>https://www.woodandstoneglue.com/v/SDS/SDS\_W&S\_Ultra\_Premium\_Sealer\_-</u> <u>USA2EN.PDF</u>
- Barnthouse, LW; DeAngelis, DL; Gardner, RH; O'Neill, RV; Suter, GW; Vaughan, DS. (1982). Methodology for environmental risk analysis. (ORNL/TM-8167). Oak Ridge, TN: Oak Ridge National Laboratory.
- <u>CalEPA</u> (California Environmental Protection Agency). (1999). Public health goal for 1,2dichloropropane in drinking water. California: Office of Environmental Health Hazard Assessment, California Environmental Protection Agency, Pesticide and Environmental Toxicology Section. <u>https://oehha.ca.gov/media/downloads/water/chemicals/phg/12dcpf.pdf</u>
- <u>CDC</u> (Centers for Disease Control and Prevention). (2018). Fourth national report on human exposure to environmental chemicals, updated tables, March 2018, volume one. Atlanta, GA: U.S. Department of Health and Human Services.
  - https://www.cdc.gov/exposurereport/pdf/FourthReport\_UpdatedTables\_Volume1\_Mar2018.pdf
- <u>CDPR</u> (California Department of Pesticide Regulation). (2013). DPR databases chemical ingredient application. <u>https://www.cdpr.ca.gov/dprdatabase.htm</u>
- <u>Chem Service.</u> (2018). Safety data sheet for 1,2-dichloropropane. <u>http://cdn.chemservice.com/product/msdsnew/External/English/N-</u>10156%20English%20SDS%20US.pdf
- <u>Cherrie, JW; Semple, S; Christopher, Y; Saleem, A; Hughson, GW; Philips, A.</u> (2006). How important is inadvertent ingestion of hazardous substances at work? Ann Occup Hyg 50: 693-704. <u>http://dx.doi.org/10.1093/annhyg/mel035</u>
- Chiou, CT; Peters, LJ; Freed, VH. (1979). A physical concept of soil-water equilibria for nonionic organic compounds. 206: 831-832. <u>http://dx.doi.org/10.1126/science.206.4420.831</u>
- <u>CITI</u> (Chemicals Inspection Testing Institute Japan). (1992). Biodegradation and bioaccumulation data of existing chemicals based on the CSCL Japan (pp. 2-18). Tokyo, Japan: Chemical Industry Ecology-Toxicology and Information Center.
- <u>Cohen SZ; Creeger SM; Carsel RF; Enfield CG.</u> (1984). Potential for pesticide contamination of groundwater resulting from agricultural uses. In Krueger RF; Seiber JN (Eds.), Treatment and disposal of pesticide wastes: American Chemcial Society Symposium Series (pp. 297-325). Washington D.C.: American Chemical Society.
- Elsevier. (2019). Reaxys: physical-chemical property data for 1,2-dichloropropane. CAS Registry Number: 78-87-5. Available online
- Hartwig, JS, H; Müller, F. (2008). Nematicides [Type of Work]. https://onlinelibrary.wiley.com/doi/abs/10.1002/14356007.a17\_125.pub2
- Howard, BE; Phillips, J; Miller, K; Tandon, A; Mav, D; Shah, MR; Holmgren, S; Pelch, KE; Walker, V; <u>Rooney, AA; Macleod, M; Shah, RR; Thayer, K.</u> (2016). SWIFT-Review: a text-mining workbench for systematic review. Syst Rev 5: 87. http://dx.doi.org/10.1186/s13643-016-0263-z
- Howard, PH. (1990). Handbook of environmental fate and exposure data for organic chemicals: Volume II: Solvents. Syracuse, NY: CRC Press.
- Mackay, D; Shiu, WY; Ma, KC. (1993). Illustrated handbook of physical-chemical properties and environmental fate for organic chemicals. Volume III. Boca Raton, FL: Lewis Publishers.

- NIOSH. (2016). Updated Immediately Dangerous To Life or Health (IDLH) Values (as of 2016). Atlanta, GA. <u>http://www.cdc.gov/niosh/idlh/intridl4.html</u>
- NIOSH. (2020). NIOSH Potential Carcinogen List [Database]. https://www.cdc.gov/niosh/topics/cancer/npotocca.html
- NLM (National Institutes of Health, National Library of Medicine). (2001). Hazardous Substances Data Bank (HSDB): A TOXNET Database: 1,2-Dichloropropane. <u>https://toxnet.nlm.nih.gov/cgi-bin/sis/search2/r?dbs+hsdb:@term+@rn+@rel+78-87-5</u>
- NLM (National Institutes of Health, National Library of Medicine). (2019). PubChem: Hazardous Substance Data Bank: 1,2-Dichloropropane, 78-87-5 [Website]. https://pubchem.ncbi.nlm.nih.gov/compound/6564#source=HSDB
- O'Neil, MJ. (2013). Propylene dichloride. 78-87-5. [1,2-Dichloroethane]. In The Merck Index (15th ed.). Cambridge, UK: The Royal Society of Chemistry.
- <u>OECD</u> (Organisation for Economic Co-operation and Development). (2006). SIDS initial assessment report: 1,2-Dichloropropane (CASRN: 78-87-5) (pp. 219). Paris, France: United Nations Environment Programme, Organisation for Economic Co-operation and Development. <u>https://hpvchemicals.oecd.org/ui/handler.axd?id=E6CE93C6-9DD5-41DF-ADE6-86ECABD0F533</u>
- <u>OECD</u> (Organisation for Economic Co-operation and Development). (2011). Emission scenario document on the chemical industry. (JT03307750). <u>http://www.oecd.org/env/ehs/risk-assessment/48774702.pdf</u>
- <u>OECD</u> (Organisation for Economic Co-operation and Development). (2015). Emission scenario document on industrial use of industrial cleaners (JT03373556). Paris, France. <u>http://www.oecd.org/officialdocuments/publicdisplaydocumentpdf/?cote=ENV/JM/MONO(2015)</u> <u>3&doclanguage=en</u>
- <u>OECD</u> (Organisation for Economic Co-operation and Development). (2018). OECD Monitoring Database [Database]. <u>http://oecd.org</u>
- OSHA (Occupational Safety and Health Administration). (2009). Occupational Safety and Health Administration (OSHA) permissible exposure limits (PELs). https://www.osha.gov/dsg/annotated-pels/tablez-1.html
- <u>RIVM</u> (Dutch National Institute for Public Health and the Environment). (2007). Ecotoxicologically based environmental risk limits for several volatile aliphatic hydrocarbons (pp. 217). (601782002/2007). Bilthoven, Netherlands: National Institute for Public Health and the Environment (RIVM). <u>https://www.rivm.nl/bibliotheek/rapporten/601782002.pdf</u>
- <u>RSC</u> (Royal Society of Chemistry). (2019). ChemSpider: 1,2-Dichloropropane. Available online at <u>http://www.chemspider.com/Chemical-Structure.6316.html?rid=7f5765b6-12e1-46af-9f9e-17daf91b9682</u>
- Rumble, JR. (2018). CRC handbook of chemistry and physics. In JR Rumble (Ed.), (99 ed.). Boca Raton, FL: CRC Press.
- Sen, D; Kim, M. (2009). Excess molar volumes and excess molar enthalpies of the binary mixtures of 1,2-dichloropropane with di- and triethylene glycol mono-alkyl ethers at T=298.15 K. Fluid Phase Equilibria 285: 30-35. <u>http://dx.doi.org/10.1016/j.fluid.2009.06.018</u>
- Steele, WV; Chirico, RD; Knipmeyer, SE; Nguyen, A. (1997). Vapor pressure, heat capacity, and density along the saturation line, measurements for cyclohexanol, 2-cyclohexen-1-one, 1,2-dichloropropane, 1,4-di-tert-butylbenzene, (+/-)-2-ethylhexanoic acid, 2-(methylamino)ethanol, perfluoro-n-heptane, and sulfolane. Journal of Chemical and Engineering Data 42: 1021-1036. http://dx.doi.org/10.1021/je9701036

- <u>SYKE</u> (Finnish Environment Institute). (2018). Data bank of environmental properties of chemicals 1,2dichloropropane. The Finnish Environment Institute. <u>http://wwwp.ymparisto.fi/scripts/Kemrek/Kemrek\_uk.asp?Method=MAKECHEMdetailsform&t</u> <u>xtChemId=544</u>
- <u>Tenax.</u> (2019). Safety data sheet: Cera Fluida. Revision number 1. <u>https://www.tenax4you.com/v/SDS/EN\_CERA%20FLUIDA-DCP\_GHS\_USA.pdf</u>
- <u>Tenax Spa.</u> (2015a). Safety data sheet: Cera Distaccante. Revision no. 1. Tenax Spa. (2015b). Safety data sheet: Cera Tewax. Revision no. 4.
  - https://www.tenax4you.com/v/SDS/US TEWAX GHS USA.pdf
- <u>Tenax Spa.</u> (2015c). Safety data sheet: Lustro Italiano Stone Polish. Revision no. 1. <u>https://www.lustroitaliano.com/v/sds/us-lustro-spray-stone-sealer-ghs-usa.pdf</u>
- <u>Tenax USA.</u> (2015). Cera Distaccante safety data sheet. <u>https://www.tenax4you.com/v/SDS/US\_CERA%20DISTACCANTE\_GHS\_USA.pdf</u>
- Tenax USA. (2020a). Cera Fluida Liquid Wax 1QT. <u>https://www.tenax4you.com/Liquid-Wax-1-QT-New-p/1maa00bg60.htm</u>
- <u>Tenax USA.</u> (2020b). Lustro Italiano Stone Cleaner. <u>https://www.lustroitaliano.com/natural-stone-</u> <u>cleaner-p/lustroclean.htm</u>
- <u>Tenax USA.</u> (2020c). Tenax Wax Paste Clear 1QT (TeWax). <u>https://www.tenax4you.com/Wax-Paste-Clear-1-QT-p/1mca00bg50.htmCera%20Distaccante</u>
- Tomer, A; Kane, J. (2015). The great port mismatch. U.S. goods trade and international transportation. The Global Cities Initiative. A joint project of Brookings and JPMorgon Chase. https://www.brookings.edu/wp-content/uploads/2015/06/brgkssrvygcifreightnetworks.pdf
- U.S. EPA (U.S. Environmental Protection Agency). (1979). Water-related environmental fate of 129 priority pollutants: Volume II: Halogenated aliphatic hydrocarbons, halogenated ethers, monocyclic aromatics, phthalate esters, polycyclic aromatic hydrocarbons, nitrosamines, and miscellaneous compounds [EPA Report]. (EPA-440/4-79-029b). Washington, DC. http://nepis.epa.gov/Exe/ZyPURL.cgi?Dockey=2000K6JL.txt
- U.S. EPA (U.S. Environmental Protection Agency). (1990). EPA Ambient Monitoring Technology Information Center: Air toxics data [Database]. <u>https://www3.epa.gov/ttnamti1/toxdat.html</u>
- U.S. EPA (U.S. Environmental Protection Agency). (1994). Methods for derivation of inhalation reference concentrations and application of inhalation dosimetry [EPA Report]. (EPA/600/8-90/066F). Research Triangle Park, NC: U.S. Environmental Protection Agency, Office of Research and Development, Office of Health and Environmental Assessment, Environmental Criteria and Assessment Office.

https://cfpub.epa.gov/ncea/risk/recordisplay.cfm?deid=71993&CFID=51174829&CFTOKEN=2 5006317

- <u>U.S. EPA</u> (U.S. Environmental Protection Agency). (1996). EPA Unregulated Contaminant Monitoring Rule (UCMR) Database [Database]. <u>https://www.epa.gov/dwucmr</u>
- U.S. EPA (U.S. Environmental Protection Agency). (1998). Guidelines for ecological risk assessment [EPA Report]. (EPA/630/R-95/002F). Washington, DC: U.S. Environmental Protection Agency, Risk Assessment Forum. <u>https://www.epa.gov/risk/guidelines-ecological-risk-assessment</u>
- U.S. EPA (U.S. Environmental Protection Agency). (2000). Science policy council handbook: Risk characterization handbook. (EPA/100/B-00/002). Washington, DC: U.S. Environmental Protection Agency, Science Policy Council. <u>https://www.epa.gov/risk/risk-characterization-handbook</u>
- <u>U.S. EPA</u> (U.S. Environmental Protection Agency). (2002). Guidelines for ensuring and maximizing the quality, objectivity, utility, and integrity of information disseminated by the Environmental

Protection Agency. (EPA/260/R-02/008). Washington, DC: U.S. Environmental Protection Agency, Office of Environmental Information. <u>https://www.epa.gov/sites/production/files/2017-03/documents/epa-info-quality-guidelines.pdf</u>

- U.S. EPA (U.S. Environmental Protection Agency). (2004). Risk Assessment Guidance for Superfund (RAGS), Volume I: Human health evaluation manual, (Part E: Supplemental guidance for dermal risk assessment). (EPA/540/R/99/005). Washington, DC: U.S. Environmental Protection Agency, Risk Assessment Forum. <u>http://www.epa.gov/oswer/riskassessment/ragse/index.htm</u>
- U.S. EPA (U.S. Environmental Protection Agency). (2005a). Guidelines for carcinogen risk assessment [EPA Report]. (EPA/630/P-03/001B). Washington, DC: U.S. Environmental Protection Agency, Risk Assessment Forum. <u>https://www.epa.gov/sites/production/files/2013-</u>09/documents/cancer\_guidelines\_final\_3-25-05.pdf
- <u>U.S. EPA</u> (U.S. Environmental Protection Agency). (2005b). Supplemental guidance for assessing susceptibility from early-life exposure to carcinogens [EPA Report]. (EPA/630/R-03/003F). Washington, DC: U.S. Environmental Protection Agency, Risk Assessment Forum. https://www3.epa.gov/airtoxics/childrens\_supplement\_final.pdf
- <u>U.S. EPA.</u> (U.S. Environmental Protection Agency). (2006a). Approaches for the application of physiologically based pharmacokinetic (PBPK) models and supporting data in risk assessment (Final Report) [EPA Report] (pp. 1-123). (EPA/600/R-05/043F). Washington, DC: U.S. Environmental Protection Agency, Office of Research and Development, National Center for Environmental Assessment. <u>http://cfpub.epa.gov/ncea/cfm/recordisplay.cfm?deid=157668</u>
- U.S. EPA (U.S. Environmental Protection Agency). (2006b). A framework for assessing health risk of environmental exposures to children (pp. 1-145). (EPA/600/R-05/093F). Washington, DC: U.S. Environmental Protection Agency, Office of Research and Development, National Center for Environmental Assessment. <u>http://cfpub.epa.gov/ncea/cfm/recordisplay.cfm?deid=158363</u>
- <u>U.S. EPA</u> (U.S. Environmental Protection Agency). (2011a). Exposure factors handbook: 2011 edition [EPA Report]. (EPA/600/R-090/052F). Washington, DC. http://cfpub.epa.gov/ncea/cfm/recordisplay.cfm?deid=236252
- U.S. EPA (U.S. Environmental Protection Agency). (2011b). Recommended use of body weight 3/4 as the default method in derivation of the oral reference dose (pp. 1-50). (EPA/100/R-11/0001). Washington, DC: U.S. Environmental Protection Agency, Risk Assessment Forum, Office of the Science Advisor. <u>https://www.epa.gov/sites/production/files/2013-09/documents/recommendeduse-of-bw34.pdf</u>
- U.S. EPA (U.S. Environmental Protection Agency). (2012a). Benchmark dose technical guidance. (EPA/100/R-12/001). Washington, DC: U.S. Environmental Protection Agency, Risk Assessment Forum. https://www.epa.gov/risk/benchmark-dose-technical-guidance
- U.S. EPA (U.S. Environmental Protection Agency). (2012b). Estimation Programs Interface Suite<sup>™</sup> for Microsoft® Windows, v 4.11. Washington, DC. Retrieved from <u>https://www.epa.gov/tsca-</u>screening-tools/epi-suitetm-estimation-program-interface
- <u>U.S. EPA</u> (U.S. Environmental Protection Agency). (2015a). ChemSTEER user guide chemical screening tool for exposures and environmental releases. Washington, DC. <u>https://www.epa.gov/sites/production/files/2015-05/documents/user\_guide.pdf</u>
- <u>U.S. EPA</u> (U.S. Environmental Protection Agency). (2015b). Peer review handbook [EPA Report] (4th ed.). (EPA/100/B-15/001). Washington, DC: U.S. Environmental Protection Agency, Science Policy Council. <u>https://www.epa.gov/osa/peer-review-handbook-4th-edition-2015</u>
- U.S. EPA (U.S. Environmental Protection Agency). (2016a). Non-confidential 2016 Chemical Data Reporting (CDR) Database. <u>http://www.epa.gov/cdr/</u>

- U.S. EPA (U.S. Environmental Protection Agency). (2016b). Provisional peer-review toxicity values for 1,2-dichloropropane (CASRN 78-87-5). Cincinnati, OH: U.S. Environmental Protection Agency, National Center for Environmental Assessment, Superfund Health Risk Technical Support Center. <u>https://hhpprtv.ornl.gov/issue\_papers/Dichloropropane12.pdf</u>
- U.S. EPA (U.S. Environmental Protection Agency). (2017). Toxics Release Inventory (TRI), reporting year 2015. <u>https://www.epa.gov/toxics-release-inventory-tri-program/tri-data-and-tools</u>
- <u>U.S. EPA</u> (U.S. Environmental Protection Agency). (2018). Application of systematic review in TSCA risk evaluations. (740-P1-8001). Washington, DC: U.S. Environmental Protection Agency, Office of Chemical Safety and Pollution Prevention.
   <u>https://www.epa.gov/sites/production/files/2018-</u>06/documents/final application of sr in tsca 05-31-18.pdf
- <u>U.S. EPA</u> (U.S. Environmental Protection Agency). (2019a). Chemical and Products Database (CPDat). <u>https://www.epa.gov/chemical-research/chemical-and-products-database-cpdat</u>
- U.S. EPA (U.S. Environmental Protection Agency). (2019b). Chemical data reporting (2012 and 2016 CBI CDR database). U.S. Environmental Protection Agency, Office of Pollution Prevention and Toxics.
- U.S. EPA (U.S. Environmental Protection Agency). (2019c). Chemistry Dashboard Information for 1,2-Dichloropropane. 78-87-5 [Website].

https://comptox.epa.gov/dashboard/dsstoxdb/results?search=DTXSID0020448

- <u>U.S. EPA</u> (U.S. Environmental Protection Agency). (2019d). CPCat: Chemical and product categories. <u>http://actor.epa.gov/cpcat</u>
- U.S. EPA (U.S. Environmental Protection Agency). (2019e). Proposed designation of 1,2dichloropropane (CASRN 78-87-5) as a high-priority substance for risk evaluation. Washington, DC. <u>https://www.epa.gov/sites/production/files/2019-08/documents/12-dichloropropane\_78-87-</u> <u>5\_high-priority\_proposal\_designation\_082219.pdf</u>
- U.S. EPA (U.S. Environmental Protection Agency). (2019f). TRI Explorer (2018 dataset released November 2019). Washington, DC: U.S. Environmental Protection Agency. https://enviro.epa.gov/triexplorer/
- U.S. EPA (U.S. Environmental Protection Agency). (2020a). Chemical data reporting (2012 and 2016 Public CDR database). Washington, DC: U.S. Environmental Protection Agency, Office of Pollution Prevention and Toxics. ChemView: July 2020. <u>https://chemview.epa.gov/chemview</u>
- U.S. EPA. (U.S. Environmental Protection Agency). (2020b). Draft risk evaluation for carbon tetrachloride (methane, tetrachloro-); CASRN: 56-23-5 (pp. 1-301). (EPA-740-R1-8014). Office of Chemical Safety and Pollution Prevention, U.S. Environmental Protection Agency. https://nepis.epa.gov/Exe/ZyPDF.cgi/P100YHUW.PDF?Dockey=P100YHUW.PDF
- U.S. EPA. (U.S. Environmental Protection Agency). (2020c). Draft Scope of the risk evaluation for 1,2-Dichloropropane CASRN 78-87-5 [EPA Report]. (EPA-740-D-20-006). <u>https://www.epa.gov/sites/production/files/2020-04/documents/casrn-78-87-5\_12-</u> <u>dichloropropane\_draft\_scope.pdf</u>
- U.S. EPA. (U.S. Environmental Protection Agency). (2020d). Risk evaluation for methylene chloride (dichloromethane, dcm); CASRN: 75-09-2 (pp. 1-753). (EPA-740-R1-8010). Office of Chemical Safety and Pollution Prevention, U.S. Environmental Protection Agency. <u>https://www.epa.gov/sites/production/files/2020-</u>06/documents/1\_mecl\_risk\_evaluation\_final.pdf
- <u>USGS</u> (U.S. Geological Survey). (1991a). USGS Monitoring Data: National Water Quality Monitoring Council [Database]. <u>http://www.waterqualitydata.us/portal/</u>

<u>USGS</u> (U.S. Geological Survey). (1991b). USGS Monitoring Data: National Water Quality Monitoring Council - Air [Database].

https://www.waterqualitydata.us/portal/#sampleMedia=Air&mimeType=csv

- <u>USGS</u> (U.S. Geological Survey). (1991c). USGS Monitoring Data: National Water Quality Monitoring Council - Groundwater [Database]. <u>https://www.waterqualitydata.us/portal/#siteType=Aggregate%20groundwater%20use&sample</u> Media=Water&mimeType=csv&dataProfile=activityAll
- <u>USGS</u> (U.S. Geological Survey). (1991d). USGS Monitoring Data: National Water Quality Monitoring Council Sediment [Database].

https://www.waterqualitydata.us/portal/#sampleMedia=Sediment&mimeType=csv

<u>USGS</u> (U.S. Geological Survey). (1991e). USGS Monitoring Data: National Water Quality Monitoring Council - Soil [Database].

https://www.waterqualitydata.us/portal/#sampleMedia=Soil&mimeType=csv

- <u>USGS</u> (U.S. Geological Survey). (1991f). USGS Monitoring Data: National Water Quality Monitoring Council - Surface Water [Database]. <u>https://www.waterqualitydata.us/portal/#siteType=Aggregate%20surface-water-use&sampleMedia=Water&mimeType=csy</u>
- <u>USGS</u> (U.S. Geological Survey). (1991g). USGS Monitoring Data: National Water Quality Monitoring Council Tissue [Database].

https://www.waterqualitydata.us/portal/#sampleMedia=Tissue&mimeType=csv

Van Dijk, H. (1980). Dissipation rates in soil of 1,2-dichloropropane and 1,3- and 2,3-dichloropropenes. 11: 625-632.

### APPENDICES

# Appendix A ABBREVIATED METHODS FOR SEARCHING AND SCREENING

#### A.1 Literature Search of Publicly Available Databases

#### A.1.1 Search Term Genesis and Chemical Verification

To develop the chemical terms for the subsequent literature search for 1,2-dichloropropane, several online sources were queried.

- California Department of Pesticide Regulation: https://www.cdpr.ca.gov/docs/chemical/monster2.htm
- USEPA Chemistry Dashboard: <u>https://comptox.epa.gov/dashboard</u>
- University of Hertfordshire PPDB: Pesticide Properties DataBase: <u>https://sitem.herts.ac.uk/aeru/ppdb/en/search.htm</u>
- USEPA Reregistration Eligibility Decision (RED) documents: https://archive.epa.gov/pesticides/reregistration/web/html/status.html
- Office of Pesticide Programs Pesticide Chemical Search: https://ofmpub.epa.gov/apex/pesticides/f?p=CHEMICALSEARCH:1
- Food and Agriculture Organization of the United Nations: <u>http://www.fao.org/home/en/</u>
- PAN Pesticides Database: <u>http://www.pesticideinfo.org/Search\_Chemicals.jsp</u>

Prior to inclusion in the search term string, all forms of chemical names were subjected to verification from several potential sources (*e.g.*, US EPA Chemistry Dashboard, STN International-CAS; see complete list of sources for chemical verification in Table\_Apx A-1). From these sources, all chemical names, synonyms, CAS number(s), trade names, *etc.* were documented and used to generate terms for database searches.

CHEMICAL SOURCE	CONTENTS	DOCUMENT LOCATION
Chemistry Dashboard	CAS Numbers, Synonyms, Structures, Properties,	Online
(https://comptox.epa.gov/dashboard)	Environmental Fate and Transport.	
Dictionary of Chemical Names and Synonyms	Wide assortment of chemical compounds by chemical name and synonym, has CAS index and some structure data	ECOTOX
Farm Chemicals Handbook-1992	Pesticide information, CAS numbers and synonyms, some structure data ***Sometimes CAS number presented for a compound is for the main constituent only	ECOTOX
OPPT SMILES Verification Source	Structure Data	Electronic verification
RTECS (Registry of Toxic Effects of chemical substance, 1983-84 ed., 2 vols)	Chemical names, synonyms and CAS numbers	ECOTOX
Sigma – Aldrich website58784 http://www.sigma-aldrich.com	Organic and inorganic Compounds by chemical name, has CAS index and some structure and Physical Property data	Online
STN International (CAS) 1994	***Most complete source of chemical name, synonym and structure information, no physical properties	Online

Table\_Apx A-1. Sources of Verification for Chemical Names and Structures

CHEMICAL SOURCE	CONTENTS	DOCUMENT LOCATION
The Pesticide Manual 10th edition, 1994	Pesticide Compounds by chemical name, synonym, product code, has CAS index and some structure and Physical Property data	ECOTOX
TSCA (Toxic Substances Control Act Chemical Substance Inventory, 1985 ed., 5 vols)	Chemical names, synonyms and CAS numbers	ECOTOX
World Wide Web (misc. web sources) A copy of the verification page is saved to the Attachments tab of the chemical entry. This includes company MSDS sheets or Chemical Labels.	Chemical names, synonyms and CAS numbers	Online
California Department of Pesticide Regulation (http://www.cdpr.ca.gov/dprdatabase.htm)	Multiple databases containing chemicals, pesticides, companies, products, <i>etc</i> .	Online
PAN Pesticide Database (http://www.pesticideinfo.org/Search Chemic als.jsp)	Pesticides searchable by name or CAS #. Includes CAS #, Name, synonyms, targets, toxicity data, related chemicals and regulatory information.	Online
US EPA Office of Pesticide Programs Pesticide Fate Database – No web access available. An electronic copy of the data file is located at the Contractor site: PFATE 37 Tables.mdb.	Multiple databases containing chemicals, pesticides, companies, products, <i>etc</i> .	Online

#### A.1.2 Publicly Available Database Searches

The databases listed below were searched for literature containing the chemical search terms. Database searching occurred during April and May of 2019 by an information specialist and the results were stored in the Health and Environmental Research Online (HERO) database and assigned a HERO reference identification number.<sup>9</sup> The present literature search focused only on the chemical name (including synonyms and trade names) with no additional limits. Full details of the search strategy for each database are presented in Appendix A.1.2.1.

After initial deduplication in HERO<sup>10</sup>, these studies were imported into <u>SWIFT Review</u> software (<u>Howard et al., 2016</u>) to identify those references most likely to be applicable to each discipline area (*i.e.* consumer, environmental, and general population exposure, occupational exposure and environmental releases, environmental hazards, human health hazards, and fate and physical chemistry).

#### A.1.2.1 Query Strings for the Publicly-Available Database Searches on 1,2-Dichloropropane

Table\_Apx A-2 presents a list of the data sources, the search dates and number of peer-reviewed references resulting from the searches for 1,2-dichloropropane. The sources are found as online databases and the resulting references were gathered and uploaded into the EPA Health and Environmental Research Online (HERO) database for literature screening.

<sup>&</sup>lt;sup>9</sup>EPA's HERO database provides access to the scientific literature behind EPA science assessments. The database includes more than 600,000 scientific references and data from the peer-reviewed literature used by EPA to develop its regulations.

<sup>&</sup>lt;sup>10</sup> Deduplication in HERO involves first determining whether a matching unique ID exists (e.g., PMID, WOSid, or DOI). If one matches one that already exists in HERO, HERO will tag the existing reference instead of adding the reference again. Second, HERO checks if the same journal, volume, issue and page number are already in HERO. Third, HERO matches on the title, year, and first author. Title comparisons ignore punctuation and case.

Literature Search Results for 1,2-Dichloropropane				
Source	Date of Search	Number of References		
Current Contents	05/16/2019	242		
Web of Science	09/10/2019	417		
ProQuest CSA	05/16/2019	411		
Dissertation Abstracts	05/17/2019	5		
Science Direct	05/20/2019	134		
Agricola	05/17/2019	47		
TOXNET	05/20/2019	684		
PubMed	07/02/2019	182		
UNIFY	05/20/2019	144		
Totals:		2266		

# Table\_Apx A-2. Summary of Data Sources, Search Dates and Number of Peer-Reviewed Literature Search Results for 1,2-Dichloropropane

#### General:

General search terms were compiled and used in the search strategies for each of the databases/sources listed below. Based upon the online search manuals for the respective databases/sources, it was necessary to construct searches as noted for each of the sources. The search terms are listed below in full for each source and noted if the general search terms or other search terms were used.

"1,1-Dichloropropane" OR "1,2-Dichloropropane" OR "1,3-Dichloropropane" OR "11Dichloropropane" OR "12dichloropropane" OR "13Dichloropropane" OR "2,2-Dichloropropane" OR "22Dichloropropane" OR "alpha,alpha-Dichloropropane" OR "alpha,alpha-Propylene dichloride" OR "alpha,beta-Dichloropropane" OR "alpha,beta-Propylene dichloride" OR "BRN 1718880" OR "BRN 1731152" OR "Caswell No. 324" OR "Dichloro-1,2 propane" OR "Dichlorodimethylmethane" OR "Dicoloropropane" OR "Dimethyldichloromethane" OR "NCI-C55141" OR "NSC 1237" OR "NSC 6204" OR "Propylene chloride" OR "Propylene dichloride" OR "Propylene chloride" OR "Trimethylene dichloride" OR "UNII-AJ1HQ2GUCP" OR "UNII-C5V432N6XB" OR "UNII-RRZ023OFWL" OR "UNII-SR71OVZ2OZ"

<u>**CURRENT CONTENTS CONNECT</u>**: (access.webofknowledge.com) General Search Terms applied to the search strategy for Current Contents.</u>

Date Searched: 05/16/19Date Range of Search: 1970 to Present N = 242

TS=("1,1-Dichloropropane" OR "1,2-Dichloropropane" OR "1,3-Dichloropropane" OR "11Dichloropropane" OR "12dichloropropane" OR "13Dichloropropane" OR "2,2-Dichloropropane" OR "22Dichloropropane" OR "alpha,alpha-Dichloropropane" OR "alpha,alpha-Propylene dichloride" OR "alpha,beta-Dichloropropane" OR "alpha,beta-Propylene dichloride" OR "BRN 1718880" OR "BRN 1731152" OR "Caswell No. 324" OR "Dichloro-1,2 propane" OR "Dichlorodimethylmethane" OR "Dicoloropropane" OR "Dimethyldichloromethane" OR "NCI-C55141" OR "NSC 1237" OR "NSC 6204" OR "Propylene chloride" OR "Propylene dichloride" OR "Trimethylene chloride" OR "Trimethylene dichloride" OR "UNII-AJ1HQ2GUCP" OR "UNII-C5V432N6XB" OR "UNII-RRZ023OFWL" OR "UNII-SR71OVZ2OZ") N = 242

#### **WOS Core Collection:**

Web of Science Core Collection may be accessed through EPA Desktop Library (<u>https://intranet.epa.gov/desktop/databases.htm</u>) by clicking on the Web of Science Link or copying and pasting (https://apps.webofknowledge.com).

Date Searched: 09/10/2019Date Range of Search: 1970 to Present N = 417

TS=("1,1-Dichloropropane" OR "1,2-Dichloropropane" OR "1,3-Dichloropropane" OR "11Dichloropropane" OR "12dichloropropane" OR "13Dichloropropane" OR "2,2-Dichloropropane" OR "22Dichloropropane" OR "alpha,alpha-Dichloropropane" OR "alpha,alpha-Propylene dichloride" OR "alpha,beta-Dichloropropane" OR "alpha,beta-Propylene dichloride" OR "BRN 1718880" OR "BRN 1731152" OR "Caswell No. 324" OR "Dichloro-1,2 propane" OR "Dichlorodimethylmethane" OR "Dicoloropropane" OR "Dimethyldichloromethane" OR "NCI-C55141" OR "NSC 1237" OR "NSC 6204" OR "Propylene chloride" OR "Propylene dichloride" OR "Trimethylene chloride" OR "Trimethylene dichloride" OR "UNII-AJ1HQ2GUCP" OR "UNII-C5V432N6XB" OR "UNII-RRZ023OFWL" OR "UNII-SR71OVZ2OZ") N = 417

#### PROQUEST Agricultural and Scientific Database: (www.csa.com)

General Search Terms applied to the search strategy for ProQuest Agricultural and Scientific Database.

Date Searched: Date Range of Search: 1900 to Present N = 411

ALL("1,1-Dichloropropane" OR "1,2-Dichloropropane" OR "1,3-Dichloropropane" OR "11Dichloropropane" OR "12dichloropropane" OR "13Dichloropropane" OR "2,2-Dichloropropane" OR "22Dichloropropane" OR "alpha,alpha-Dichloropropane" OR "alpha,alpha-Propylene dichloride" OR "alpha,beta-Dichloropropane" OR "alpha,beta-Propylene dichloride" OR "BRN 1718880" OR "BRN 1731152" OR "Caswell No. 324" OR "Dichloro-1,2 propane" OR "Dichlorodimethylmethane" OR "Dicoloropropane" OR "Dimethyldichloromethane" OR "NCI-C55141" OR "NSC 1237" OR "NSC 6204" OR "Propylene chloride" OR "Propylene dichloride" OR "Propylene chloride" OR "Trimethylene dichloride" OR "UNII-AJ1HQ2GUCP" OR "UNII-C5V432N6XB" OR "UNII-RRZ023OFWL" OR "UNII-SR71OVZ2OZ") AND STYPE("Scholarly Journals" OR Reports OR Thesis OR "Government Documents") AND LA(ENG) N = 411

#### **PROQUEST Dissertations and Theses:** (search.proquest.com)

General Search Terms applied to the search strategy for ProQuest Dissertations and Theses.

Date Searched: 05/17/2019Date Range of Search: 1900 to Present N = 5 ALL("1,1-Dichloropropane" OR "1,2-Dichloropropane" OR "1,3-Dichloropropane" OR "11Dichloropropane" OR "12dichloropropane" OR "13Dichloropropane" OR "2,2-Dichloropropane" OR "22Dichloropropane" OR "alpha,alpha-Dichloropropane" OR "alpha,alpha-Propylene dichloride" OR "alpha,beta-Dichloropropane" OR "alpha,beta-Propylene dichloride" OR "BRN 1718880" OR "BRN 1731152" OR "Caswell No. 324" OR "Dichloro-1,2 propane" OR "Dichlorodimethylmethane" OR "Dicoloropropane" OR "Dimethyldichloromethane" OR "NCI-C55141" OR "NSC 1237" OR "NSC 6204" OR "Propylene chloride" OR "Propylene dichloride" OR "Trimethylene chloride" OR "Trimethylene dichloride" OR "UNII-AJ1HQ2GUCP" OR "UNII-C5V432N6XB" OR "UNII-RRZ023OFWL" OR "UNII-SR71OVZ2OZ") AND LA(ENG) N = 5

#### SCIENCE DIRECT: (www.sciencedirect.com)

General Search Terms applied to the search strategy for Science Direct

Date Searched: 05/20/2019Date Range of Search: 1823 to Present N = 134

Science Direct 01:

"1,1-Dichloropropane" OR "1,2-Dichloropropane" OR "1,3-Dichloropropane" OR "11Dichloropropane" OR "12dichloropropane" OR "13Dichloropropane" OR "2,2-Dichloropropane" OR "22Dichloropropane" OR "alpha,alpha-Dichloropropane" N = 131

Science Direct 02:

"alpha,alpha-Propylene dichloride" OR "alpha,beta-Dichloropropane" OR "alpha,beta-Propylene dichloride" OR "BRN 1718880" OR "BRN 1731152" OR "Caswell No. 324" OR "Dichloro-1,2 propane" OR "Dichlorodimethylmethane" OR "Dicoloropropane" N = 1

Science Direct 03:

"Dimethyldichloromethane" OR "NCI-C55141" OR "NSC 1237" OR "NSC 6204" OR "Propylene chloride" OR "Propylene dichloride" OR "Propylidene chloride" OR "Trimethylene chloride" OR "Trimethylene dichloride"

N = 2

Science Direct 04: "UNII-AJ1HQ2GUCP" OR "UNII-C5V432N6XB" OR "UNII-RRZ023OFWL" OR "UNII-SR71OVZ2OZ" N = 0

#### AGRICOLA: (www.nal.usda.gov)

General Search Terms applied to the search strategy for Agricola. The Agricola database contains a significant amount of gray literature including proceedings, symposia, and progress reports from government and educational institutions. Agricola is not used when conducting a search for the Office of Water.

Date Searched: 05/17/2019

Date Range of Search:  $15^{\text{th}}$  century to the Present N = 47

Agricola 01: 1,1-Dichloropropane 1,2-Dichloropropane 1,3-Dichloropropane 11Dichloropropane 12dichloropropane 2,2-Dichloropropane 22Dichloropropane alpha,alpha-Dichloropropane alpha,alpha-Propylene dichloride N = 47

Agricola 02: alpha,beta-Dichloropropane alpha,beta-Propylene dichloride BRN 1718880 BRN 1731152 Caswell No. 324 Dichloro-1,2 propane Dichlorodimethylmethane Dicoloropropane Dimethyldichloromethane NCI-C55141 N = 0

Agricola 03: NSC 1237 NSC 6204 Propylene chloride Propylene dichloride Propylidene chloride Trimethylene chloride Trimethylene dichloride UNII-AJ1HQ2GUCP UNII-C5V432N6XB UNII-RRZ023OFWL N = 0

Agricola 04: UNII-SR71OVZ2OZ N = 0

**TOXNET:** (toxnet.nlm.nih.gov/cgi-bin/sis/htmlgen?TOXLINE)

General Search Terms applied to the search strategy for TOXNET.

Date Searched: 05/20/2019Date Range of Search: 1900 to Present N = 684

TOXNET 01: 78-87-5 OR 78-99-9 OR 142-28-9 OR 594-20-7 OR 26198-63-0

#### PubMed:

PubMed may be accessed through the EPA Desktop Library (<u>https://www.ncbi.nlm.nih.gov/pubmed/</u>)

Date Searched: 07/02/2019Date Range of Search: 1900 to present N = 182

."1,1-Dichloropropane" OR "1,2-Dichloropropane" OR "1,3-Dichloropropane" OR "11Dichloropropane" OR "12dichloropropane" OR "13Dichloropropane" OR "2,2-Dichloropropane" OR "22Dichloropropane" OR "alpha,alpha-Dichloropropane" OR "alpha,alpha-Propylene dichloride" OR "alpha,beta-Dichloropropane" OR "alpha,beta-Propylene dichloride" OR "BRN 1718880" OR "BRN 1731152" OR "Caswell No. 324" OR "Dichloro-1,2 propane" OR "Dichlorodimethylmethane" OR "Dicoloropropane" OR "Dimethyldichloromethane" OR "NCI-C55141" OR "NSC 1237" OR "NSC 6204" OR "Propylene chloride" OR "Propylene dichloride" OR "Propylidene chloride" OR "Trimethylene chloride" OR "Trimethylene dichloride" OR "UNII-AJ1HQ2GUCP" OR "UNII-C5V432N6XB" OR "UNII-RRZ023OFWL" OR "UNII-SR71OVZ2OZ" N = 182

#### **ECOTOX UNIFY:**

This is an internal EPA database that is not accessible to the public. Results from the ECOTOX Unify search strategy.

Date Searched: 05/20/2019Date Range of Search: all years N = 144

12DPA,11DPA,13DPA,22DPA

# A.1.2.2 Data Prioritization for Environmental Hazard, Human Health Hazard, Fate and Physical Chemistry

In brief, SWIFT Review has pre-set literature search strategies ("filters") developed by information specialists that can be applied to identify studies that are more likely to be useful for identifying human health and ecotoxicity content from those that likely do not (*e.g.*, analytical methods). The filters function like a typical search strategy where studies are tagged as belonging to a certain filter if the terms in the filter literature search strategy appear in title, abstract, keyword or medical subject headings (MeSH) fields content. The applied SWIFT Review filters focused on lines of evidence: human, animal models for human health, ecological taxa (which includes ecotoxicological animal models, plants, and other taxa), and *in vitro* studies. The details of the search strategies that underlie the filters are available online. Studies not retrieved using these filters were not considered further. Studies that included one or

more of the search terms in the title, abstract, keyword, or MeSH fields were exported as a RIS file for screening in Swift-ActiveScreener or <u>DistillerSR</u><sup>11</sup>.

#### A.1.2.3 Data Prioritization for Occupational Exposures and Environmental Releases and General Population, Consumer and Environmental Exposures

To prioritize references related to occupational exposure, environmental release, general population exposure, consumer exposure, and environmental exposure, EPA used positive and negative seed studies to build a classification model in SWIFT Review. The positive seeds were identified using relevant literature pool for the first ten TSCA risk evaluations, while the negative seeds were identified from a subset of literature for the current high-priority substances. The model was then applied to the unclassified literature to generate a classification score for each references. Scores above a certain threshold value were then prioritized for further review in SWIFT-ActiveScreener.

### A.2 Peer-Reviewed Screening Process

The studies identified from publicly available database searches and SWIFT-Review filtering/ prioritization were housed in HERO system and imported into SWIFT-ActiveScreener or DistillerSR for title/abstract and full-text screening. Both title/abstract and full-text screening were conducted by two independent reviewers. Screening is initiated with a pilot phase of screening (between 10 and 50) studies to identify areas where clarification in screening criteria might be needed or chemical-specific supplemental material tags might be identified. Records that met PECO (or equivalent criteria (Appendix A.2.1) during title and abstract screening were considered for full-text screening. At both the title/abstract and full-text review levels, screening conflicts were resolved by topic-specific experts and/or discussion among the primary screeners. For citations with no abstract, the articles are initially screened based on all or some of the following: title relevance (titles that suggest a record is not relevant can be excluded rather than marked as unclear), and page numbers (articles two pages in length or less were assumed to be conference reports, editorials, or letters). During title/abstract or full-text level screening in DistillerSR, studies that did not meet the PECO criteria, but which could provide supporting information were categorized (or "tagged") as supplemental information.

It is important to emphasize that being tagged as supplemental material does not mean the study would necessarily be excluded from consideration in an assessment. The initial screening level distinctions between a study meeting the PECO criteria and a supplemental study are often made for practical reasons and the tagging structure (as seen in the literature inventory trees and heat maps in Section 2.1 of this document) are designed to ensure the supplemental studies are categorized for easy retrieval if needed while conducting the assessment. The impact on the assessment conclusions of individual studies tagged as supporting material is often difficult to assess during the screening phase of the assessment. These studies may emerge as being critically important to the assessment and need to be evaluated and summarized at the individual study level (e.g., cancer MOA mechanistic or non-English-language studies), or be helpful to provide context (e.g., summarize current levels of exposure, provide hazard evidence from routes or durations of exposure not pertinent to the PECO), or not be cited at all in the assessment (e.g., individual studies that contribute to a well-established scientific conclusion). Studies maybe be tagged as supplemental material during either title and abstract or full-text screening. When tagged as supplemental material during title and abstract screening, it may not be completely clear whether the chemical of interest is reported in the study (*i.e.*, abstracts may not describe all chemicals investigated). In these cases, studies are still tagged with the expectation that if full-text retrieval is pursued, then additional screening would be needed to clarify if the study is pertinent.

<sup>&</sup>lt;sup>11</sup><u>DistillerSR</u> is a web-based systematic review software used to screen studies available at <u>https://www.evidencepartners.com/products/distillersr-systematic-review-software</u>.

#### A.2.1 Inclusion/Exclusion Criteria

A PECO statement is typically used to focus the research question(s), search terms, and inclusion/exclusion criteria in a systematic review. PECO criteria were developed *a priori* to screening and modified to fit the various discipline areas supporting the TSCA risk evaluations. Variations include the RESO (receptor, exposure, scenario/setting, and outcome) used for the occupational exposure and environmental releases discipline, and PESO (pathways/processes, exposures, setting/scenario, and outcomes) used by the fate and transport discipline. All PECOs and PECO-equivalent criteria can be found in the following sections.

#### A.2.1.1 PECO for Environmental and Human Health Hazards

The PECO used in this evidence map to identify literature pertinent to 1,2-dichloropropane effects on human health and environmental hazard is presented in Table\_Apx A-3. In addition to the PECO criteria, studies containing potentially relevant supplemental material were tracked and categorized during the literature screening process as outlined in Table\_Apx A-4.

Table\_Apx A-3. Hazards Title and Abstract and Full-text PECO Criteria for 1,2-Dichloropropane

PECO Element	Evidence				
Р	<ul> <li>Human: Any population and life stage (e.g., occupational or general population, including children and other sensitive populations).</li> <li>Animal: Aquatic and terrestrial species (live, whole organism) from any life stage (e.g., preconception, in utero, lactation, peripubertal, and adult stages). Tests of the single toxicants in <i>in vitro</i> systems or on live, whole, taxonomically verifiable organisms (e.g., gametes, embryos, or plant or fungal sections capable of forming whole, new organisms) that are not bacteria, humans, monkeys, viruses, or yeast. In most cases, transgenic animal models will get screened as "yes" or "unclear" at the Title and Abstract (TIAB) screening level. Although certain non-mammalian model systems are increasing used to identify potential human health hazards (e.g., Xenopus and zebrafish), for simplicity animal models will be further inventoried according to the categorization below:         <ul> <li><u>Human health models</u>: rat, mouse, rabbit, dog, hamster, guinea pig, cat, non-human primate, pig, hen (neurotoxicity only)</li> <li><u>Environmental models</u>: invertebrates (e.g., insects, spiders, crustaceans, mollusks, and worms) and vertebrates (e.g., mammals and all amphibians, birds, fish, and reptiles). All hen studies (including neurotoxicity studies) will be included for ecotoxicological models.</li> </ul> </li> <li>Plants: All aquatic and terrestrial species (live), including algal, moss, lichen and fungi species.</li> <li>Screener note:         <ul> <li>To identify potential environmental hazards, other organisms not listed above in their respective categories can also be used. Non-mammalian model systems are increasingly used to identify potential human health hazards (e.g., Xenopus, zebrafish), and traditional human health models (e.g., OECD 418 and 419) are considered relevant to both human and eco hazard.</li> <li>PECO considerations should be directed toward effects on target species only and not</li></ul></li></ul>				
Е	Relevant forms and isomers         • 1,2-Dichloropropane (CASRN 78-87-5)         • Related isomers:         • 1,1-Dichloropropane (CASRN 78-99-9)         • 1,3-Dichloropropane (CASRN 142-28-9)         • 2,2-Dichloropropane (CASRN 594-20-7)				

PECO Element	Evidence						
	For synonyms see the EPA Chemistry Dashboard.						
	<ul> <li>Human: Any exposure to 1,2-Dichloropropane.</li> <li>Animal: Any exposure to 1,2-Dichloropropane, including via water, soil or sediment, injection (<i>i.e.</i>, oral or topical), gavage, diet, dermal, and inhalation.</li> <li>Plants: Exposure to 1,2-Dichloropropane via water and/or soil, with reported concentration and duration. Studies involving exposures to mixtures will be included only if they also include exposure to one of these solvents alone.</li> </ul>						
	Screener note: Field studies with media concentrations (surface water, interstitial water, soil) and/or body/tissue concentrations of animals or plants are to be identified as <u>Supplemental</u> if any biological effects are reported.						
	<ul> <li>Human: A comparison or referent population exposed to lower levels (or no exposure/exposure below detection limits) of 1,2-Dichloropropane, or exposure to one of these solvents for shorter periods of time. case-crossover, case-referent, case-only, case-specular, case-cohort, case-parent, nested case-control study designs are all included.</li> <li>Animal and Plants: A concurrent control group exposed to vehicle-only treatment and/or untreated control (control could be a baseline measurement).</li> </ul>						
С	<ul> <li>Screener note:</li> <li>If no control group is explicitly stated or implied (<i>e.g.</i> by mention of statistical results that could only be obtained if a control group was present), the study will be marked as <u>unclear</u> during Title/Abstract Screening.</li> <li>All case reports and case studies/series describing findings in a sample size of less than 20 people in any setting (<i>e.g.</i>, occupation, general population) will be tracked as "potentially relevant supplemental information".</li> </ul>						
0	<ul> <li>Human: All health outcomes (cancer and noncancer).</li> <li>Animal and Plants: All biological effects (including bioaccumulation from laboratory studies with concurrently measured water and tissue concentrations).</li> <li>Screener note:</li> <li>Measurable biological effects relevant for humans, animals and plants may include but are not limited to: mortality, behavioral, population, cellular, physiological, growth, reproduction of an acceptable organism to a chemical toxicant.</li> </ul>						

#### Table\_Apx A-4. Major Categories of Potentially Relevant Supplemental Materials for 1,2-Dichloropropane

Category	Evidence		
Mechanistic studies	Studies reporting measurements related to a health outcome that inform the biological or chemical events associated with phenotypic effects, in both mammalian and non-mammalian model systems, including <i>in vitro</i> , <i>in vivo</i> (by various non-inhalation routes of exposure), ex vivo, and <i>in silico</i> studies.		
ADME, PBPK, and toxicokinetic	Studies designed to capture information regarding absorption, distribution, metabolism, and excretion (ADME), toxicokinetic studies, or physiologically based pharmacokinetic (PBPK) models.		
Susceptible populations (no health outcome)	Studies that identify potentially susceptible subgroups; for example, studies that focus on a specific demographic, life stage, or genotype.		
Mixture studies	Mixture studies that are not considered PECO-relevant because they do not contain an exposure or treatment group assessing only the chemical of interest.		

Category	Evidence
IL ASE FENALIS AF PASE SELIES	Case reports ( $n \le 3$ cases) and case series/studies (<20 cases) will be tracked as potentially relevant supplemental information.
	Records that do not contain original data, such as other agency assessments, informative scientific literature reviews, editorials or commentaries.
IC ONTERENCE ADSTRACTS	Records that do not contain sufficient documentation to support study evaluation and data extraction.
RIEIA SILIAIES	Field studies where there are accompanying body/tissue concentrations of animals without any biological effects reported

#### A.2.1.2 PECO for Consumer, Environmental, and General Population Exposures

# Table\_Apx A-5. Generic Inclusion Criteria for the Data Sources Reporting Exposure Data on General Population, Consumers and Environmental Receptors

PECO Element	Evidence
<b>P</b> opulation	<b>Human:</b> General population; consumers; bystanders in the home; near-facility populations (includes industrial and commercial facilities manufacturing, processing, or using the chemical substance); children; susceptible populations (life stages, preexisting conditions, genetic factors); pregnant women; lactating women; women of child bearing age. Many human population groups may be exposed. No chemical-specific exclusions are suggested at this time.
	<b>Environmental:</b> aquatic species, terrestrial species, terrestrial plants, aquatic plants (field studies only)
	Expected Primary Exposure Sources, Pathways, Routes:
<u>E</u> xposure	<u>Pathways:</u> indoor air/vapor/mist; indoor dust; particles; outdoor/ambient air; surface water; biosolids; sediment; breastmilk; food items containing 1,2-dichloropropane including fish; consumer product uses in the home (including consumer product containing chemical);
	Routes of Exposure: Inhalation, Oral, Dermal
Comparator	<b>Human:</b> Consider media-specific background exposure scenarios and use/source specific exposure scenarios as well as which receptors are and are not reasonably exposed across the projected exposure scenarios.
(Scenario)	<b>Environmental</b> Consider media-specific background exposure scenarios and use/source specific exposure scenarios as well as which receptors are and are not reasonably exposed across the projected exposure scenarios.
<u>O</u> utcomes for Exposure Concentration or	<b>Human:</b> Acute, subchronic, and/or indoor air and water concentration estimates (mg/m <sup>3</sup> or mg/L). Both external potential dose and internal dose based on biomonitoring and reverse dosimetry mg/kg/day will be considered. Characteristics of consumer products or articles (weight fraction, emission rates, etc) containing 1,2-dichloropropane.
Dose	<b>Environmental:</b> A wide range of ecological receptors will be considered (range depending on available ecotoxicity data) using surface water concentrations, sediment concentrations.

Chemical	Drinking Water	Ambient Air	Air Disposal	Land Disposal	Underground Disposal	Ground Water
1,2-Dichloropropane	Х	Х	Х	Х	Х	Х

#### Table\_Apx A-6. Pathways Identified as Supplemental for 1,2-Dichloropropane<sup>a</sup>

<sup>a</sup> Supplemental pathways refer to pathways addressed by other EPA administered statutes (see Section 2.6.3.1). Studies tagged under these pathways provide media information that is not prioritized in the screening process.

#### A.2.1.3 RESO for Occupational Exposure and Environmental Releases

EPA developed a generic RESO statement to guide the screening of engineering and occupational exposure data or information sources for the TSCA risk evaluations. Data or information sources that comply with the inclusion criteria specified in the RESO statement are eligible for inclusion, considered for evaluation, and possibly included in the environmental release and occupational exposure assessments. On the other hand, data or information sources that fail to meet the criteria in the RESO statement are excluded from further consideration.

Assessors seek information on various chemical-specific engineering and occupational exposure data needs as part of the process of developing the exposure assessment for each risk evaluation. EPA uses the RESO statement (Table\_Apx A-7) along with the information in Table\_Apx A-8 when screening the engineering and occupational exposure data and information.

<b>RESO Element</b>	Evidence
<u>R</u> eceptors	<u>Humans</u> : Workers, including occupational non-users
	• <u>Environment</u> : All environmental receptors (relevant release estimates input to Exposure)
	Please refer to the conceptual models for more information about the environmental and human receptors included in the TSCA risk evaluation.
<u>E</u> xposure	<ul> <li>Worker exposure to and relevant environmental releases of the chemical substance from occupational scenarios:         <ul> <li>Dermal and inhalation exposure routes (as indicated in the conceptual model)</li> <li>Oral route (as indicated in the conceptual model)</li> </ul> </li> </ul>
	Please refer to the conceptual models for more information about the routes and media/pathways included in the TSCA risk evaluation.
<u>S</u> etting or <u>Scenario</u>	• Any occupational setting or scenario resulting in worker exposure and relevant environmental releases (includes all manufacturing, processing, use, disposal.
<u>O</u> utcomes	<ul> <li>Quantitative estimates* of worker exposures and of relevant environmental releases from occupational settings</li> <li>General information and data related and relevant to the occupational estimates*</li> <li>(kg/day or mg/m<sup>3</sup> for worker exposures kg/site/day for releases) are determined by toxicologists for</li> </ul>

 Table\_Apx A-7. Inclusion Criteria for Data Sources Reporting Engineering and Occupational

 Exposure Data

\* Metrics (*e.g.*, mg/kg/day or mg/m<sup>3</sup> for worker exposures, kg/site/day for releases) are determined by toxicologists for worker exposures and by exposure assessors for releases; also, the Engineering, Release and Occupational Exposure Data Needs (Table\_Apx A-8) provides a list of related and relevant general information. TSCA=Toxic Substances Control Act

Table_Apx A-8. Engineering, Environmental Release and Occupational Data Necessary to
Develop the Environmental Release and Occupational Exposure Assessments

Objective Determined during Scoping	Type of Data <sup>a</sup>
General Engineering Assessment (may apply to Occupational Exposures and / or Environmental Releases)	<ul> <li>Description of the life cycle of the chemical(s) of interest, from manufacture to end-of-life (<i>e.g.</i>, each manufacturing, processing, or use step), and material flow between the industrial and commercial life cycle stages.</li> <li>The total annual U.S. volume (lb/yr or kg/yr) of the chemical(s) of interest manufactured, imported, processed, and used; and the share of total annual manufacturing and import volume that is processed or used in each life cycle step.</li> <li>Description of processes, equipment, and unit operations during each industrial/ commercial life cycle step.</li> <li>Material flows, use rates, and frequencies (lb/site-day or kg/site-day and days/yr; lb/site-batch and batches/yr) of the chemical(s) of interest during each industrial/ commercial life cycle step. Note: if available, include weight fractions of the chemicals (s) of interest and material flows of all associated primary chemicals (especially water).</li> <li>Number of sites that manufacture, process, or use the chemical(s) of interest for each industrial/ commercial life cycle step and site locations.</li> </ul>
Occupational Exposures	<ul> <li>Concentration of the chemical of interest</li> <li>Description of worker activities with exposure potential during the manufacture, processing, or use of the chemical(s) of interest in each industrial/commercial life cycle stage.</li> <li>Potential routes of exposure (<i>e.g.</i>, inhalation, dermal).</li> <li>Physical form of the chemical(s) of interest for each exposure route (<i>e.g.</i>, liquid, vapor, mist) and activity.</li> <li>Breathing zone (personal sample) measurements of occupational exposures to the chemical(s) of interest, measured as time-weighted averages (TWAs), short-term exposures, or peak exposures in each occupational life cycle stage (or in a workplace scenario similar to an occupational life cycle stage).</li> <li>Area or stationary measurements of airborne concentrations of the chemical(s) of interest in each occupational setting and life cycle stage (or in a workplace scenario similar to the life cycle stage of interest).</li> <li>For solids, bulk and dust particle size characterization data.</li> <li>Dermal exposure data.</li> <li>Exposure frequency (days/yr).</li> <li>Number of workers who potentially handle or have exposure to the chemical(s) of interest in each occupational life cycle stage.</li> <li>PPE types employed by the industries within scope.</li> <li>ECs employed to reduce occupational exposures in each occupational life cycle stage (or in a workplace scenario similar to the life cycle stage (or in a workplace scenario similar to the life cycle stage of interest).</li> </ul>
Environmental Releases (to relevant environmental media)	<ul> <li>Description of sources of potential environmental releases, including cleaning of residues from process equipment and transport containers, involved during the manufacture, processing, or use of the chemical(s) of interest in each life cycle stage.</li> <li>Estimated mass (lb or kg) of the chemical(s) of interest released from industrial and commercial sites to each environmental medium (water) and treatment and disposal methods (POTW), including releases per site and aggregated over all sites (annual release rates, daily release rates)</li> <li>Release or emission factors.</li> <li>Number of release days per year.</li> <li>Waste treatment methods and pollution control devices employed by the industries within scope and associated data on release/emission reductions.</li> </ul>
	s included in the full-text screening form. The screener makes a selection from these specific tags, which ific types of data or information.

Objective Determined	Type of Data <sup>a</sup>					
during Scoping	Type of Data					
In addition to the d	ata types listed above, EPA may identify additional data needs for mathematical modeling. These data					
needs will be deter	mined on a case-by-case basis.					
Abbreviations:						
hr=Hour						
kg=Kilogram(s)	=Kilogram(s)					
lb=Pound(s)						
yr=Year						
PV=Particle volum	e					
POTW=Publicly ov	wned treatment works					
PPE=Personal prot	ection equipment					
PSD=Particle size distribution						
TWA=Time-weigh	ted average					

#### A.2.1.4 PESO for Fate and Transport

EPA developed a generic PESO statement to guide the screening of environmental fate data or information sources for the TSCA risk evaluations. Data or information sources that comply with the inclusion criteria in the PESO statement are eligible for inclusion, considered for evaluation, and possibly included in the environmental fate assessment. On the other hand, data or information sources that fail to meet the criteria in the PESO statement are excluded from further consideration.

Assessors seek information on various chemical-specific fate endpoints and associated fate processes, environmental media and exposure pathways as part of the process of developing the environmental fate assessment for each risk evaluation. EPA uses the PESO statement (Table\_Apx A-9) along with the information in Table\_Apx A-10 when screening the fate data or information sources to ensure complete coverage of the processes, pathways and data or information relevant to the environmental fate and transport of the chemical substance undergoing risk evaluation.

# Table\_Apx A-9. Inclusion Criteria for Data or Information Sources Reporting Environmental Fate and Transport Data

PESO Element	Evidence
<u>P</u> athways and <u>P</u> rocesses	<ul> <li>Environmental fate, transport, partitioning and degradation behavior across environmental media to inform exposure pathways of the chemical substance of interest</li> <li>Exposure pathways included in the conceptual models: air, surface water, groundwater, wastewater, soil, sediment and biosolids.</li> <li>Processes associated with the target exposure pathways</li> <li>Bioconcentration and bioaccumulation</li> <li>Destruction and removal by incineration</li> <li>Please refer to the conceptual models for more information about the exposure pathways included in each TSCA risk evaluation.</li> </ul>
<u>E</u> xposure	<ul> <li>Environmental exposure of environmental receptors (<i>i.e.</i>, aquatic and terrestrial organisms) to the chemical substance of interest, mixtures including the chemical substance, and/or its degradation products and metabolites</li> <li>Environmental exposure of human receptors, including any potentially exposed or susceptible subpopulations, to the chemical substance of interest, mixtures including the chemical substance, and/or its degradation products and metabolites</li> <li>Please refer to the conceptual models for more information about the environmental and human receptors included in each TSCA risk evaluation.</li> </ul>
<u>S</u> etting or <u>S</u> cenario	Any setting or scenario resulting in releases of the chemical substance of interest into the natural or built environment ( <i>e.g.</i> , buildings including homes or workplaces, or wastewater treatment facilities) that would expose environmental ( <i>i.e.</i> , aquatic and terrestrial organisms) or human receptors ( <i>i.e.</i> , general population, and potentially exposed or susceptible subpopulation)
<u>O</u> utcomes	Fate properties which allow assessments of exposure pathways: Abiotic and biotic degradation rates, mechanisms, pathways, and products Bioaccumulation magnitude and metabolism rates Partitioning within and between environmental media (see Pathways and Processes)

# Table\_Apx A-10. Fate Endpoints and Associated Processes, Media and Exposure Pathways Considered in the Development of the Environmental Fate Assessment

		Associated Media/Exposure Pathways			
Fate Data Endpoint	Associated Process(es)	Surface Water, Wastewater, Sediment	Soil, Biosolids	Groundwater	Air
<b>Required Environmental Fate</b>	Data		-		-
Abiotic reduction rates or half- lives	Abiotic reduction, Abiotic dehalogenation	Х			
Aerobic biodegradation rates or half-lives	Aerobic biodegradation	Х	Х		
Anaerobic biodegradation rates or half-lives	Anaerobic biodegradation	Х	Х	Х	
Aqueous photolysis (direct and indirect) rates or half-lives	Aqueous photolysis (direct and indirect)	Х			

		Associate	ed Media/Exj	posure Pathways	
Fate Data Endpoint	Associated Process(es)	Surface Water, Wastewater, Sediment	Soil, Biosolids	Groundwater	Air
Atmospheric photolysis (direct and indirect) rates or half-lives	Atmospheric photolysis (direct and indirect)				X
Bioconcentration factor (BCF), Bioaccumulation factor (BAF)	Bioconcentration, Bioaccumulation	X	Х		X
Biomagnification and related information	Trophic magnification	Х			
Desorption information	Sorption, Mobility	X	Х	Х	
Destruction and removal by incineration	Incineration				Х
Hydrolysis rates or half-lives	Hydrolysis	X	Х	Х	
K <sub>OC</sub> and other sorption information	Sorption, Mobility	X	Х	Х	
Wastewater treatment removal information	Wastewater treatment	Х	Х		
Supplemental (or Optional) En	vironmental Fate Data			F	
Abiotic transformation products	Hydrolysis, Photolysis, Incineration	X			X
Aerobic biotransformation products	Aerobic biodegradation	X	Х		
Anaerobic biotransformation products	Anaerobic biodegradation	X	Х	Х	
Atmospheric deposition information	Atmospheric deposition				X
Coagulation information	Coagulation, Mobility	X		Х	
Incineration removal information	Incineration				X

#### A.2.1.5 Generation of Hazard Heat Maps

As stated in Appendix A.1.2.2, SWIFT Review has pre-set literature search strategies ("filters") developed by information specialists that can be applied to identify studies that are more likely to be useful for identifying human health and ecotoxicity content. The filters function like a typical search strategy where studies are tagged as belonging to a certain filter if the terms in the filter literature search strategy appear in title, abstract, keyword or MeSH fields content.

After the completion of full-text screening for hazard data, all references tagged as included (or "PECOrelevant) were uploaded to the SWIFT Review tool for further filtering. The SWIFT Review filters applied at this phase focused on types of health outcomes included: "ADME", "PBPK", "cancer", "cardiovascular", "developmental", "endocrine", "gastrointestinal", "hematological and immune", "hepatic", "mortality", "musculoskeletal", "neurological", "nutritional and metabolic", "ocular and sensory", "renal", "reproductive", "respiratory", and "skin and connective tissue". The details of these health outcome search strategies that underlie the filters are available <u>online</u>. Studies that included one or more of the search terms in the title, abstract, keyword, or MeSH fields were exported and used to populate the Hazard Heat Map (Figure 2-10). Studies that were not retrieved using these filters were tagged as "No Tag". The evidence type listed in the heat map (*e.g.*, human, animal-human health model, animal- environmental model, and plant) was manually assigned to each reference by screeners during the full-text screening.

The health outcome tags were originally designed for vertebrate systems, and as such, did not conform well to plant evidence. Therefore, any plant studies tagged for: "cancer", "cardiovascular", "gastrointestinal", "hematological and immune", "hepatic", "musculoskeletal", "neurological", "ocular and sensory" and "renal and respiratory" were manually reviewed and re-tagged to more appropriate health outcomes.


## A.3 Gray Literature Search and Screening Strategies

EPA conducted a gray literature search for available information to support the TSCA risk evaluations for the next twenty TSCA risk evaluations. Gray literature is defined as the broad category of data/information sources not found in standard, peer-reviewed literature databases (*e.g.*, PubMed and Web of Science). Gray literature includes data/information sources such as white papers, conference proceedings, technical reports, reference books, dissertations, information on various stakeholder websites, and other databases. Given the nature of how gray literature is searched and collected, results may not come with a bibliographic citation or abstract and were therefore processed using a decision tree logic described in Appendix A.3.1 for potential relevance prior to entering full text screening where a discipline-specific PECO is applied.

Search terms were variable dependent on source and based on knowledge of a given source to provide discipline-specific information. A summary of sources is provided in Appendix A.3.4. The criteria for determining the potential relevance of documents identified from gray literature sources is described in the following sections for each discipline.

#### A.3.1 Screening of Gray Literature

To reduce the overall burden of processing gray literature results, EPA developed a screening process to determine the potential relevance of gray literature. This step was introduced prior to collecting the resulting documents. Figure\_Apx A-1 describes the decision logic used to screen gray literature results.



Figure\_Apx A-1. Decision Logic Tree Used to Screen Gray Literature Results

#### A.3.2 Initial Screening of Sources using Decision Logic Tree

The purpose of the inclusion/exclusion decision logic tree in Figure\_Apx A-1 is to provide a broad, general screening technique to determine whether each gray literature source should be included and further screened or excluded with no additional screening necessary. The diamonds in the decision tree require analysis by the screener, whereas the rectangular boxes are used to classify the type of source. All the questions used in the decision process are provided in Table\_Apx A-11.

Step	Metric	Questions to Consider
1	Potential Relevance	Does the result have information (qualitative or quantitative) related to TSCA risk evaluations? *Apply Discipline relevancy metric
2.1.1		Is it a secondary data source (assessment, robust summary, TSCA submission databases, etc.)?
2.1.2		Is the document from a peer reviewed/published journal?
2.2		Is there an established procedure for data collection, communication, peer review, and/or reporting?
2.2.1	Complete / Available	Has the data been provided by a US governmental/state source?
2.2.2		Has the data been provided by an international governmental source?
2.3		Are these data publicly available/accessible?
2.3.1		Is the source TSCA CBI, proprietary, TSCA or NGO stakeholder submission?
3	Duplicate	Does the result contain any duplicative information found in other sources?

Table\_Apx A-11. Decision Logic Tree Overview

Results of the gray literature search and decision tree process are included in Appendix A.3.4.

#### A.3.3 TSCA Submission Searching and Title Screening

EPA screens information submitted under TSCA Sections 4, 5, 8(e), and 8(d), as well as for your information (FYI) submissions. In the gray literature process defined in Appendix A.3.2, EPA considers the databases that contain TSCA submissions to be secondary sources (Step 1.1) because the metadata in the databases are secondary. These databases then advance to Step 2.3.1 and then to Process C. The Process C steps are described here.

EPA first screens the titles using two screeners per title. EPA conducts this step primarily to reduce the number of full studies to be obtained because some studies are available only on microfiche or in long-term storage. Screening is done using the inclusion and exclusion criteria within the relevant PECOs, PESOs or RESOs for each topic area (Appendix A.2.1). EPA excludes interim reports (*e.g.*, interim sacrifices for toxicity studies) and only final reports are further considered. If the title is not clear regarding the document's contents, EPA obtains the full text and advances to the next steps.

After full texts are obtained, EPA plans to review\_some sources (prior to full-text screening) based on whether they have several factors; primary data, an established procedure for peer review, data collection, communication and/or reporting and are publicly available. Sources that have these factors will move on to full text screening. Other sources will go straight to full text screening using PECO-type criteria without going through this extra step.

EPA may decide to initiate a backwards search on sources that are deemed to have secondary data. In situations where parameters such as procedures for peer review and data collection are unclear, EPA may reach out to the authors to retrieve information to gauge whether the source should be included or excluded. Studies that are not publicly available (such as proprietary or CBI sources) may undergo additional screening steps.

During the full-text screening step, two individuals screen each source according to the PECOs, PESOs and RESOs (Appendix A.2.1).

Results of the TSCA submission search and decision tree process are included in Appendix A.3.4.

#### A.3.4 Gray Literature Search Results for 1,2-Dichloropropane

Table\_Apx A-12 provides a list of gray literature sources that yielded results for 1,2-dichloropropane.

Source Agency	Source Name	Source Type	Source Category	Source Website
ATSDR	ATSDR Tox Profile Updates and Addendums	Other US Agency Resources	Assessment or Related Document	https://www.atsdr.cdc.gov/tox profiles/profilesaddenda.asp
ATSDR	ATSDR Toxicological Profiles (original publication)	Other US Agency Resources	Assessment or Related Document	https://www.atsdr.cdc.gov/tox profiles/index.asp
Australian Government Department of Health	NICNAS Assessments (human health, Tier I, II or III)	International Resources	Assessment or Related Document	https://www.nicnas.gov.au/che mical-information/imap- assessments/imap-assessments
CAL EPA	Technical Support Documents for regulations: Drinking Water Public Health Goals	Other US Agency Resources	Assessment or Related Document	https://oehha.ca.gov/chemicals
CAL EPA	Technical Support Documents for regulations: Proposition 65, Cancer	Other US Agency Resources	Assessment or Related Document	https://oehha.ca.gov/chemicals
CDC	CDC Biomonitoring Tables	Other US Agency Resources	Database	www.cdc.gov/exposurereport/i ndex.html/
ECHA	ECHA Documents	International Resources	Assessment or Related Document	https://echa.europa.eu/informat ion-on-chemicals
ECHA	European Union Risk Assessment Report	International Resources	Assessment or Related Document	https://echa.europa.eu/informat ion-on-chemicals/information- from-existing-substances- regulation
Env Canada	Chemicals at a Glance (fact sheets)	International Resources	Assessment or Related Document	https://www.canada.ca/en/heal th-canada/services/chemical- substances/fact- sheets/chemicals-glance.html
EPA	OPPT: TSCATS database maintained at SRC (TSCA submissions)	US EPA Resources	Database	
EPA	OPPT: Chemview (TSCA submissions - chemical test rule data and substantial risk reports)	US EPA Resources	Database	https://chemview.epa.gov/che mview

 Table\_Apx A-12. Gray Literature Sources that Yielded Results for 1,2-Dichloropropane

Source Agency	Source Name	Source Type	Source Category	Source Website
EPA	OPPT: CIS (CBI LAN) (TSCA submissions)	US EPA Resources	Database	
EPA	Office of Air: AQS, Annual	US EPA Resources	Database	https://aqs.epa.gov/aqsweb/air data/download_files.html#Ann ual
EPA	Office of Air: National Emissions Inventory (NEI) - National Emissions Inventory (NEI) Data (2014, 2011, 2008)	US EPA Resources	Database	https://www.epa.gov/air- emissions-inventories/2014- national-emissions-inventory- nei-data
EPA	Office of Water: STORET and WQX	US EPA Resources	Database	https://www.waterqualitydata. us/portal/
EPA	PPRTV Derivation Support Document	US EPA Resources	Assessment or Related Document	https://hhpprtv.ornl.gov/quickv iew/pprtv_papers.php
EPA	EPA Office of Water: Ambient Water Quality Criteria documents	US EPA Resources	Assessment or Related Document	www.epa.gov/wqc/
EPA	IRIS Summary	US EPA Resources	Assessment or Related Document	https://cfpub.epa.gov/ncea/iris _drafts/atoz.cfm?list_type=alp ha
EPA	Office of Air: TRI	US EPA Resources	Database	https://www.epa.gov/toxics- release-inventory-tri- program/tri-data-and-tools
EPA	Office of Water: Drinking Water Standards Health Effects Support Documents	US EPA Resources	Regulatory Document or List	https://www.epa.gov/eg
EPA	Office of Water: CFRs	US EPA Resources	Regulatory Document or List	https://www.epa.gov/eg
EPA	Office of Air: CFRs and Dockets	US EPA Resources	Regulatory Document or List	https://www.epa.gov/stationar y-sources-air-pollution
IARC	IARC Monograph	International Resources	Assessment or Related Document	http://monographs.iarc.fr/ENG /Monographs/PDFs/index.php
ILO	International Chemical Safety Cards (ICSCs)	International Resources	Database	https://www.ilo.org/safework/i nfo/publications/WCMS_1131 34/langen/index.htm
Japan	Japanese Ministry of the Environment Assessments - Environmental Risk Assessments (Class I Designated Chemical Substances Summary Table)	International Resources	Regulatory Document or List	https://www.env.go.jp/en/che mi/prtr/substances/

Source Agency	Source Name	Source Type	Source Category	Source Website
KOECT	Kirk-Othmer Encyclopedia of Chemical Technology Journal Article	Other Resource	Encyclopedia	https://onlinelibrary.wiley.com/do i/book/10.1002/0471238961
NIOSH	CDC NIOSH - Occupational Health Guideline Documents	Other US Agency Resources	Assessment or Related Document	www.cdc.gov/niosh/topics/che mical.html/
NIOSH	CDC NIOSH - Pocket Guide	Other US Agency Resources	Database	https://www.cdc.gov/niosh/np g/default.html
NLM	National Library of Medicine's PubChem	Other US Agency Resources	Database	https://pubchem.ncbi.nlm.nih.g
NLM	National Library of Medicine's HazMap	Other US Agency Resources	Database	https://haz-map.com/
NLM	National Library of Medicine's Hazardous Substance Databank	Other US Agency Resources	Database	https://www.nlm.nih.gov/datab ases/download/hsdb.html
NTP	Additional NTP Reports	Other US Agency Resources	Assessment or Related Document	https://ntp.niehs.nih.gov/public ations/index.html
NTP	Technical Reports	Other US Agency Resources	Assessment or Related Document	https://ntp.niehs.nih.gov/public ations/reports/index.html?type =Technical+Report
OECD	OECD SIDS	International Resources	Assessment or Related Document	https://hpvchemicals.oecd.org/ ui/Publications.aspx
OECD	OECD: General Site	International Resources	General Search	https://www.oecd.org/
OECD	OECD Emission Scenario Documents	International Resources	Assessment or Related Document	http://www.oecd.org/document /46/0,2340,en 2649 201185 2 412462_1_1_1_1,00.html
OSHA	OSHA Chemical Exposure Health Data	Other US Agency Resources	Database	www.osha.gov/opengov/health samples.html/
RIVM	RIVM Reports: Risk Assessments	International Resources	Assessment or Related Document	https://www.rivm.nl/en
TERA	Toxicology Excellence for Risk Assessment	Other Resources	Assessment or Related Document	www.tera.org/

## Appendix B PHYSICAL AND CHEMICAL PROPERTIES OF 1,2-DICHLOROPROPANE

Table\_Apx B-1 summarizes statistics for the physical and chemical property values identified through systematic review as of June 2020. The "N" column indicates the number of unique primary sources of data for that endpoint. That is, if multiple sources presented equivalent values and cited the same primary source, only one of those was included in these statistics and included in the statistical calculations. All physical and chemical property values that were extracted and evaluated as of June 2020 are presented in the supplemental file *Data Extraction and Data Evaluation Tables for Physical and Chemical Property Studies* (EPA-HQ-OPPT-2018-0428).

Property or Endpoint	N	Unit	Mean	Standard Deviation	Min	Max
Molecular formula	-	-	NA	NA	NA	NA
Molecular weight	-	g/mol	NA	NA	NA	NA
Physical state	4	-	NA	NA	NA	NA
Physical properties	4	-	NA	NA	NA	NA
Melting point	12	°C	-97.7	8.7	-101	-70
Boiling point	16	°C	96.0	0.96	94.5	98.5
Density	7	g/cm <sup>3</sup>	1.153	0.012	1.13	1.16
Vapor pressure	6	mm Hg	48.1	6.38	40	53.3
Vapor density	1	-	3.9	-	3.9	3.9
Water solubility	6	mg/L	2788	114.6	2677	3000
Octanol/water partition coefficient (log Kow)	3	-	1.99	0.006	1.98	1.99
Henry's Law constant	2	atm·m <sup>3</sup> /mol	0.0574	0.0772	0.00282	0.112
Flash point	6	°C	13.3	18.5	-15.6	40
Auto flammability	0	°C	-	-	-	-
Viscosity	0	cP	-	-	-	-
Refractive index	6	-	1.4429	0.0123	1.4364	1.468
Dielectric constant	1	-	8.92	-	8.92	8.92

Table\_Apx B-1. Summary Statistics for Reviewed Physical Properties

NA = Not applicable

## Appendix C ENVIRONMENTAL FATE AND TRANSPORT PROPERTIES OF 1,2-DICHLOROPROPANE

Table\_Apx C-1 provides the environmental fate characteristics that EPA identified and considered in developing the scope for 1,2-dichloropropane. This information was presented in the *Proposed Designation of 1,2-Dichloropropane (CASRN 78-87-5) as a High-Priority Substance for Risk Evaluation* (U.S. EPA, 2019e) and may be updated as EPA collects additional information through systematic review methods.

Property or Endpoint	Value <sup>a</sup>	Reference
Direct Photodegradation	Not expected to be susceptible to direct photolysis by sunlight because the chemical structure of 1,2-dichloropropane does not contain chromophores that absorb at wavelengths >290 nm	<u>NLM (2019)</u> citing <u>U.S.</u> <u>EPA (1979)</u>
	Vapor-phase photolysis under simulated sunlight did not occur after prolonged exposure	<u>NLM (2019)</u> citing <u>Cohen</u> <u>SZ et al. (1984)</u>
Indirect Photodegradation	$t_{1/2} = 24$ days (based on $\cdot$ OH reaction rate constant of $4.4 \times 10^{-13}$ cm <sup>3</sup> /molecule second at 25 °C and an OH concentration of $1.5 \times 10^{6}$ OH/cm <sup>3</sup> ; estimated) <sup>b</sup>	<u>U.S. EPA (2012b)</u>
Hydrolysis	$t_{1/2} = 15.8$ years (rate constant of $5.0 \times 10^{-6}$ hours <sup>-1</sup> at pH 7–9 and 25 °C)	<u>OECD (2006)</u> citing <u>Mackay et al. (1993)</u>
Biodegradation (Aerobic)	0%/14 days based on biological oxygen demand (Japanese MITI test)	<u>NLM (2019)</u> citing <u>CITI</u> (1992)
	$t_{1/2} = 52$ days in a closed system with fresh soil at 15 °C	<u>NLM (2019)</u> citing <u>Van</u> <u>Dijk (1980)</u>
Wastewater Treatment	59% total removal (12% by biodegradation, 1.1 by sludge and 46% by volatilization to air; estimated) <sup>b</sup>	<u>U.S. EPA (2012b)</u>
Bioconcentration Factor	1.2–3.2 ( <i>Cyprinus caprio</i> at 0.4 mg/L test substance concentration); 0.5–6.9 ( <i>C. caprio</i> at 0.04 mg/L test substance concentration)	<u>SYKE (2018); OECD</u> (2006) citing <u>Howard</u> (1990); <u>Mackay et al.</u> (1993)
Bioaccumulation Factor	7.1 (estimated) <sup>b</sup>	<u>U.S. EPA (2012b)</u>
Soil Organic Carbon:Water Partition Coefficient (Log K <sub>OC</sub> )	1.67 (K <sub>OC</sub> = 47) measured in silt loam	<u>NLM (2019)</u> citing <u>Chiou</u> et al. (1979)

Table_Apx C-1. Environmental Fate and Transport Properties of 1,2-Dichloropropane
---

<sup>a</sup> Measured unless otherwise noted; <sup>b</sup>EPI Suite<sup>TM</sup> physical property inputs: Log  $K_{OW} = 1.98$ , BP = 95.5 °C, MP = -100 °C, VP = 53.3 mm Hg at 25 °C, WS = 2,800 mg/L, HLC =  $2.82 \times 10^{-3}$  atm·m<sup>3</sup>/mol at 25 °C, BioP 120, BioA 30, BioS 30, SMILES ClCC(Cl)C; ·OH = hydroxyl radical; OECD = Organisation for Economic Co-operation and Development; TG = test guideline; GC = gas chromatography; MITI = Ministry of International Trade and Industry; BCF = bioaccumulation factor; BOD = biochemical oxygen demand; HPLC = high performance liquid chromatography

#### **Appendix D REGULATORY HISTORY**

The chemical substance, 1,2-dichloropropane, is subject to federal and state laws and regulations in the United States Table\_Apx D-1 and Table\_Apx D-2. Regulatory actions by other governments, tribes and international agreements applicable to 1,2-dichloropropane are listed in Table\_Apx D-3.

#### **Federal Laws and Regulations D.1**

Statutes/Regulations	Description of Authority/Regulation	Description of Regulation
EPA Statutes/Regulations		
Toxic Substances Control Act (TSCA) – Section 6(b)	EPA is directed to identify high- priority chemical substances for risk evaluation; and conduct risk evaluations on at least 20 high priority substances no later than three and one- half years after the date of enactment of the Frank R. Lautenberg Chemical Safety for the 21st Century Act.	1,2-Dichloropropane is one of the 20 chemicals EPA designated as a High-Priority Substance for risk evaluation under TSCA ( <u>84 FR 71924</u> , December 30, 2019). Designation of 1,2- dichloropropane as high-priority substance constitutes the initiation of the risk evaluation on the chemical.
Toxic Substances Control Act (TSCA) – Section 8(a)	The TSCA Section 8(a) CDR Rule requires manufacturers (including importers) to give EPA basic exposure-related information on the types, quantities and uses of chemical substances produced domestically and imported into the United States.	1,2-Dichloropropane manufacturing (including importing), processing and use information is reported under the CDR rule (85 FR 20122, April 2, 2020).
Toxic Substances Control Act (TSCA) – Section 8(b)	EPA must compile, keep current and publish a list (the TSCA Inventory) of each chemical substance manufactured (including imported), or processed, in the United States.	1,2-Dichloropropane was on the initial TSCA Inventory and therefore was not subject to EPA's new chemicals review process under TSCA Section 5 (60 FR 16309, March 29, 1995).
Toxic Substances Control Act (TSCA) – Section 8(e)	Manufacturers (including importers), processors, and distributors must immediately notify EPA if they obtain information that supports the conclusion that a chemical substance or mixture presents a substantial risk of injury to health or the environment.	One risk report received for 1, 2-dichloropropane received: 1993) (U.S. EPA, <u>ChemView</u> . Accessed [April 11, 2019]).

1 T . . . 1 5 1 ... 

Statutes/Regulations	Description of Authority/Regulation	Description of Regulation
Toxic Substances Control Act (TSCA) – Section 4	Provides EPA with authority to issue rules and orders requiring manufacturers (including importers) and processors to test chemical substances and mixtures.	Eleven chemical data submissions from test rules were received for 1,2- dichloropropane: Acute aquatic plant toxicity (2), Acute aquatic toxicity (1), Chronic aquatic toxicity (2), Developmental toxicity (2), Metabolism and Pharmacokinetics (3), Mutagenicity/Genetic toxicity (1), Neurological toxicity (2), Reproductive toxicity (1). (Submissions were received between 1988-2005) (U.S. EPA, <u>ChemView</u> . Accessed April 18, 2019).
Emergency Planning and Community Right-To-Know Act (EPCRA) – Section 313	Requires annual reporting from facilities in specific industry sectors that employ 10 or more full-time equivalent employees and that manufacture, process or otherwise use a TRI-listed chemical in quantities above threshold levels. A facility that meets reporting requirements must submit a reporting form for each chemical for which it triggered reporting, providing data across a variety of categories, including activities and uses of the chemical, releases and other waste management ( <i>e.g.</i> , quantities recycled, treated, combusted) and pollution prevention activities (under Section 6607 of the Pollution Prevention Act). These data include on- and off-site data as well as multimedia data ( <i>i.e.</i> , air, land and water)	1,2-Dichloropropane is a listed substance subject to reporting requirements under <u>40 CFR</u> <u>372.65</u> effective as of January 01, 1987.
Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA) - Sections 3 and 6	FIFRA governs the sale, distribution and use of pesticides. Section 3 of FIFRA generally requires that pesticide products be registered by EPA prior to distribution or sale. Pesticides may only be registered if,	<ul><li>1,2-Dichloropropane was registered as a conventional chemical in April 23, 1984.</li><li>EPA removed 1,2- dichloropropane from its list of</li></ul>

Statutes/Regulations	Description of Authority/Regulation	Description of Regulation
	among other things, they do not cause " <i>unreasonable adverse effects on the</i> <i>environment</i> ." Section 6 of FIFRA provides EPA with the authority to cancel pesticide registrations if either (1) the pesticide, labeling, or other material does not comply with FIFRA; or (2) when used in accordance with widespread and commonly recognized practice, the pesticide generally causes unreasonable adverse effects on the environment.	pesticide product inert ingredients used in pesticide products on June 24, 1998, it is allowed as an impurity only. ( <u>Pesticides Chemical Search,</u> <u>OPP, Accessed April 16, 2019)</u>
Clean Air Act (CAA) – Section 111(b)	Requires EPA to establish new source performance standards (NSPS) for any category of new or modified stationary sources that EPA determines causes, or contributes significantly to, air pollution, which may reasonably be anticipated to endanger public health or welfare. The standards are based on the degree of emission limitation achievable through the application of the best system of emission reduction (BSER) which (taking into account the cost of achieving reductions and environmental impacts and energy requirements) EPA determines has been adequately demonstrated.	1,2-Dichloropropane is subject to the NSPS for equipment leaks of volatile organic compounds (VOCs) in the synthetic organic chemicals manufacturing industry for which construction, reconstruction or modification began after January 5, 1981 (40 CFR Part 60, Subpart VV).
Clean Air Act (CAA) – Section 112(b)	Defines the original list of 189 hazardous air pollutants (HAPs). Under 112(c) of the CAA, EPA must identify and list source categories that emit HAP and then set emission standards for those listed source categories under CAA Section 112(d). CAA Section 112(b)(3)(A) specifies that any person may petition the Administrator to modify the list of HAP by adding or deleting a substance. Since 1990, EPA has removed two pollutants from the original list leaving 187 at present.	1,2-Dichloropropane is listed as a HAP. ( <u>42 U.S. Code Section 7412</u> )

Statutes/Regulations	Description of Authority/Regulation	Description of Regulation
Clean Air Act (CAA) – Section 112(d)	Directs EPA to establish, by rule, NESHAPs for each category or subcategory of listed major sources and area sources of HAPs (listed pursuant to Section 112(c)). For major sources, the standards must require the maximum degree of emission reduction that EPA determines is achievable by each particular source category. This is generally referred to as maximum achievable control technology (MACT). For area sources, the standards must require generally achievable control technology (GACT) though may require MACT.	EPA has established <u>NESHAP</u> for a number of source categories that emit 1,2- dichloropropane to air.
Clean Air Act (CAA) – Section 112(d) and 112(f)	Risk and technology review (RTR) of Section 112(d) national emission standards for hazardous air pollutants (NESHAP). Section 112(f)(2) requires EPA to conduct risk assessments for each source category subject to Section 112(d) NESHAP that require maximum achievable control technology (MACT), and to determine if additional standards are needed to reduce remaining risks. Section 112(d)(6) requires EPA to review and revise the emission standards, as necessary, taking into account developments in practices, processes and control technologies.	EPA has promulgated a number of RTR <u>NESHAP</u> and will do so, as required, for the remaining source categories with NESHAP.
Clean Water Act (CWA) - Section 304(a)(1)	Requires EPA to develop and publish ambient water quality criteria (AWQC) reflecting the latest scientific knowledge on the effects on human health that may be expected from the presence of pollutants in any body of water.	In 2015, EPA published updated <u>AWQC</u> for 1,2- dichloropropane, including a recommendation of 0.90 ( $\mu$ g/L) for "Human Health for the consumption of Water + Organism" and 31 ( $\mu$ g/L) for "Human Health for the consumption of Organism Only" for states and authorized tribes to consider when adopting criteria into their water quality standards.

Statutes/Regulations	Description of Authority/Regulation	Description of Regulation
Clean Water Act (CWA) – Section 301, 304, 306, 307, and 402	Clean Water Act Section 307(a) establishes a list of toxic pollutants or combination of pollutants under the CWA. The statute specifies a list of families of toxic pollutants also listed in the Code of Federal Regulations at 40 CFR Part 401.15. The "priority pollutants" specified by those families are listed in 40 CFR Part 423 Appendix A. These are pollutants for which best available technology effluent limitations must be established on either a national basis through rules (Sections 301(b), 304(b), 307(b), 306) or on a case-by-case best professional judgement basis in NPDES permits, see Section 402(a)(1)(B). EPA identifies the best available technology that is economically achievable for that industry after considering statutorily prescribed factors and sets regulatory requirements based on the performance of that technology.	1,2-Dichloropropane is designated as a priority pollutant under Section 307(a)(1) of the CWA and as such is subject to effluent limitations. Under CWA Section 304, 1,2- dichloropropane is included in the list of total toxic organics (TTO) (40 CFR 413.02(i)).
Clean Water Act (CWA) – Section 311(b)(2)(A) and 501(a) of the Federal Water Pollution Control Act.	Requires EPA to develop, promulgate, and revise as may be appropriate, regulations designating as hazardous substances, other than oil, which, when discharged present an imminent and substantial danger to the public health or welfare, including, but not limited to, fish, shellfish, wildlife, shorelines, and beaches.	1,2-Dichloropropane is a designated hazardous substance in accordance with Section 311(b) (2)(A) of the Federal Water Pollution Control Act ( <u>40</u> <u>FR 116.4</u> , March 13, 1978).
Safe Drinking Water Act (SDWA) – Section 1412	Requires EPA to publish a non- enforceable maximum contaminant level goal (MCLG) for a contaminant for which EPA makes the determination that the contaminant: 1. may have an adverse effect on the health of persons; 2. is known to occur or there is a substantial likelihood that the contaminant will occur in public water systems with a frequency and at levels of public health concern; and 3. in the sole judgement of the	1,2-Dichloropropane is subject to NPDWR under the SDWA with a MCLG of zero mg/L and an enforceable MCL of 0.005 mg/L (Section 1412) (52 FR 25690, January 30, 1991).

Statutes/Regulations	Description of Authority/Regulation	Description of Regulation
	Administrator, regulation of the contaminant presents a meaningful opportunity for health risk reductions for persons served by public water systems. When EPA publishes an MCLG, EPA must also promulgate a National Primary Drinking Water Regulation (NPDWR) which includes either an enforceable maximum contaminant level (MCL), or a required treatment technique. Public water systems are required to comply with NPDWRs.	
Clean Water Act (CWA) – Section 311(b)(2)(A) and 501(a) of the Federal Water Pollution Control Act.	Requires EPA to develop, promulgate, and revise as may be appropriate, regulations designating as hazardous substances, other than oil, which, when discharged present an imminent and substantial danger to the public health or welfare, including, but not limited to, fish, shellfish, wildlife, shorelines, and beaches.	1,2-Dichloropropane is a designated hazardous substance in accordance with Section 311(b)(2)(A) of the Federal Water Pollution Control Act (40 FR 116.4, March 13, 1978).
Resource Conservation and Recovery Act (RCRA) – Section 3001	Directs EPA to develop and promulgate criteria for identifying the characteristics of hazardous waste, and for listing hazardous waste, taking into account toxicity, persistence, and degradability in nature, potential for accumulation in tissue and other related factors such as flammability, corrosiveness, and other hazardous characteristics.	1,2-Dichloropropane is included on the list of hazardous wastes pursuant to RCRA 3001. RCRA Hazardous Waste Code: U083 ( <u>40 CFR 261.33</u> ).
Comprehensive Environmental Response, Compensation and Liability Act (CERCLA) – Sections 102(a) and 103	Authorizes EPA to promulgate regulations designating as hazardous substances those substances which, when released into the environment, may present substantial danger to the public health or welfare or the environment. EPA must also promulgate regulations establishing the quantity of any hazardous substance the release of	1,2-Dichloropropane is a hazardous substance under CERCLA. Releases of 1,2- dichloropropane in excess of 1000 pounds must be reported ( <u>40 CFR 302.4</u> ).

Statutes/Regulations	Description of Authority/Regulation	Description of Regulation
	which must be reported under Section 103. Section 103 requires persons in charge of vessels or facilities to report to the National Response Center if they have knowledge of a release of a hazardous substance above the reportable quantity threshold.	
Superfund Amendments and Reauthorization Act (SARA) –	Requires the Agency to revise the hazardous ranking system and update the National Priorities List of hazardous waste sites, increases state and citizen involvement in the superfund program and provides new enforcement authorities and settlement tools.	1,2-Dichloropropane is listed on SARA, an amendment to CERCLA and the CERCLA Priority List of Hazardous Substances. This list includes substances most commonly found at facilities on the CERCLA National Priorities List (NPL) that have been deemed to pose the greatest threat to public health.
Other Federal Statutes/Reg	gulations	
Occupational Safety and Health Act (OSHA)	Requires employers to provide their workers with a place of employment free from recognized hazards to safety and health, such as exposure to toxic chemicals, excessive noise levels, mechanical dangers, heat or cold stress or unsanitary conditions (29 U.S.C Section 651 et seq.). Under the Act, OSHA can issue occupational safety and health standards including such provisions as Permissible Exposure Limits (PELs), exposure monitoring, engineering and administrative control measures, and respiratory protection.	On June 30, 1993, OSHA issued occupational safety and health standards for 1,2- dichloropropane that included a PEL of 75 ppm TWA, exposure monitoring, control measures and respiratory protection (29 <u>CFR 1910.1000</u> ). OSHA Annotated Table Z-1, Accessed April 16, 2019.
Federal Hazardous Materials Transportation Act (HMTA)	<ul> <li>Section 5103 of the Act directs the Secretary of Transportation to:</li> <li>Designate material (including an explosive, radioactive material, infectious substance, flammable or combustible liquid, solid or gas, toxic, oxidizing or corrosive material, and compressed gas) as</li> </ul>	1,2-Dichloropropane is listed as a hazardous material with regard to transportation and is subject to regulations prescribing requirements applicable to the shipment and transportation of listed

Statutes/Regulations	<b>Description of Authority/Regulation</b>	Description of Regulation
	<ul> <li>hazardous when the Secretary determines that transporting the material in commerce may pose an unreasonable risk to health and safety or property.</li> <li>Issue regulations for the safe transportation, including security, of hazardous material in intrastate, interstate and foreign commerce.</li> </ul>	hazardous materials ( <u>70 FR</u> <u>34381</u> , June 14, 2005).

# **D.2** State Laws and Regulations

State Actions	Description of Action	
State Air Regulations	Allowable Ambient Levels New Hampshire 232 24-Hr AAL (µg/m3)	
	4.0 Annual AALB (µg/m3) (Env-A 1400: Regulated Toxic Air	
	Pollutants). Rhode Island 0.1 Annual (µg/m3) (Air Pollution Regulation	
	<u>No. 22</u> ).	
State Drinking Water	Arizona ( <u>14 Ariz. Admin. Register 2978</u> , August 1, 2008), California	
Standards and Guidelines	(Cal Code Regs. Title 26, § 22-64444), Delaware (Del. Admin. Code	
	<u>Title 16, § 4462</u> ), Connecticut (Conn. Agencies Regs. <u>§ 19-13-B102</u> ),	
	Florida (Admin. Code R. Chap. 62-550), Maine (10 144 Me. Code R.	
	Chap. 231), Massachusetts (310 Code Mass. Regs. § 22.00), Michigan	
	(Mich. Admin. <u>Code r.299.44 and r.299.49</u> , 2017), Minnesota (Minn R.	
	Chap. 4720) (MN specifies VOC's, but does not go beyond that), New	
	Jersey (7:10 N.J Admin. Code § 5.2), Pennsylvania (25 Pa. Code §	
	<u>109.202</u> ), Rhode Island (Rules and Regulations Pertaining to Public	
	Drinking Water <u>R46-13-DWQ</u> ), Texas ( <u>30 Tex. Admin. Code §</u>	
	<u>290.104</u> ).	
State PELs	California (PEL of 75 ppm and a STEL of 110) (Cal Code Regs. <u>Title 8</u> ,	
	<u>§ 5155</u> ).	
	Hawaii PEL: 75 ppm ( <u>Hawaii Administrative Rules Section 12-60-50</u> ).	
State Right-to-Know	Massachusetts (105 Code Mass. Regs. § 670.000 Appendix A) and New	
Acts	Jersey (8:59 N.J. Admin. Code § 9.1).	
Chemicals of High	Several states have adopted reporting laws for chemicals in children's	
Concern to Children	products containing 1,2-dichloropropane, including Maine's list of	
	Chemical of Concern ( <u>38 MRSA Chapter 16-D</u> ).	
Other	California listed 1,2-dichloropropane on Proposition 65 in January 1,	
	1990 due to cancer ( <u>Cal Code Regs. Title 27, § 27001</u> ).	
	1,2-Dichloropropane is listed as a Candidate Chemical under	
	California's Safer Consumer Products Program established under Health	

State Actions	Description of Action	
	and Safety Code § 25252 and 25253. (California, Candidate Chemicals	
	List. Accessed April 18, 2019) (CDTSC, 2017).	
	California issued a Health Hazard Alert for 1,2-dichloropropane (Hazard Evaluation System and Information Service, 2016).	
	California lists 1,2-dichloropropane as a designated priority chemical for biomonitoring under criteria established by <u>California SB 1379</u> (CDPH, 2015) (Accessed February 2019).	
	1,2-Dichloropropane is on the MA Toxic Use Reduction Act (TURA) list of 1989 ( <u>301 Code Mass. Regs. § 41.03</u> ).	

## **D.3** International Laws and Regulations

<b>Country/Organization</b>	Requirements and Restrictions	
Canada	Canada requires Reportable to National Pollutant Release Inventory for 1,2-dichloropropane if manufactured, processed, or otherwise used at quantities greater than: 10 tonnes. NPRI Part (Threshold Category): <u>1A</u> , <u>2018-01-20</u> . Canada added 1,2-dichloropropane to the Domestic Substances List as an "Existing Substance", <i>i.e.</i> , already in commerce in Canada in 1984-1986 and not subject to the New Substance Notification Regulations. <u>Canada</u> <u>Gazette Part II, Vol. 128, No. 9</u> , May 04 1994, SOR/94-311. 1,2-Dichloropropane did not meet Canada's criteria under Subsection 73(1) of the Canadian Environmental Protection Act, 1999 ( <u>CEPA</u> ). Canadian Environmental Protection Act: categorization of existing	
	substances. 2006-09-14.	
European Union	1,2-Dichloropropane is registered for use in the EU. (European Chemicals Agency (ECHA) database. Accessed April 17, 2019.)	
Australia	<ul> <li>1,2-Dichloropropane can be manufactured or imported into Australia for commercial purposes without notifying us, provided that the Australian importer/manufacturer is currently registered with us. This chemical may be subject to other Australian Government and State or Territory Government regulations.</li> <li>1,2-Dichloropropane was assessed under Human Health Tier II of the Inventory Multi-Tiered Assessment and Prioritisation (IMAP).</li> <li>No specific Australian use, import, or manufacturing information has been identified.</li> </ul>	
	The following international uses have been identified through Galleria Chemica, European Union Registration, Evaluation, Authorisation and Restriction of Chemicals dossier (REACH), Substances and Preparations in the Nordic countries (SPIN) database, United States (US) Department of Health National Toxicology Program (NTP), US Environmental Protection Agency Chemical and Product Categories (CPCat), Hazardous Substances Data Bank (HSDB), Organisation for Economic Co-operation and Development (OECD, 2003), International Agency for Research on Cancer (IARC) Agency for Toxic Substances and Disease Registry <u>ATSDR (1989)</u> and the National Institute of standards and Technology (NIST).	

 Table\_Apx D-3. Regulatory Actions by other Governments, Tribes, and International Agreements

 Country/Organization
 Requirements and Restrictions

<b>Country/Organization</b>	Requirements and Restrictions
	The chemical may have domestic uses including: as a solvent for glues, adhesives, degreasers, cleaning and painting products; as a stain remover for fabrics; in car care products; and as a paint remover. ( <u>NICNAS</u> , 2017, <i>Propane, 1,2-dichloro-: Human health tier II</i>
Japan	<ul> <li>assessment, Accessed April 17, 2019.)</li> <li>1,2-Dichloropropane is regulated in Japan under the following legislation: <ul> <li>Act on the Evaluation of Chemical Substances and Regulation of Their Manufacture, <i>etc.</i> (Chemical Substances Control Law; CSCL)</li> <li>Act on Confirmation, <i>etc.</i> of Release Amounts of Specific Chemical Substances in the Environment and Promotion of Improvements to the Management Thereof</li> <li>Industrial Safety and Health Act (ISHA)</li> <li>Air Pollution Control Law</li> <li>Water Pollution Control Law</li> <li>(National Institute of Technology and Evaluation [NITE] Chemical Risk Information Platform [CHRIP], Accessed April 17, 2019).</li> </ul> </li> </ul>
World Health Organization (WHO)	EHC on 1,2-dichloropropane found that for the general population there is low or non-existent exposure to 1,2-dichloropropane from air and food. However, in certain areas, exposure may occur when groundwater is contaminated. Occupational exposure: With good work practices, hygienic measures, and safety precautions, the use of 1,2-dichloropropane is unlikely to present a risk for those occupationally exposed to it. (EHC <u>146</u> , 1993).
Australia, Belgium, Canada, Denmark, Finland, France, Hungary, Ireland, Japan, New Zealand, People's Republic of China, Poland, Romania, Singapore, South Korea, Spain, Switzerland	Occupational exposure limits for 1,2-dichloropropane (GESTIS International limit values for chemical agents (Occupational exposure limits, OELs) database, Accessed April 18, 2019.

# Appendix E PROCESS, RELEASE AND OCCUPATIONAL EXPOSURE INFORMATION

This appendix provides information and data found in preliminary data identification and evaluation for 1,2-dichloropropane.

### **E.1 Process Information**

Process-related information potentially relevant to the risk evaluation may include process diagrams, descriptions and equipment. Such information may inform potential release sources and worker exposure activities.

#### E.1.1 Manufacture (Including Import)

#### E.1.1.1 Manufacture

A commenter informed EPA that methods to manufacture 1,2-dichloropropane include a chemical reaction of propylene chloride and antimony pentachloride. According to the same commenter, it is also a byproduct of the reaction of propene and chlorine in epichlorohydrin manufacturing (EPA-HQ-OPPT-2018-0428-0011). EPA plans to further evaluate the method(s) of domestic manufacture of 1,2-dichloropropane during the risk evaluation.

#### E.1.1.2 Import

Commodity chemicals such as 1,2-dichloropropane may be imported into the United States in bulk via water, air, land, and intermodal shipments (Tomer and Kane, 2015). These shipments take the form of oceangoing chemical tankers, railcars, tank trucks, and intermodal tank containers. Chemicals shipped in bulk containers may be repackaged into smaller containers for resale, such as drums or bottles. Domestically manufactured commodity chemicals may be shipped within the United States in liquid cargo barges, railcars, tank trucks, tank containers, intermediate bulk containers (IBCs)/totes, and drums. Both imported and domestically manufactured commodity chemicals may be repackaged by wholesalers for resale; for example, repackaging bulk packaging into drums or bottles. The type and size of container will vary depending on customer requirement. In some cases, QC samples may be taken at import and repackaging sites for analyses. Some import facilities may only serve as storage and distribution locations, and repackaging/sampling may not occur at all import facilities.

In the 2016 CDR, companies reported importing a formulation containing at least 90 percent 1,2-dichloropropane (U.S. EPA, 2016a).

#### E.1.2 Processing and Distribution

#### E.1.2.1 Processing as a Reactant

Processing as a reactant or intermediate is the use of 1,2-dichloropropane as a feedstock in the production of another chemical via a chemical reaction in which 1,2-dichloropropane is consumed to form the product. In the 2016 CDR, companies reported use of 1,2-dichloropropane as an intermediate in the manufacture of other basic organic chemicals (U.S. EPA, 2016a). EPA has not identified specific process information for the processing of 1,2-dichloropropane as a reactant.

#### E.1.2.2 Incorporation into Formulation, Mixture, or Reaction Product

Incorporation into a formulation, mixture or reaction product refers to the process of mixing or blending of several raw materials to obtain a product or mixture (*e.g.*, cleaning products). EPA has not identified formulation processes specific to 1,2-dichloropropane.

#### E.1.3 Uses

#### E.1.3.1 Industrial Use – Non-Incorporative Activities

1,2-Dichloropropane is used at Solenis LLC's Courtland, Virginia plant as a solvent in the production of alkaline paper surface sizing agents. At the end of this batch process, 1,2-dichloropropane is evaporated under vacuum and recovered (EPA-HQ-OPPT-2018-0428-0015).

#### E.1.3.2 Commercial Laboratory Use

A SDS for 1,2-dichloropropane (100 percent purity) states the chemical has laboratory use. However, specific laboratory use activities are unknown (<u>Chem Service, 2018</u>). A commenter (<u>EPA-HQ-OPPT-2018-0428-0031</u>) provided descriptions of their use of 1,2-dichloropropane in analytical standard, research, equipment calibration and sample preparation applications, including reference sample for analysis of terrestrial and extraterrestrial material samples, which the commenter also indicated was a critical use, further informing EPA's understanding of this condition of use.

#### E.1.3.3 Commercial Cleaning and Furnishing Care Products

1,2-Dichloropropane is present in various liquid spray cleaners, all-purpose liquid cleaners, waxes and polishes at concentrations ranging from 1 to 54 percent concentration. According to product webpages, these products are first applied to surfaces via spray, sponge, or cloth, then wiped or buffed off (<u>Tenax</u> <u>USA</u>, 2020a, b, c, 2015).

#### E.1.4 Disposal

Each of the conditions of use of 1,2-dichloropropane may generate waste streams of the chemical that are collected and transported to third-party sites for disposal, treatment, or recycling. Industrial sites that treat or dispose onsite wastes that they themselves generate will be assessed in each condition of use. Similarly, point source discharges of 1,2-dichloropropane to surface water will be assessed in each condition of use. Wastes of 1,2-dichloropropane that are generated during a condition of use and sent to a third-party site for treatment or disposal may include wastewater and solid waste. 1,2-Dichloropropane may be contained in wastewater discharged to POTW or other, non-public treatment works for treatment. Wastewater discharges to POTWs and non-public treatment works of 1,2-dichloropropane will be evaluated during risk evaluation.

Solid wastes are defined under RCRA as any material that is discarded by being: abandoned; inherently waste-like; a discarded military munition; or recycled in certain ways (certain instances of the generation and legitimate reclamation of secondary materials are exempted as solid wastes under RCRA).

## E.2 Preliminary Occupational Exposure Data

EPA presents below examples of occupational exposure-related information from the preliminary data identification and evaluation. EPA plans to consider this information and data in combination of other data and methods for use in the risk evaluation. Note there are no OSHA Chemical Exposure and Health Data (CEHD) for 1,2-dichloropropane within the last ten years. In addition, there are no NIOSH Health Hazard Evaluations specific to this chemical.

Table\_Apx E-1. Summary of Industry Sectors with 1,2-Dichloropropane Personal Monitoring Air Samples Obtained from OSHA Inspections Conducted since 1984

NAICS Code	NAICS Description	Number of Data Points					
327991	Cut Stone and Stone Product Manufacturing	2					
Jumber of data points in Table Apx E-1 was populated from data found at							

Number of data points in Table\_Apx E-1was populated from data found at https://www.osha.gov/opengov/healthsamples.html

# Appendix F SUPPORTING INFORMATION - CONCEPTUAL MODEL FOR INDUSTRIAL AND COMMERCIAL ACTIVITIES AND USES

Life Cycle Stage	Category	Subcategory	Release / Exposure Scenario	Exposure Pathway	Exposure Route	Receptor / Population	Plans to Evaluate	Rationale
			Manufacture of 1,2- dichloropropane	Liquid Contact	Dermal	Worker	Yes	Workers are expected to routinely handle liquids containing 1,2-Dichloropropane.
	Domestic	Domestic		Vapor	Inhalation	Worker	Yes	Due to high volatility (53.3 mmHg at room temperature), EPA plans to evaluate inhalation exposure to vapor.
Manufacturing	manufacturing Import	manufacturing Import		Liquid Contact	Dermal	ONU	No	Dermal exposure by ONU is not expected for this condition of use as they are not expected to directly handle the chemical.
				Vapor	Inhalation	ONU	Yes	Due to high volatility (53.3 mmHg at room temperature), EPA plans to evaluate inhalation exposure to vapor.
	As a reactant	Intermediate in all other basic organic chemical manufacturing	Manufacture of all other basic organic chemicals	Liquid Contact	Dermal	Worker	Yes	Workers are expected to routinely handle liquids containing 1,2-Dichloropropane.
				Vapor	Inhalation	Worker	Yes	Due to high volatility (53.3 mmHg at room temperature), EPA plans to evaluate inhalation exposure to vapor.
Processing				Liquid Contact	Dermal	ONU	No	Dermal exposure by ONU is not expected for this condition of use as they are not expected to directly handle the chemical.
				Vapor	Inhalation	ONU	Yes	Due to high volatility (53.3 mmHg at room temperature), EPA plans to evaluate inhalation exposure to vapor.

 Table\_Apx F-1. Worker and Occupational Non-User Exposure Conceptual Model Supporting Table

Life Cycle Stage	Category	Subcategory	Release / Exposure Scenario	Exposure Pathway	Exposure Route	Receptor / Population	Plans to Evaluate	Rationale
				Liquid Contact	Dermal	Worker	Yes	Workers are expected to routinely handle liquids containing 1,2-Dichloropropane.
	Incorporation into	Intermediate in all other chemical	Manufacture of all other	Vapor	Inhalation	Worker	Yes	Due to high volatility (53.3 mmHg at room temperature), EPA plans to evaluate inhalation exposure to vapor.
Processing	formulation, mixture, or reaction product	other chemical product and preparation manufacturing	all other chemical product and preparation	Liquid Contact	Dermal	ONU	No	Dermal exposure by ONU is not expected for this condition of use as they are not expected to directly handle the chemical.
				Vapor	Inhalation	ONU	Yes	Due to high volatility (53.3 mmHg at room temperature), EPA plans to evaluate inhalation exposure to vapor.
Distribution in Commerce	Distribution in Commerce	Distribution in Commerce	Distribution of bulk shipments of 1,2- dichloropropane and formulated products	Liquid Contact, Vapor	Dermal, Inhalation	Worker, ONU	Yes	EPA plans to evaluate activities resulting in exposures associated with distribution in commerce ( <i>e.g.</i> , loading, unloading) throughout the various lifecycle stages and conditions of use ( <i>e.g.</i> , manufacturing, processing, industrial use, commercial use, disposal) rather than as a single distribution scenario.
			Processing solvent used in the production of paper sizing agent	Liquid Contact	Dermal	Worker	Yes	Workers are expected to routinely handle liquids containing 1,2-Dichloropropane.
	Non-	Processing aids, not otherwise listed		Vapor	Inhalation	Worker	Yes	Due to high volatility (53.3 mmHg at room temperature), EPA plans to evaluate inhalation exposure to vapor.
Industrial Use	incorporative activities			Liquid Contact	Dermal	ONU	No	Dermal exposure by ONU is not expected for this condition of use as they are not expected to directly handle the chemical.
				Vapor	Inhalation	ONU	Yes	Due to high volatility (53.3 mmHg at room temperature), EPA plans to evaluate inhalation exposure to vapor.

Life Cycle Stage	Category	Subcategory	Release / Exposure Scenario	Exposure Pathway	Exposure Route	Receptor / Population	Plans to Evaluate	Rationale
			Laboratory chemical use	Liquid Contact	Dermal	Worker	Yes	Workers are expected to routinely handle liquids containing 1,2-Dichloropropane.
Commercial		Laboratoria		Vapor	Inhalation	Worker	Yes	Due to high volatility (53.3 mmHg at room temperature), EPA plans to evaluate inhalation exposure to vapor.
use	Other use	Laboratory chemicals		Liquid Contact	Dermal	ONU	No	Dermal exposure by ONU is not expected for this condition of use as they are not expected to directly handle the chemical.
				Vapor	Inhalation	ONU	Yes	Due to high volatility (53.3 mmHg at room temperature), EPA plans to evaluate inhalation exposure to vapor.
	Cleaning and furnishing care products	All-purpose liquid spray cleaner All-purpose waxes and polishes All-purpose liquid cleaner/polish	Application of liquid cleaners, polishes, and waxes	Liquid Contact	Dermal	Worker	Yes	Workers are expected to routinely handle liquids (including from mist) containing 1,2-Dichloropropane.
				Vapor	Inhalation	Worker	Yes	Due to high volatility (53.3 mmHg at room temperature), EPA plans to evaluate inhalation exposure to vapor.
Commercial use				Mist	Inhalation	Worker, ONU	Yes	Based on information from SDS, product website, and third-party site, some polishing products may be spray-applied.
				Liquid Contact	Dermal	ONU	No	Dermal exposure by ONU is not expected for this condition of use as they are not expected to directly handle the chemical.
				Vapor	Inhalation	ONU	Yes	Due to high volatility (53.3 mmHg at room temperature), EPA plans to evaluate inhalation exposure to vapor.

Life Cycle Stage	Category	Subcategory	Release / Exposure Scenario	Exposure Pathway	Exposure Route	Receptor / Population	Plans to Evaluate	Rationale
		Disposal 1	Worker handling of wastes	Liquid Contact	Dermal	Worker	Yes	Workers are expected to routinely handle liquids containing 1,2-Dichloropropane.
				Vapor	Inhalation	Worker	Yes	Due to high volatility (53.3 mmHg at room temperature), EPA plans to evaluate inhalation exposure to vapor.
Disposal	Disposal			Liquid Contact	Dermal	ONU	No	Dermal exposure by ONU is not expected for this condition of use as they are not expected to directly handle the chemical.
				Vapor	Inhalation	ONU	Yes	Due to high volatility (53.3 mmHg at room temperature), EPA plans to evaluate inhalation exposure to vapor.

## Appendix G SUPPORTING INFORMATION - CONCEPTUAL MODEL FOR CONSUMER ACTIVITIES AND USES

Life Cycle Stage	Category	Subcategory	Release from source	Exposure Pathway	Route	Receptor	Plan to Evaluate	Rationale
		All-purpose liquid spray cleaner	Application of liquid waxes and polishes	Liquid Contact	Dermal	Consumer	Yes	Consumers may potentially handle liquids containing 1,2-dichloropropane.
Consumer Use	Cleaning and furniture care products	All-purpose waxes and polishes All-purpose liquid cleaner/polish	Application of spray cleaner or polish	Vapor/Mist	Inhalation	Consumers/ Bystanders	Yes	Due to high volatility (53.3 mmHg at room temperature), EPA plans to evaluate inhalation exposure to vapor. Based on information from SDS, product website, and third-party site, some polishing products may be spray-applied.

#### Table\_Apx G-1. Consumer Exposure Conceptual Model Supporting Table

# Appendix HSUPPORTING INFORMATION – CONCEPTUAL MODEL FOR<br/>ENVIRONMENTAL RELEASES AND WASTES

Life Cycle Stage	Category	Release	Exposure Pathway / Media	Exposure Routes	Receptor / Population	Plans to Evaluate	Rationale
			Near facility ambient air concentrations	Inhalation	General Population	No	1,2-Dichloropropane is a HAP. Stationary source
	Emissions to Air	Emissions to Air	Indirect deposition to	Oral Dermal	General Population	No	releases of 1,2- dichloropropane to ambient
			nearby bodies of water and soil catchments	TBD	Aquatic and Terrestrial Receptors	No	air are under the jurisdiction of the CAA
	Wastewater or Liquid Wastes	Industrial pre- treatment and wastewater treatment, or POTW	Direct release into surface water and	TBD	Aquatic Receptors	Yes	EPA has developed Ambient Water Quality Criteria for
All			indirect partitioning to sediment	Oral Dermal	General Population	No	protection of human health for 1,2-dichloropropane.
			Drinking Water via Surface or Ground Water	Oral Dermal and Inhalation ( <i>e.g.</i> , showering)	General Population	No	The drinking water exposure pathway for 1,2- dichloropropane is currently addressed in NPDWR.
			Biosolids: application to soil and/or migration to groundwater and/or	Oral ( <i>e.g.</i> , ingestion of soil) Inhalation	General Population	No	Unlikely to be a route to general population.
			surface water	TBD	Aquatic receptors	Yes	
		Underground injection	Migration to groundwater, potential surface/drinking water	Oral Dermal Inhalation	General Population	No	1,2-Dichloropropane is released to Class I Underground Injection Wells

#### Table\_Apx H-1. General Population and Environmental Exposure Conceptual Model Supporting Table

Life Cycle Stage	Category	Release	Exposure Pathway / Media	Exposure Routes	Receptor / Population	Plans to Evaluate	Rationale
				TBD	Aquatic and Terrestrial Species		which are covered by SDWA and RCRA.
	Disposal Solid and Liquid Wastes Hazardous, Municipal landfill and other land disposal	Hazardous,	Leachate to soil,	Oral Dermal	General Population	No	1,2-Dichloropropane is included on the list of hazardous wastes pursuant to
Disposal		ground water and/or mitigation to surface water	TBD	Aquatic and Terrestrial Receptors		RCRA 3001 (40 CFR §§ 261.33).	