Development of Emissions Estimating Methodologies for Egg-laying Houses and Manure Sheds

Draft

Prepared by:

U.S. Environmental Protection Agency

Office of Air and Radiation
Office of Air Quality Planning and Standards
109 T.W. Alexander Drive
Research Triangle Park, N.C. 27709

August 2021

This document is a preliminary draft. It has not been formally released by the U.S. Environmental Protection Agency (EPA) and should not at this stage be construed to represent Agency policy. It is being circulated for comments on its technical merit.

Table of Contents

1.0	INTE	RODUCT	ΓΙΟΝ	1-1
	1.1	Site de	escriptions	1-1
		1.1.1	CA2B	1-1
		1.1.2	IN2B/IN2H	1-2
		1.1.3	NC2B	1-3
	1.2	Data S	Sampled	1-4
		1.2.1	Animal Husbandry	1-5
		1.2.2	Biomaterials Sampling Methods and Schedule	1-5
2.0	REV	ISIONS	TO DATASET AND EMISSIONS DATA SUMMARY	2-1
	2.1	Revisi	ions to the 2010 Dataset	2-1
	2.2	Data (Completeness Criteria for the Revised Dataset	2-2
	2.3	Comp	arison between the 2010 and Revised Datasets	2-3
		2.3.1	NH ₃ High Rise Dataset	2-3
		2.3.2	H ₂ S High Rise Dataset	2-3
		2.3.3	PM High Rise Dataset	2-4
		2.3.4	NH ₃ Manure Belt House Dataset	2-5
		2.3.5	H ₂ S Manure belt Dataset	2-6
		2.3.6	PM Manure Belt House Dataset	2-6
		2.3.7	Manure Shed Dataset	2-7
	2.4		arison Between the Revised Datasets and NAEMS Datasets Used ved Published Papers	
		2.4.1	High Rise Houses	2-7
		2.4.2	Manure belt Houses	2-12
		2.4.3	Manure Shed	2-13
3.0	REL	ATIONS	SHIPS ESTABLISHED IN LITERATURE	3-1
	3.1	NH ₃ a	nd H ₂ S Emissions from Houses and Manure Sheds	3-1
	3.2	PM E1	missions from House and Manure Sheds	3-4
4.0	SITE	COMP	ARISON, TRENDS, AND ANALYSIS	4-1
	4.1	Emiss	ions Data	4-2
		4.1.1	High Rise Houses	4-2
		4.1.2	Manure belt Houses	4-3
		4.1.3	Manure Shed	4-5

	4.2	Enviro	onmental Parameters	4-5
		4.2.1	High Rise Houses	4-5
		4.2.2	Manure Belt Houses	4-9
		4.2.3	Manure Shed	4-12
	4.3	Manur	re Parameters	4-14
		4.3.1	High Rise Houses	4-14
		4.3.2	Manure belt Houses	4-16
		4.3.3	Manure Shed	4-17
5.0	DEV	ELOPM	ENT AND SELECTION OF MODELS FOR DAILY EMISSIONS	5-1
	5.1	High F	Rise Operations	5-1
	5.2	Manur	e Belt Operations	5-4
	5.3	Manur	e Sheds	5-8
6.0	MOD	EL COI	EFFICIENT EVALUATION	6-1
	6.1	High F	Rise Layer House Models	6-1
		6.1.1	NH ₃ Model Evaluation	6-1
		6.1.2	H ₂ S Model Evaluation	6-3
		6.1.3	PM ₁₀ Model Evaluation	6-5
		6.1.4	PM _{2.5} Model Evaluation	6-7
		6.1.5	TSP Model Evaluation	6-9
	6.2	Manur	e belt Layer House Models	6-11
		6.2.1	NH ₃ Model Evaluation	6-11
		6.2.2	H ₂ S Model Evaluation	6-13
		6.2.3	PM ₁₀ Model Evaluation	6-14
		6.2.4	PM _{2.5} Model Evaluation	6-15
		6.2.5	TSP Model Evaluation	6-16
	6.3	Manur	e Shed Models	6-17
7.0	ANN	UAL EM	IISSION ESTIMATES AND MODEL UNCERTAINTY	7-1
8.0	MOD	EL APP	PLICATION AND ADDITIONAL TESTING	8-1
	8.1	Model	Application Example	8-1
		8.1.1	High Rise Example	8-2
		8.1.2	Manure Belt House Example	8-3
		8.1.3	Manure Shed Example	8-3
		8.1.4	Combining Structures	8-4
	8.2	Model	Sensitivity Testing	8-5
		8.2.1	Sensitivity to Inventory	8-5

10.0	REFE	RENCE	ES	10-1
9.0	SUMN	MARY A	AND CONCLUSIONS	9-1
		8.5.3	Manure Shed	8-30
		8.5.2	Manure belt	8-28
		8.5.1	High Rise	8-27
	8.5	Data C	Concerns	8-27
	8.4	Replic	cation of Independent Measurements	8-23
		8.3.3	Manure Shed	8-23
		8.3.2	Manure belt House	8-22
		8.3.1	High Rise House	8-21
	8.3	Compa	arison to literature	8-21
		8.2.3	Model Limitations	8-11
		8.2.2	Sensitivity to Climate	8-6

List of Tables

Table 1-1. Layer Confinement Sites Monitored Under NAEMS	-1
Table 2-1. Number of valid ADM and mean NH ₃ emission values (at 75% data completenes	ss)
between the 2010 and revised high rise datasets.	-3
Table 2-2. Number of valid ADM and mean H ₂ S emission values (at 75% data completenes	
between the 2010 and revised high rise datasets.	-4
Table 2-3. Number of valid ADM and mean PM ₁₀ emission values between the 2010 and revise	ed
high rise site datasets.	-4
Table 2-4. Number of valid ADM and mean PM _{2.5} emission values between the 2010 and revise	ed
high rise site datasets.	-5
high rise site datasets	ed
high rise site datasets.	-5
Table 2-6. Number of valid ADM and mean NH ₃ emission values between the 2010 and revise	ed
manure belt house datasets2-	
Table 2-7. Number of valid ADM and mean H ₂ S emission values between the 2010 and revise	ed
manure belt house datasets2-	
Table 2-8. Number of valid ADM and mean PM ₁₀ emission values between the 2010 and revise	ed
manure belt house datasets	-6
Table 2-9. Number of valid ADM and mean PM _{2.5} emission values between the 2010 and revise	ed
manure belt house datasets	
Table 2-10. Number of valid ADM and mean TSP emission values between the 2010 and revise	ed
manure belt house datasets2-	-7
Table 2-11. Number of valid ADM and mean emission values between the 2010 and revise	
manure shed datasets	-7
Table 2-12. Comparison of NH ₃ emissions in the EEM dataset to datasets published in peer-revie	w
journal papers2	
Table 2-13. Comparison of H ₂ S emissions in the EEM dataset to datasets published in peer-revie	w
journal papers	
Table 2-14. Comparison of PM ₁₀ emissions in the EEM dataset to datasets published in pee	
review journal papers. 2-1	
Table 2-15. Comparison of PM _{2.5} emissions in the EEM dataset to datasets published in pee	
review journal papers. 2-1	
Table 2-16. Comparison of TSP emissions in the EEM dataset to datasets published in peer-revie	
journal papers.	
Table 2-2-17. Comparison of emissions in the manure belt EEM dataset to datasets published	
peer-review journal papers	
Table 4-1. Bird specific parameters regression analysis for high rise houses	
Table 4-2. House specific parameters regression analysis for high rise houses	
Table 4-3. Ambient parameters regression analysis for high rise houses	
Table 4-4. Bird specific parameters regression analysis for manure belt houses4-1	
Table 4-5. House specific parameters regression analysis for manure belt houses	
Table 4-6. Ambient parameter regression analysis for manure belt houses	
Table 4-8. House and ambient parameter regression analysis for manure sheds	
Table 4-10. Manure parameter regression analysis for manure belt houses4-1	
Table T-10. Intallife parameter regression analysis for manufe out nouses	ı /

Table 4-11. Manure parameter regression analysis for manure belt houses	4-19
Table 5-1. Parameter combinations tested as models for NH ₃ and H ₂ S emissions	5-1
Table 5-2. Parameter combinations tested as models for PM ₁₀ , PM _{2.5} , and TSP emissions	
Table 5-3. Selected daily models for high rise layer houses	
Table 5-4. Parameter combinations tested as models for manure belt house NH ₃ and	
emissions.	
Table 5-5. Parameter combinations tested as models for manure belt house PM ₁₀ , PM _{2.5} , and	
emissions.	
Table 5-6. Selected daily models for manure belt layer houses.	
Table 5-7. Parameter combinations tested as models for layer manure shed NH ₃ and	
emissions	
Table 5-8. Parameter combinations tested as models for layer manure shed PM ₁₀ , PM _{2.5} , and	
emissions.	
Table 5-9. Selected daily models for layer manure sheds.	5-11
Table 6-1. Model coefficients developed using the jackknife approach for NH ₃ emissions	
high rise houses	
Table 6-2. Model fit statistics for the high rise house NH ₃ jackknife	
Table 6-3. Model coefficients developed using the jackknife approach for H ₂ S emissions from	
rise houses.	
Table 6-4. Model fit statistics for the high rise house H ₂ S jackknife.	6-4
Table 6-5. Model coefficients developed using the jackknife approach for PM ₁₀ emissions	
high rise houses.	
Table 6-6. Model fit statistics for the high rise house PM ₁₀ jackknife	6-6
Table 6-7. Model coefficients developed using the jackknife approach for PM _{2.5} emissions	from
high rise houses	
Table 6-8. Model fit statistics for the high rise house PM _{2.5} jackknife	6-8
Table 6-9. Model coefficients developed using the jackknife approach for TSP emissions	from
high rise houses	6-10
Table 6-10. Model fit statistics for the high rise house TSP jackknife	6-10
Table 6-11. Model coefficients developed using the jackknife approach for NH ₃ emissions	from
manure belt houses	6-12
Table 6-12. Model fit statistics for the manure belt house NH ₃ jackknife	6-12
Table 6-13. Model coefficients developed using the jackknife approach for H ₂ S emissions	from
manure belt houses	
Table 6-14. Model fit statistics for the manure belt house H ₂ S jackknife	6-13
Table 6-15. Model coefficients developed using the jackknife approach for PM ₁₀ emissions	from
manure belt houses	
Table 6-16. Model fit statistics for the manure belt house PM ₁₀ jackknife	6-15
Table 6-17. Model coefficients developed using the jackknife approach for PM _{2.5} emissions	from
manure belt houses	
Table 6-18. Model fit statistics for the manure belt house PM _{2.5} jackknife	6-16
Table 6-19. Model coefficients developed using the jackknife approach for TSP emissions	from
manure belt houses	
Table 6-20. Model fit statistics for the manure belt house TSP jackknife	6-17
Table 7-1. Back transformation parameters	
Table 7-2. Annual Uncertainty Model Details	7-2

Table 8-1. Daily calculation parameter values	8-2
Table 8-2. Summary of temperature at the two meteorological sites	8-8
Table 8-3. Summary of relative humidity at the two meteorological sites	8-8
Table 8-4. Total annual emission from the theoretical high rise houses in IA and GA	8-9
Table 8-5. Total annual emission from a theoretical manure belt house in IA and GA	8-10
Table 8-6. Total annual emission from a theoretical manure shed in IA and GA	8-11
Table 8-7. Parameter ranges tested for the high rise house models	8-12
Table 8-8. Parameter ranges tested for the manure belt model	8-15
Table 8-9. Parameter ranges tested for the manure shed model.	8-18
Table 8-10. NH ₃ Emission factors for high rise houses from literature	8-21
Table 8-11. Comparison of resulting high rise house NH ₃ emission from various	estimation
methods	8-22
Table 8-12. NH ₃ Emission factors for manure belt houses from literature	8-22
Table 8-13. Comparison of resulting manure belt house NH ₃ emission from various	estimation
methods	8-23
Table 8-14. Model performance evaluation statistics for high rise houses	8-23
Table 8-15. Model performance evaluation statistics by season	8-24

List of Figures

Figure 1-1. CA2B Farm layout.	1-2
Figure 1-2. IN2B/IN2H Facility layout.	
Figure 1-3. NC3B facility layout. Monitored buildings were houses 1, 2 and 3	
Figure 2-1. Ratio of mean predicted emissions for portion of day with valid	
measurements to mean predicted emissions for the complete day at the finishing (A) a	and sow (B)
farm. Error plotted against number of valid 30-minute measurements (from Grant et	
	2-2
Figure 6-1. Comparison of variation in coefficients and standard errors for NH ₃ high	n rise house
model	
Figure 6-2. Comparison of variation in coefficients and standard errors for H ₂ S high	ı rise house
model	
Figure 6-3. Comparison of variation in coefficients and standard errors for PM ₁₀ high	1 rise house
model	
Figure 6-4. Comparison of variation in coefficients and standard errors for PM _{2.5} high	
model	
Figure 6-5. Comparison of variation in coefficients and standard errors for TSP high	
model	6-11
Figure 6-6. Comparison of variation in coefficients and standard errors for NH ₃ manure	e belt house
model	
Figure 6-7. Comparison of variation in coefficients and standard errors for H ₂ S manure	
model	
Figure 6-8. Comparison of variation in coefficients and standard errors for PM_{10} manur	
model	6-15
Figure 6-9. Comparison of variation in coefficients and standard errors for PM _{2.5} manur	e belt house
model	
Figure 6-10. Comparison of variation in coefficients and standard errors for TSP manur	
model	
Figure 8-1. 2017 Census of Agriculture plot indicating areas of broiler sales.	
Figure 8-2. Comparison on temperatures at test locations in Iowa (IA) and Georgia (G	
Figure 8-3. Comparison of deily emission at test locations IA and GA	
Figure 8-4. Comparison of daily emission at test high rise locations IA and GA	
Figure 8-5. Comparison of daily emission at a theoretical manure belt house in IA and Figure 8-6. Comparison of daily emission at a theoretical manure shed in IA and GA.	
Figure 8-7. High rise house limitation tests for gaseous pollutants.	
Figure 8-8. High rise house limitation tests for particulate matter	
Figure 8-10. Manure belt limitation tests for gaseous pollutants	
Figure 8-11. Manure belt limitation tests for particulate matter.	
Figure 8-12. Maximum values at which the manure belt house models yield negative	
rigure 8-12. Waximum values at which the manure bert house models yield negative	
Figure 8-13. Manure shed limitation tests for gaseous pollutants.	
Figure 8-14. Manure shed limitation tests for particulate matter	
Figure 8-15. Maximum values at which the manure belt house models yield negative	
Tigure 6 15. Waximam values at which the manare best house models yield negative	

Figure 8-16. Scatter plot of the observed NH ₃ emissions at the APECAB IN high rise site versus
the emission model estimates8-25
Figure 8-17. Scatter plot of the observed H ₂ S emissions at the APECAB IN high rise site versus
the emission model estimates8-26
Figure 8-18. Scatter plot of the observed PM ₁₀ emissions at the APECAB IN high rise site versus
the emission model estimates8-27
Figure 8-19. Overhead view of IN2H/IN2B, with in the PM inlet measurement location indicated
by a triangle. (Ni et al. 2010a)
Figure 8-20. Overhead (top) and side (bottom) view of sensor measurements at IN2H (Ni et al
2010a)
Figure 8-21. Overhead (a.) and side (b.) view of sensor measurements at WA5A barn (Ramirez-
Dorronsoro et al. 2010)
Figure 8-22. Overhead view of sensor measurements at IN2B (Ni et al. 2010b)8-31

GLOSSARY / ACRONYMS

-2LogL negative twice the likelihood

ADMs average daily means
AFO animal feeding operation
AIC Akaike information criterion

AICc adjusted Akaike information criterion
BIC Schwarz Bayesian Information Criterion
FANS Fan Assessment Numeration System

H₂S hydrogen sulfide LAW live animal weight

MB mean bias
ME mean error

NAEMS National Air Emissions Monitoring Study

NH₃ ammonia

NMB normalized mean bias

NME normalized mean error

PI Principal Investigator

PM particulate matter

PM₁₀ particulate matter with aerodynamic diameters less than 10 micrometers

PM_{2.5} PM with aerodynamic diameters less than 2.5 micrometers

QAPP quality assurance project plan

QC quality control

TAN total ammoniacal nitrogen

TEOM tapered element oscillating microbalance

TKN total Kjeldahl nitrogen
TSP total suspended particulate
USDA U.S. Department of Agriculture
VOCs volatile organic compounds

1.0 INTRODUCTION

1.1 Site descriptions

There were eight layer houses and one manure shed monitored during NAEMS. The site locations were in California (CA2B), Indiana (IN2H/IN2B) and North Carolina (NC2B). Table 1-1 summarizes sites and the structures monitored. The following section provides additional detail on the sites.

Table 1-1. Layer Confinement Sites Monitored Under NAEMS

Site	Site Type	Monitoring Period	Ventilation type	Number of units measured	Manure Collection	Manure storage ²
CA2B	High-rise	10/17/07 - 10/31/09	MV (sidewall)	2	DB	Inside
IN2B ¹	High-rise	6/1/07 - 5/31/09	MV (sidewall)	2	CBC	First floor
IINZD	Manure belt	1/1/08 - 12/31/09	MV (sidewall)	2	Belt	Shed
IN2H	Manure shed	1/1/08 - 12/31/09	MV	1	Loader	-
NC2B	High-rise	10/17/07 - 10/31/09	MV (tunnel)	2	СВС	Inside

¹House sites that also have measured manure shed

MV = mechanically ventilated

CBC = Curtain backed cages

DB = dropping boards under cages

1.1.1 CA2B

The CA2B layer site was located in central California. The monitored houses, H5 and H6, were built in 2003. The cluster initially consisted of three houses built in 2003, but a fourth house was added in the summer of 2008, during the monitoring period (Heber et al., 2012). In addition, there was a storage lagoon for temporarily holding egg-wash water (not monitored in NAEMS). At this four-house layer cluster, one building (144 m long, 15 m wide, 6.7 m high sidewalls, and 9.1 m high ridge) was selected as the monitoring site (Liang, 2015), which contained two separate (distinct) and individually-ventilated high-rise houses (H5 and H6) (7.5 m wide), each of identical design, and capacity of 38,000 hens (Lohmann LSL Lite) (Cortus et al., 2010; Lin et al., 2012b). Importantly, H5 and H6 are identical in building design, feed regime, manure management, and ventilation (Lin et al., 2012b). H5 and H6 use board scraper systems for manure collection. In this system, the manure collects on dropping boards under the cages and is then scraped into the first floor, where it is stored for six to eight months. H5 and H6 are mechanically ventilated and have misting systems that are used in the summer. A schematic of CA2B and the monitored houses is provided in Figure 1-1. The particulate matter sampling schedule is provided in Appendix A, Table A-1.

²Characterizes type of farm, not necessarily a measurement location.




Figure 1-1. CA2B Farm layout.

1.1.2 IN2B/IN2H

The IN2H/IN2B layer site was in eastern Indiana. This site provided the unique opportunity to monitor the two most common housing and manure management types at one location. The egg production farm consists of an egg-processing plant, two high-rise caged-layer houses, seven manure belt caged-layer houses, two cage-free layer houses, and one free-standing manure shed (Ni et al., 2010a; Ni et al., 2010b; Ni et al., 2017a). A schematic of IN2H/IN2B is provided in Figure 1-2. The high rise site, IN2H, consisted of monitoring at the two high rise houses, H6 and H7. The belt-battery, or manure belt system, monitoring site, IN2B, consisted of two of the farm's manure belt houses, H8 and H9, and the manure shed.

The monitored high rise houses were built in 1997. Each high-rise house is 198 m long and 30.5 m wide (Ni et al., 2010a). All houses are oriented east-west and spaced 17 to 18 m apart (Ni et al., 2010a). The high-rise houses had a capacity of 218,050 hens in ten rows of [Big Dutchman 520N] A-frame cages (5 tiers high) on the upper floor (Ni et al., 2017a). The houses were mechanically ventilated and had a direct drop manure collection method where manure drops off slanted boards behind the cages directly into the first floor, where it is stored for up to one year. The particulate matter sampling schedule is provided in Appendix A, Table A-2.

The monitored manure belt houses were built in 2004. The monitored houses were mechanically ventilated and measured 140 m long and 19.5 m wide, with 7-m sidewalls (Ni et al., 2010b). The houses had capacities of 280,000 birds housed in seven 10-tier rows with H-

frame cages (Ni et al., 2010b). The manure shed for houses 8 and 9 was 85 m long and 30.5 m wide and naturally ventilated via two 0.6-m (2-ft) wide ventilation openings that run the length of the east and west sides (Ni et al., 2010). Manure was collected on 1.21-m wide plastic belts that were under each tier of cages. The manure belt system was manually operated for approximately 4 hours in the morning to move the manure 1/3 of the total belt length from west to east. The longest time that any manure stayed in the house was three days. The manure was then conveyed into manure drying tunnels by three belts at the east end of the house. The particulate matter sampling schedule is provided in Appendix A, Table A-3.




Figure 1-2. IN2B/IN2H Facility layout.

1.1.3 NC2B

The NC2B layer site was located in eastern North Carolina. The monitored houses here, H3 and H4, were built in 2002 and are 18 m wide and 175 m long. At the time of NAEMS, this farm consisted of nine egg-layer houses, one egg-processing (packing) plant, two wastewater treatment lagoons with solid traps, and a wastewater spray field (Wang-Li et al., 2013a). The aerated pond, used for temporarily holding egg-wash water at this facility, was not monitored in

NAEMS. At NC2B, two monitored tunnel-ventilated high-rise houses (H3 and H4) had an inventory of 95,000 hens and 34 exhaust fans (SW-NE) on opposite end-walls with sixteen of these exhaust fans located at the cage level (top floor) (Wang Li et al., 2013a; 2013b). Layers were placed in six rows of 4-tiered curtain-backed cages on the upper floor. Manure falls onto the curtain backed cages and then down into the first floor, where it is stored for up to one year.

The NC2B site was a comprehensive environmental management system (EMS) and complied with International Organization for Standardizations (ISO) 14000 standards (Wang-Li et al., 2013a). The ISO 14000 is a series of international, voluntary environmental management standards, guides, and technical reports (Wall et al., 2001; Feldman and Tibor, 1996). For example, the monitored houses at NC2B are controlled by a computerized environmental control system with ISO 14000 implementations (Wang et al., 2010). A schematic of NC2B is provided in Figure 1-3. The particulate matter sampling schedule is provided in Appendix A, Table A-4.




Figure 1-3. NC3B facility layout. Monitored buildings were houses 1, 2 and 3.

1.2 Data Sampled

NAEMS collected a host of data from the sites. Data collected included gaseous pollutant samples, particulate matter samples, meteorological data, confinement parameters, and biomaterial samples. All procedures were outlined in the project Quality Assurance Project Plan (QAPP) (Heber, 2008), and are summarized in Section 4 of the main report. The following section outlines any collection specific to the layer sites.

1.2.1 Animal Husbandry

Weekly layer inventories, feed and water consumption, egg production and characteristics and bird mass data were collected from the farm's computer system for each site.

1.2.2 Biomaterials Sampling Methods and Schedule

Surface manure samples were collected to determine pH, moisture content and total ammoniacal nitrogen. In addition to surface manure, loadout manure was sampled during each full cleanout of the houses and were analyzed for pH, moisture content, total N, and ammoniacal N. All analyses of biomaterials were performed by an independent laboratory (Midwest Laboratories, Omaha, NE).

Sampling frequency varied between the sites. For CA2B, manure sampled from the first floor storage 5 times at H5 and 8 times at H6. Sampling of load out material occurred 3 times at CA1B H5 and 4 times at CA1B H6. At IN2H, the in-house manure sampling was approximately every three months, on a total of 8 days for each house. The load-out manure was only sampled when the manure was loaded out, which only occurred once during the two year monitoring period. For NC2B, H3 and H4 were sampled on 6 and 5 days, respectively. Dates were randomly spaced across the study period. Load out material was sampled on three different dates from each house.

For the manure belt site, IN2B, samples were collected from 5 locations: 1) the belts in the house, 2) the drying tunnel inlet, 3) the drying tunnel outlet, 4) the manure shed, and 5) the manure shed load out material. Manure from the houses, drying tunnel, and shed were sampled every 60 days. Manure from the shed load out were sampled twice during NAEMS. Per the SOP (Hanni & Bogan 2008), a block random sampling procedure was used to take the manure surface samples. Each windrow was divided into multiple sections per house. A computer program randomly selected sections to be sampled. The samples of approximately equal weight were randomly collected from each section. These samples were mixed thoroughly, and 12 to 15 samples (about ½ kg each) were taken from the mixture and sent to the lab for analysis.

Loadout manure samples were taken from random locations in the manure piles outside of each house or from the trucks used during load out event. Multiple samples were collected, and then combined and mixed. A subsample from this mixed collection sample was sent to the lab for analysis.

2.0 REVISIONS TO DATASET AND EMISSIONS DATA SUMMARY

2.1 Revisions to the 2010 Dataset

As described in Section 4.2 of the main report, NAEMS monitoring data were submitted to EPA in 2010, with revisions submitted in 2015. Revisions included an adjustment to methodology to determine house gas inlet concentrations, which reduced the number of negative emission calculations due to occasionally high inlet concentrations. A more detailed description of these changes can be found in the main report.

Further site-specific revisions include re-calculating negative emission data at IN2H;, PM concentrations and emissions at NC2B; and airflow at IN2B to dry standard conditions.

In 2018, EPA received additional data associated with the NC2B dataset from Dr. Albert Heber. At NC2B, monitoring continued for an additional three months (until 12/31/09) beyond the NAEMS monitoring dates (9/25/07-9/30/09) and were included since the investigators continued to follow NAEMS QA/QC procedures. In addition, revised environmental and production data were received for NC2B, which included revised values for a range of variables such as inventory, live animal weight, exhaust temperature, house temperature and ambient temperature. Revised values for inventory, live animal weight, exhaust temperature, house temperature and ambient temperature were also received for IN2H and CA2B. A description of the revisions is provided in Liang (2015).

EPA reviewed the datasets and removed a small amount of individual environmental data points that were erroneous. In addition, EPA corrected a small amount of production values where there was inconsistency between inventory, hen weight, live animal weight and flock status. Furthermore, in 2020, EPA received additional inventory data for IN2B, which was used to fill-in blank inventory data during a flock replacement event (10/4/08-10/24/08) at H9. This inventory data were also used in the Ni et al. (2017b) publication and was determined based on CO₂ production (Heber, personal communication). This information was requested by EPA as there is a limited range of inventory values in the layer-manure belt dataset and this was the only flock replacement event at the monitored houses during NAEMS.

While performing the exploratory data analysis and reviewing model performance for outliers, EPA developed criteria for the removal of additional negative values from the dataset as part of an outlier analysis. Appendix B outlines the method and the number of additional data points excluded from the layer datasets.

2.2 Data Completeness Criteria for the Revised Dataset

The appropriate data completeness criteria to use in a study depends on the size of the dataset and the accuracy needed. A study by Grant et al. (2013), in which NH₃ emissions were modeled from swine lagoons based on NAEMS data, investigated data completeness and associated accuracy. The swine lagoon NH₃ emissions dataset had limited data availability at a data completeness of 75%. Grant et al. (2013) explored how much the data completeness criteria could be relaxed but still result in data with acceptable error. The study suggested an error of $\pm 25\%$ to be acceptable and determined that a daily data completeness of 52% (or 25 out of 48 30-minute periods) gave less than $\pm 25\%$ error (see Figure 2-1). Using this relaxed daily completeness criteria resulted in a substantial increase in the size of the dataset.

Based on Figure 3-1 from the Grant et al. (2013) study, it can be observed that a daily completeness criterion of 75% (36 out of 48 30-minute periods) would give an error of approximately 10%. If it is assumed that the relationship between data completeness and error from the Grant et al. (2013) study is representative of other NAEMS datasets, the effect of relaxed data completeness criteria can be investigated for other NAEMS sources.

The following sections examine the effect of a reduced data completeness criterion on the number of valid average daily means (ADM) for both the layer houses and manure shed, based on additional analysis completed by Heber that examined the effect of different completeness criteria by comparing the number of valid ADM.

EPA reviewed this data for the egg-layer sites and retained the 75% completeness criterion for all sites. The full analysis can be found in Appendix D.




Figure 2-1. Ratio of mean predicted emissions for portion of day with valid emissions measurements to mean predicted emissions for the complete day at the finishing (A) and sow (B) farm. Error plotted against number of valid 30-minute measurements (from Grant et al., 2013c).

2.3 Comparison between the 2010 and Revised Datasets

The influence of the previous described corrections on the revised dataset can be observed by comparing the number of valid ADM and mean emission values (at 75% data completeness) between the 2010 dataset, as summarized in the final site reports, and the revised dataset. The following sections describe the differences in the ADM for each pollutant between the 2010 data and the revised dataset used in this analysis.

2.3.1 NH₃ High Rise Dataset

The influence of the previous described corrections on the revised dataset can be observed by comparing the number of valid ADM and mean emission values (at 75% data completeness) between the 2010 and revised datasets (Table 2-1). At CA2B the number of valid ADM decreased by 13 (2.2%) for both H5 and H6 with mean NH3 emissions increasing by 1.2% for H5 and 1.8% for H6. At NC2B the number of valid ADM increased by 102 (16.6%) and 104 (17.0%) for H3 and H4 (due to additional data, see section 2.1), respectively, with the mean emissions decreasing by 7.3% for H3 and 1.0% for H4. At IN2H, there was also an increase in number of valid ADM, with number of ADM values increasing for both houses in the revised dataset (39 (7.4%) for H6 and 58 (11.3%) for H7). In terms of the effect on mean NH3 emissions, IN2H had the smallest changes in mean emissions with H6 increasing by 0.6% and H7 decreasing by 0.4%.

Table 2-1. Number of valid ADM and mean NH₃ emission values (at 75% data completeness) between the 2010 and revised high rise datasets.

Dataset	Statistic	CA2B H5	CA2B H6	IN2H H6	IN2H H7	NC2B H3	NC2B H4
2010	n of ADM	583	603	525	512	613	611
	Overall ADM (kg d ⁻¹)	32.7	31.7	223.3	249.3	62.5	58.1
Revised	n of ADM	570	590	564	570	715	715
	Overall ADM (kg d ⁻¹)	33.1	32.3	224.7	248.2	58.1	57.5

2.3.2 H₂S High Rise Dataset

The comparison of the number of valid ADM and mean emission values (at 75% data completeness) between the 2010 and revised datasets is provided in Table 2-2. At CA2B, the number of valid ADM decreased by 13 (2.1%) for both H5 and H6 with mean H₂S emissions increasing by 0.02% for H5 and 0.4% for H6. At NC2B, the number of valid ADM increased by 21 (3.3%) and 28 (4.4%) for H3 and H4, respectively, with the mean emissions increasing by 0.6% for H3 and 3.9% for H4. At IN2H, there was the largest change in number of valid ADM with the number of ADM values increasing for both houses in the revised dataset (43 (12.0%) for H6 and 41 (11.1%) for H7).

In terms of the effect on mean H_2S emissions, IN2H had the largest changes in mean emissions with H6 increasing by 8.9% and H7 increasing by 4.8%. Additional data provided for the NC2B site increased the number of valid ADM by 39 (5.6%) and 38 (5.4%), respectively. The additional data resulted in the mean emission decreasing by 0.2% at H3 and 4.7% at H4.

Table 2-2. Number of valid ADM and mean H₂S emission values (at 75% data completeness) between the 2010 and revised high rise datasets.

Dataset	Statistic	CA2B H5	CA2B H6	IN2H H6	IN2H H7	NC2B H3	NC2B H4
2010	N of ADM	614	633	358	369	641	635
	Overall ADM (kg d ⁻¹)	45.40	39.80	277	257	57.10	62.80
Revised	N of ADM	601	620	401	410	662	663
	Overall ADM (kg d ⁻¹)	45.41	39.96	301.54	269.27	57.45	65.24
Additional	N of ADM	-	-	1	-	701	701
Data	Overall ADM (kg d ⁻¹)	-	-	-	-	57.36	62.31

2.3.3 PM High Rise Dataset

Table 2-3 provides a summary of the number of valid ADM (N of ADM) and the overall ADM for PM₁₀ at each site. At CA2B, the number of valid ADM remained the same for both H5 and H6 with mean PM₁₀ emissions increasing by 0.2% for H5 and no change for H6. At IN2H, the number of valid ADM values increased for both H6 (6 days, 1.5%) and H7 (18 days, 4.5%). In terms of the effect on mean H₂S PM10 emissions, H6 decreased by 0.6% and H7 decreased by 0.9%. For NC2B, the number of valid ADM increased by 6 (1.6%) and 47 (9.1%) at H3 and H4, respectively. The additional valid ADM values resulted in the mean emission increasing by 2.5% at H3 and decreasing by 0.9% at H4.

Table 2-3. Number of valid ADM and mean PM₁₀ emission values between the 2010 and revised high rise site datasets.

Dataset	Statistic	CA2B H5	CA2B H6	IN2H H6	IN2H H7	NC2B H3	NC2B H4
2010	N of ADM	451	527	411	403	377	518
	Overall ADM (g d ⁻¹)	1,270	960	3,702	4,944	1,486	2,219
Revised	N of ADM	451	527	417	421	383	565
	Overall ADM (g d ⁻¹)	1,273	960	3,678	4,898	1,523	2,200

Table 2-4 provides a summary of the number of valid ADM and the overall ADM for PM_{2.5} at each site. The number of valid ADM values remained the same for both houses at CA2B, H6 at IN2H, and H3 at NC2B. However, the ADM did decrease at CA2B H5 by 0.4%, IN2H H6 by 3.3%, and 38% at NC2B H3. Emissions at CA2B H6 remained the same. The number of valid ADM at IN2H H7 decreased by 1 (10%), which corresponded to a 1.9% decrease in the ADM. The largest difference occurred at NC2B H4, where the number of valid ADM values increased by 15 days (45.5%), which resulted in a 12.1% decrease in the ADM.

Table 2-4. Number of valid ADM and mean PM_{2.5} emission values between the 2010 and revised high rise site datasets.

Dataset	Statistic	CA2B H5	CA2B H6	IN2H H6	IN2H H7	NC2B H3	NC2B H4
2010	N of ADM	40	43	16	10	21	33
	Overall ADM (g d ⁻¹)	238	168	214	104	50	165
Revised	N of ADM	40	43	16	9	21	48
	Overall ADM (g d ⁻¹)	237	168	207	102	31	145

Similar to the PM_{2.5} results, the number of ADM did not change much for most of the sites (see Table 2-5). Both CA2B houses and IN2H H6 saw no change in the number of ADM available. However, there were changes in the overall ADM. CA2B H5 had a 0.1% decrease, while CA2B H6 and IN2H H6 had 0.2% and 5.3% increase, respectively. IN2H H7 and NC2B H3 had small changes in the number of ADM available, increasing by 2 (10.5%) and 1 (2.3%) days, respectively. These corresponded to a 0.9% decrease in overall ADM at IN2H H7 and a 1.3% increase at NC2B H3. The final site, NC2B H4, saw the largest change, as the number of ADM increased by 45 (40.3%) which corresponded to a 5.4% decrease in overall ADM.

Table 2-5. Number of valid ADM and mean TSP emission values between the 2010 and revised high rise site datasets.

Dataset	Statistic	CA2B H5	CA2B H6	IN2H H6	IN2H H7	NC2B H3	NC2B H4
2010	N of ADM	36	32	19	19	44	62
	Overall ADM (g d ⁻¹)	2,440	2,760	7,408	4,694	3,391	4,385
Revised	N of ADM	36	32	19	21	45	87
	Overall ADM (g d ⁻¹)	2,437	2,765	7,803	4,652	3,434	4,148

2.3.4 NH₃ Manure Belt House Dataset

For NH₃ emissions at the manure belt site, the changes made in the revised dataset were relatively minor (Table 2-6). At CA2B the number of valid ADM decreased by 3 (0.5%) for both H8 while the number of valid days remained the same at H9. Mean NH₃ emissions increased by less than 1 kg at each house, for a 1.4% and 0.8% change for H8 and H9, respectively.

Table 2-6. Number of valid ADM and mean NH₃ emission values between the 2010 and revised manure belt house datasets.

Dataset	Statistic	IN2B H8	IN2B H9
2010	N of ADM	624	629
	Overall ADM (kg d ⁻¹)	70.6	66.5
Revised	N of ADM	621	629
	Overall ADM (kg d ⁻¹)	71.5	67.0

2.3.5 H₂S Manure belt Dataset

The changes in the H_2S dataset for the manure belt house made between the 2015 revision to the dataset were relatively minor (Table 2-7). At CA2B the number of valid ADM decreased by 3 (0.5%) for H8 while the number of valid days remained the same at H9. Mean H_2S emissions increased in the revised dataset by 0.6% for H3 and 0.4% for H4.

Table 2-7. Number of valid ADM and mean H₂S emission values between the 2010 and revised manure belt house datasets.

Dataset	Statistic	IN2B H8	IN2B H9
2010	N of ADM	634	645
	Overall ADM (g d ⁻¹)	489.0	469.2
Revised	N of ADM	631	645
	Overall ADM (g d ⁻¹)	492.1	471.1

2.3.6 PM Manure Belt House Dataset

The emissions dataset for PM₁₀ (Table 2-8), PM_{2.5} (Table 2-9), and TSP (Table 2-10) were unchanged between the original 2010 submission and the revision submitted in 2015 by Dr. Heber. The comparison here does not include the exclusions implemented by EPA.

Table 2-8. Number of valid ADM and mean PM₁₀ emission values between the 2010 and revised manure belt house datasets.

Dataset	Statistic	IN2B H8	IN2B H9
2010	N of ADM	346	361
	Overall ADM (g d ⁻¹)	2,209.2	6,076.2
Revised	N of ADM	346	361
	Overall ADM (g d ⁻¹)	2,209.2	6,076.2

Table 2-9. Number of valid ADM and mean PM_{2.5} emission values between the 2010 and revised manure belt house datasets.

Dataset	Statistic	IN2B H8	IN2B H9
2010	N of ADM	25	31
	Overall ADM (g d ⁻¹)	-85.1	113.2
Revised	N of ADM	25	31
	Overall ADM (g d ⁻¹)	-85.1	113.2

Table 2-10. Number of valid ADM and mean TSP emission values between the 2010 and revised manure belt house datasets.

Dataset	Statistic	IN2B H8	IN2B H9
2010	N of ADM	35	34
	Overall ADM (g d ⁻¹)	8,136.3	21,871.0
Revised	N of ADM	35	34
	Overall ADM (g d ⁻¹)	8,136.3	21,871.0

2.3.7 Manure Shed Dataset

The emissions dataset for manure sheds (Table 2-11) also remained unchanged between the original 2010 submission and the revision submitted in 2015 by Dr. Heber. The comparison here does not include the exclusions implemented by EPA.

Table 2-11. Number of valid ADM and mean emission values between the 2010 and revised manure shed datasets.

Dataset	Statistic	NH₃	H₂S	PM ₁₀	PM _{2.5}	TSP
2010	N of ADM	518	534	307	30	24
	Overall ADM (g d ⁻¹)	5	35	134	48	295
Revised	N of ADM	518	534	307	30	24
	Overall ADM (g d ⁻¹)	5	36	134	48	295

2.4 Comparison Between the Revised Datasets and NAEMS Datasets Used in Peer-reviewed Published Papers

Where possible, EPA compared the revised dataset developed for this report to values presented in peer reviewed journals to quantify any differences due to the application of the revised calculation methods and other adjustments discussed in Section 2.1.

2.4.1 High Rise Houses

Summaries of the NH₃ emissions from CA2B, IN2H, and NC2B high rise layer houses have been published in peer-reviewed journal articles (Lin et al., 2012a; Ni et al., 2017a; Wang-Li et al. 2013b). A simple comparison of the summary statistics presented in these papers and the summary statistics of the dataset used to develop the emission models is presented in Table 2-12. For CA2B, the number of ADM is less than in the article by 2%. This resulted in a 1.3% and 2.3% difference in the mean at H5 and H6, respectively. For IN2H, differences in the means are minor (less than 1%) despite an increase of 39 and 58 daily means at H6 and H7, respectively. For NC2B, the revised EEM dataset has 64 and 56 more ADM values for H3 and H4, respectively, than in comparison to the Wang-Li et al. (2013b) study. However, the number of ADMs reported by Wang-Li et al.(2013b) is for full and active bird status only, whereas the

revised EEM dataset includes ADMs for all bird status. In the revised EEM dataset, the mean ADM values were 1.5% (H3) and 4.2% (H4) higher than in the Wang-Li et al. (2013b) study.

Table 2-12. Comparison of NH₃ emissions in the EEM dataset to datasets published in peer-review journal papers.

Site	Units	Statistic	EEM Dataset	Previous Studies	Study
		Number of ADM	570	583	
CA2B H5	Emissions	Mean	0.963	0.95	Lin et al.,
CAZB IIS	(g day ⁻¹ hd ⁻¹)	Standard Deviation	0.494	0.49	2012a
		Max	2.34	2.28	
		Number of ADM	590	602	
CA2B H6	Emissions	Mean	0.962	0.94	Lin et al.,
CAZB NO	(g day ⁻¹ hd ⁻¹)	Standard Deviation	0.872	0.67	2012a
		Max	3.95	3.97	
	Emissions (kg day ⁻¹)	Number of ADM	564	525	Ni et al
IN2H H6		Mean	225	223	Ni et al., 2017a
		Standard Deviation	90	86	2017a
	Emissions	Number of ADM	570	512	Ni et al
IN2H H7	Emissions (kg day ⁻¹)	Mean	249	249	Ni et al., 2017a
	(kg day)	Standard Deviation	90	97	2017a
	Emissions	Number of ADM	715	651	
NC2B H3	(full and active;	Mean	58.1	57.2	Wang-Li et al.,
	kg day⁻¹)	Standard Deviation	20.9	19.0	2013
	Emissions	Number of ADM	715	659	
NC2B H4	(full and active;	Mean	57.5	55.1	Wang-Li et al.,
	(kg day ⁻¹)	Standard Deviation	24.4	23.3	2013

The H₂S emissions from NAEMS high rise layer houses have been published in peer-reviewed journal articles (Lin et al., 2012a; Ni et al., 2017a; Wang et al., 2016), which is presented in Table 2-13. For CA2B and NC2B, there are only small differences between the datasets with the difference in number of ADM ranging from 1.9-2.1% for all houses at CA2B and 6.9-7.5% at NC2B. Additionally the differences in mean and standard deviations are less than 10% for all houses at CA2B and NC2B. There is, though, a larger difference in the max values between the Lin et al. (2012b) study and this study, which have values of 3.72 (H5) and 4.26 (H6) and 3.80 (H5) and 4.99 (H6), respectively.

With respect to IN2H, there are large differences in the number of ADMs and in the standard deviation between the Ni et al. (2017a) study and this study. The number of available ADM reported in the Ni et al. (2017a) study are 27.7% (H6) and 27.3% (H7) less than this study. This 111 day difference in the daily values available has relatively small changes in the mean values: 4% and 7% for H6 and H7, respectively. However, it has a much larger influence on

standard deviation, with values decreasing by 41.7% (H6) and 22.1% (H7) in comparison to the dataset used to develop the models. This suggests that some of the high or low emission values in the dataset used in this study were not included in the dataset in the Ni et al. (2017a) study.

Table 2-13. Comparison of H₂S emissions in the EEM dataset to datasets published in peer-review journal papers.

			EEM	Previous	
Site	Units	Statistic	Dataset	Studies	Study
CA2B H5	Emissions	Number of ADM	601	614	Lin et al.,
	(mg day ⁻¹ hd ⁻¹)	Mean	1.33	1.33	2012a
		Standard Deviation	0.71	0.70	
		Max	3.80	3.72	
CA2B H6	Emissions	Number of ADM	620	632	Lin et al.,
	(mg day ⁻¹ hd ⁻¹)	Mean	1.20	1.20	2012a
		Standard Deviation	0.89	0.86	
		Max	4.99	4.26	
IN2H H6	Emissions	Number of ADM	401	290	Ni et al.,
	(g day ⁻¹)	Mean	302	314	2017a
		Standard Deviation	278	162	
IN2H H7	Emissions	Number of ADM	410	298	Ni et al.,
	(g day ⁻¹)	Mean	269	287	2017a
		Standard Deviation	281	219	
NC2B H3	Emissions	Number of ADM	701	656	Wang et al.,
	(full and active	Mean	57.4	59.6	2016
	status; g day ⁻¹)	Standard Deviation	35.2	34.7	
		Median	47.7	48.6	
NC2B H4	Emissions	Number of ADM	701	652	Wang et al.,
	(full and active	Mean	62.3	65.4	2016
	status; g day ⁻¹)	Standard Deviation	43.7	41.5	
		Median	49.5	50.5	

The PM₁₀ emissions for all NAEMS high rise layer houses have been published in peer-reviewed journal articles (Lin et al., 2012a; Ni et al., 2017a; Li et al., 2013), which is presented in Table 2-14. The biggest departure across the houses is NC2B H4, which has 171 (30.3%) more ADM available, which translates to a 19% increase in the average value. The main reason for this difference is that Li et al. (2013) does not report daily PM summary statistics for the whole monitoring period at house 4 due to the two different PM sampling strategies used at NC2B H4. For CA2B, the modeling dataset produces summary statistics nearly identical to the Lin et al. (2012b) paper. The exception is EPA included an additional value for house 6. This value was the maximum value in the EPA dataset, and caused an 8.2% increase in the standard deviation. For IN2H houses and NC2B H3, there are only small differences between the datasets, with the difference in number of ADM ranging from 1.4-4.3% increase in the number of ADM

available. This minor increase in ADM available translates to differences in mean and standard deviations less than 2% for these houses.

Table 2-14. Comparison of PM₁₀ emissions in the EEM dataset to datasets published in peer-review journal papers.

Site	Units	Statistic	EEM Dataset	Previous Studies	Study
		Number of ADM	451	451	
CA2B H5	Emissions	Mean	37.6	37.6	Lin et al.,
CAZB IIS	(mg day ⁻¹ hd ⁻¹)	Standard Deviation	30.3	30.4	2012a
		Max	231	231	
		Number of ADM	525	524	
CA2B H6	Emissions	Mean	29.6	29.2	Lin et al.,
CAZB NO	(mg day ⁻¹ hd ⁻¹)	Standard Deviation	26.7	24.5	2012a
		Max	276	143	
	Emissions (g day ⁻¹)	Number of ADM	417	411	
IN2H H6		Mean	3,678	3,687	Ni et al., 2017a
		Standard Deviation	3,230	3,197	2017a
	Emissions (g day ⁻¹)	Number of ADM	421	403	
IN2H H7		Mean	4,898	4,934	Ni et al., 2017a
	(g day)	Standard Deviation	4,004	3,982	2017a
		Number of ADM	383	371	
NICOD LIO	Emissions	Mean	1,523	1,528	Li et al.,
NC2B H3	(g day ⁻¹)	Standard Deviation	636	644	2013
		Median	1,481	1,501	
		Number of ADM	565	394	
NG2D II4	Emissions	Mean	2,200	1,781	Li et al.,
NC2B H4	(g day ⁻¹)	Standard Deviation	1,130	783	2013
		Median	2,016	1,693	•

The PM_{2.5} emissions for CA2B and NC2B houses have been published in peer-reviewed journal articles (Lin et al., 2012a; Li et al., 2013). Searches did not find articles that included summaries of the PM_{2.5} emissions data from IN2H. Table 2-15 presents a summary of the model development dataset and the summary values presented in the articles. For CA2B, the modeling dataset produces summary statistics nearly identical to the Lin et al. (2012a) paper. For NC2B H4, the EEM data set has 27 days more available than the Li et al. (2013) study, which is an increase of 56.3%. This produced a 72.7% increase in the mean and an 59.3% increase in the standard deviation for the house. The main reason for the difference in the number of ADMs is

that Li et al. (2013) does not report daily PM summary statistics for the whole monitoring period at house 4 due to the two different PM sampling strategies used at NC2B H4. Data for H3 at NC2B was nearly identical to the statistics presented in Li et al. (2013), with a 0.1% or less difference across all statistics.

Table 2-15. Comparison of PM_{2.5} emissions in the EEM dataset to datasets published in peer-review journal papers.

C:+-	l l'oite	Ctatiatia	EEM	Previous	CAd	
Site	Units	Statistic	Dataset	Studies	Study	
		Number of ADM	40	40		
CA2B H5	Emissions	Mean	6.7	6.7	Lin et al.,	
CAZBTIS	(mg day ⁻¹ hd ⁻¹)	Standard Deviation	14.9	14.9	2012a	
		Max	53.2	53.2		
		Number of ADM	43	43		
CA2B H6	Emissions (mg day ⁻¹ hd ⁻¹)	Mean	5.17	5.2	Lin et al., 2012a	
CAZBTIO		Standard Deviation	10.3	10.3		
		Max	35.2	35.2		
		Number of ADM	21	21		
NC2B H3	Emissions	Mean	31.1	31.1	Li et al.,	
NCZB II3	(g day ⁻¹)	Standard Deviation	79.0	79	2013	
		Median	48.9	48.9		
		Number of ADM	48	21		
NC2B H4	Emissions	Mean	144.9	39.5	Li et al.,	
INCZB П4	(g day ⁻¹)	Standard Deviation	168.8	68.7	2013	
		Median	81.1	72.3		

Similar to PM_{2.5}, TSP emissions have been published for CA2B and NC2B (Lin et al., 2012a; Li et al., 2013), and no articles were found that included summaries of the TSP emissions data from IN2H. Table 2-16 presents a summary of the model development dataset and the summary values presented in the articles. For CA2B, the modeling dataset produces summary statistics nearly identical to the Lin et al. (2012a) paper. For NC2B H4, the EEM data set has 50 more ADMs (an increase of 57.5%) available than the Li et al. (2013) study. The main reason for this difference is that Li et al. (2013) does not report daily PM summary statistics for the whole monitoring period at house 4 due to the two different PM sampling strategies used at NC2B H4. The EEM dataset had 3% lower mean and 45.6% higher standard deviation in comparison to the Li et al. (2013) study. Data for H3 at NC2B saw an 8 day (17.8%) increase in the number of daily values available, which resulted in a 7.2% decrease in the mean and 1.8% decrease in the standard deviation.

Table 2-16. Comparison of TSP emissions in the EEM dataset to datasets published in peer-review journal papers.

Site	Units	Statistic	EEM	Previous	Chudu	
			Dataset	Studies	Study	
CA2B H5	Emissions (mg day ⁻¹ hd ⁻¹)	Number of ADM	36	36		
		Mean	71.9	71.9	Lin et al.,	
		Standard Deviation	41.0	41	2012a	
		Max	177	177		
СА2В Н6	Emissions (mg day ⁻¹ hd ⁻¹)	Number of ADM	32	32		
		Mean	84.0	84	Lin et al.,	
		Standard Deviation	44.4	44.4	2012 a	
		Max	226	226		
NC2B H3	Emissions (g day ⁻¹)	Number of ADM	45	37		
		Mean	3,434	3,680	Li et al.,	
		Standard Deviation	1,515	1,543	2013	
		Median	3,296	3,606		
NC2B H4	Emissions (g day ⁻¹)	Number of ADM	87	37		
		Mean	4,148	4,273	Li et al.,	
		Standard Deviation	2,429	1,322	2013	
		Median	4,415	4,348		

2.4.2 Manure belt Houses

Summaries of the emissions from the manure belt layer houses monitored during NAEMS have been published in a peer-reviewed journal article by Ni et al. (2017b). The model development dataset is slightly different from the summaries presented in the article, with all statistics reported in Table 2-17 varying by less than 2%.

Table 2-2-17. Comparison of emissions in the manure belt EEM dataset to datasets published in peer-review journal papers.

Pollutant	Site	Units	Statistic	EEM Dataset	Previous Studies	Study
NH ₃	IN2B H8	Emissions (kg day ⁻¹)	Number of ADM	621	624	Ni et al., 2017b
			Mean	71.5	70.6	
			Standard Deviation	37.5	36.8	
	IN2B H9	Emissions (kg day ⁻¹)	Number of ADM	629	629	Ni et al., 2017b
			Mean	67.0	66.5	
			Standard Deviation	43.0	42.2	
H₂S	IN2B H8	Emissions (g day ⁻¹)	Number of ADM	631	634	Ni et al., 2017b
			Mean	492	489	
			Standard Deviation	246	241	
	IN2B H9	Emissions (g day ⁻¹)	Number of ADM	645	645	Ni et al., 2017b
			Mean	471	469	
			Standard Deviation	268	265	
PM ₁₀	IN2B H8	Emissions (g day ⁻¹)	Number of ADM	251	248	Ni et al., 2017b
			Mean	3,039	3,086	
			Standard Deviation	4,813	4,812	
	IN2B H9	Emissions (g day ⁻¹)	Number of ADM	361	361	Ni et al., 2017b
			Mean	6,076	6,076	
			Standard Deviation	8,238	8,226	

2.4.3 Manure Shed

Searches by EPA did not find any articles that included data from the manure shed at IN2B.

3.0 RELATIONSHIPS ESTABLISHED IN LITERATURE

Developing EEMs for AFOs is complex as many variables potentially influence emissions. Therefore, to be efficient in this study, a focused approach was used to develop EEMs. The focused approach involved developing models based on variables that could potentially have a major influence on air emissions. This assessment was made based on theoretical considerations and observations reported by previous studies that have investigated the influence of variables on emissions from swine AFOs.

3.1 NH₃ and H₂S Emissions from Houses and Manure Sheds

The amount of manure produced at a layer house is a key factor influencing NH₃ and H₂S emissions, since this will affect the total amount of NH₃ and H₂S that is generated in the manure (due to microbial degradation of urea, undigested proteins, and amino acids (Mackie et al., 1998)) and released (e.g., movement of gas from manure into the air). Proxies for the amount of fresh manure produced at a layer house are inventory and live animal weight (LAW). Thus, these variables, which were determined daily, were selected for further investigation. For the layermanure shed, there is on average a 5-day gap between manure production in the house and the manure arriving in the manure-shed. Therefore, variables were created and selected for further investigation for the layer manure shed, which represent a 5-day lag of inventory (adjusted inventory) and LAW (adjusted LAW) from the house values. For the manure-shed, inventory, LAW, adjusted LAW and adjusted inventory represented the sum of inventory and LAW from the two houses the shed was receiving manure from. The amount and content of fresh manure produced by layers can also be influenced by feed characteristics, which can be different for layers depending on their age (Wu-Hann et al. 2007; Li et al. 2013). Accordingly, hen age was selected for further investigation by creating a numeric variable that indicated how many days since the hens were brought into the house. For the layer-manure shed, this variable was the weighted average age of the hens in the two houses that the shed was receiving manure from.

In layer high rise houses, the manure is stored on the bottom floor of the house and thus the amount of manure in the house increases with time, which may influence gas emissions (Liang et al. 2005; Lin et al. 2012a; Li et al. 2013). Therefore, a variable for manure age was created with a numeric value that indicated how many days since the date of the last manure cleanout. This variable was similarly created for layer manure sheds, but not for layer-manure belt houses since this manure management system does not store the manure in the house.

Manure characteristics such as nitrogen and sulfur content, solid and moisture content and pH can influence NH₃ and H₂S emissions. Common measurement of nitrogen and sulfur content that relate to NH₃ and H₂S emissions are total kjeldahl nitrogen (TKN; NH₃-N + organic

N), total ammoniacal nitrogen (TAN; NH₃-N), and sulfide. Higher concentrations of these nitrogen and sulfur components indicate a greater potential for NH₃ emissions (Groot Koerkamp, 1994) and H₂S emissions. Manure pH can influence emissions as it can affect both the generation of NH₃ in manure and the concentration of NH₃ in layer manure (Tong et al., 2020). Groot Koerkamp, (1994) reported that as manure pH increases above 5.5, the rate of manure degradation increases. Manure pH also influences NH₃ and H₂S concentrations due to its effect on the chemical equilibrium between NH₃ and NH₄⁺ and HS⁻ and H₂S, respectively. Fresh layer manure is excreted with approximately 75% moisture content (Xin et al., 2011) and dries through movement of air caused by ventilation fans. The rate of drying affects the manure solid and moisture content, which influences NH₃ generation within the manure and thus influences emissions (Groot Koerkamp, 1994; Xin et al. 2011). Lower manure solid content and thus higher moisture content increase the generation of NH₃ increases due to the impact of moisture on microbe growth. It is hypothesized that moisture would have a similar effect on H₂S generation. In NAEMS, manure samples at layer sites were taken at a frequency of 2-5 months on average. Manure samples were analyzed for TAN in all samples, but TKN and sulfide were not measured in surface manure in any of the houses, with the exception of one sample taken at CA2B-H6. TAN was selected for further analysis, but TKN and sulfide could not be selected due to the low number of measurements. All collected manure samples were analyzed for solid content and pH, therefore manure solid content and pH were selected for further investigation. For the manureshed, manure samples were regularly analyzed for TKN concentration in addition to pH, solids content and TAN. Accordingly, for the manure shed, TKN was also selected for further investigation.

House airflow or ventilation rate is a variable that can have a major influence on the emissions of NH₃ and H₂S from manure as it affects the air flow above the manure surface (Tong et al. 2020; Rumsey and Aneja, 2014). An increase in air velocity reduces the boundary layer thickness above the manure surface, thereby lowering the resistance to volatilization and causing an increase in the transfer of NH₃ and H₂S across the air-manure interface (Arogo et al. 1999; Rumsey and Aneja, 2014). However, higher ventilation rates can also dry out layer manure (i.e., increase manure solid content) resulting in lower emission rates (Ni et al. 2017a). House air flow was measured continuously during NAEMS and was chosen for further analysis. The layer-manure shed was naturally ventilated, and therefore the airflow is not necessarily a function of temperature but could be related to wind speed (Joo et al. 2014). Accordingly, air flow and wind speed, which were measured continuously, were selected for further investigation.

Temperature plays a key role in many of the biological, physical, and chemical processes involved in NH₃ and H₂S generation and release processes, and thus influences NH₃ and H₂S emissions from layer manure. Manure temperature influences the microbial degradation of layer

manure, with increasing temperatures resulting in increasing degradation rates (Groot Koerkamp, 1994; Zhao et al. 2013). Increasing manure temperature will also increase the Henry's law constant and dissociation constant for NH₃ and H₂S (Tong et al. 2020; Montes et al. 2009; Rumsey and Aneja, 2014). For NH₃, this increases the potential amount that can be released from the manure into the air. However, for H₂S, an increasing Henry's law constant and dissociation have conflicting effects on the potential amount available, meaning that the overall influence of temperature on H₂S emissions may not be as strong as for NH₃. Increasing manure temperature and air temperature can also increase the transfer of NH₃ and H₂S across the manure-air interface (Ni, 1999, and references within; Montes et al., 2009, and references within; Tong et al. 2020; Rumsey and Aneja, 2014). Note that while the release of NH₃ is controlled by the convective mass transfer release mechanism, the release of H₂S is additionally influenced by bubble-release (ebullition) mechanisms, which can be triggered by manure disturbances (Ni et al., 2009) from animal or management activities inside the house. The effect of temperature on emissions from layer manure is complicated by the relationship between temperature and ventilation rate. In mechanically ventilated houses, increasing ambient and house temperature will result in higher ventilation rates, which as previously described could reduce emissions due to drying of the manure, thus countering the effect temperature has as on other emission processes. During NAEMS, researchers took continuous measurements of house exhaust temperature (temperature at house fan exhaust) and ambient temperature, and both were chosen for further analysis. For the layer manure-shed, no measurements of shed temperature were made, therefore only ambient temperature was selected for further analysis.

Relative humidity may influence NH₃ and H₂S emissions from layer manure through its effect on manure solids content/moisture content as a higher relative humidity may reduce the evaporation of water from the manure surface, resulting in wetter manure and thus higher NH₃ emissions (Ni et al. 2017a). This effect of relative humidity on NH₃ emissions has also been identified in broiler litter, where increasing relative humidity that varied from 40% to 80% (similar to house exhaust relative humidity measured at NAEMS sites (see Appendix D)) was found to increase NH₃ levels (Weaver and Meijerhof, 1991). It is proposed that relative humidity could affect H₂S emissions similarly. During NAEMS, researchers took continuous measurements of exhaust relative humidity and ambient relative humidity, and both were chosen for further analysis. For the layer manure-shed, no measurements of shed relative humidity were made, therefore only ambient relative humidity was selected for further analysis.

Management activities can also influence gas emissions from layer houses (Lim et al. 2003). There are three major management activities that occur in layer-HR houses: flock emptying and replacement, manure cleanouts, and molting. During flock emptying and replacement there will be different numbers of hens in the house, thus influencing the amount of

fresh manure produced. Manure cleanouts involve the removal of manure from the bottom level of the house where the manure is stored. In NAEMS, the manure cleanouts did not occur during periods of flock replacement. During manure cleanouts there is the potential for increases in gas emissions due to the disturbance of manure (Ni et al. 2009). Theoretically, molting can influence NH₃ and H₂S emissions as birds can be given different amounts of feed during molting. In addition, different types of feed can be given to the birds, which could have different nutrient content, thus influencing gas emissions. The influence of flock emptying and replacement, manure cleanouts and molting on layer-HR houses was selected for further investigation. To achieve this, each day for each house was assigned a status of full (F), empty (E), transition to empty or full (T), manure cleanout (C) or molting (M). Similarly, the same statuses with the exception of C status (since manure is not stored in a manure-belt house) were assigned to each day of each house for the manure belt houses. For manure sheds, the only management event that occurs is the removal or cleanout of manure since there are no birds in the shed. Manure cleanouts only occurred for two days during the NAEMS monitoring period. However, house management activities that influence the amount and composition of manure can also potentially affect gas emissions from the manure shed once the manure arrives 5-days later. Therefore, for each day, the manure shed was assigned a status that was a combination of the statuses from 5 days beforehand at house 8 and 9. For example, if house 8 had a status of F and house 9 a status of M, the combined shed status would be FM. Accordingly, each day had a two letter status with the exception of the two days when the manure shed manure clean out occurred. On these days, the letter "C" was added to the two-letter status.

3.2 PM Emissions from House and Manure Sheds

The release of particulate matter into layer house air is caused by the physical suspension of a range of different materials in layer houses including feed, manure, and feathers (Cambra-Lopez et al. 2011). In the manure shed, there are no birds, feed, or recently excreted manure, therefore, the deposition of manure from the belt to the manure pile is probably the main process that causes particulate matter emissions. The amount of particulate matter source materials being transferred from the belt to the manure pile will likely be related to house LAW and inventory, therefore these variables were explored further. Similar to the gases, in addition to the LAW and inventory values, the variables, adjusted inventory and adjusted LAW, which represent a 5-day lag of inventory and LAW from both of the houses were created for further investigation for the layer-manure sheds.

Physical suspension of particulate matter from house surfaces can be caused by animal activity, human activity, and air flow (Aarnink and Ellen, 2007). However, house activity measurements were not provided to EPA, therefore the influence of this variable could not be explored further. Ventilation rate or airflow influences house particulate matter emissions by

controlling the amount of particulate matter sedimentation in a house (Shepherd et al. 2015). In animal houses or houses, mechanical ventilation is typically a function of ambient temperature and thus house temperature, meaning that temperature could serve as a potential surrogate variable for airflow. The physical suspension of particulate matter can also be influenced by moisture conditions and relative humidity (Cambra-Lopez et al. 2010). The moisture content of feces can be affected by bird water consumption with increased water consumption during warmer months leading to wetter feces (Shepherd et al. 2015). While there was no direct measure of this in NAEMS, measurements of manure solid content could be an indicator, therefore manure solid content was selected for investigation. As mentioned, feed characteristics can be different for layers depending on their age (Li et al. 2013). In addition, hen activity can increase with hen age after initial placement (Li et al. 2013). Both of these factors could potentially influence particulate matter emissions; therefore, hen age was selected for further investigation.

The moisture content of the house air can also influence particulate matter emissions. Takai et al. (1998) reported that a relative humidity greater than 70% results in a high equilibrium moisture content and may contribute to particles aggregating together, resulting in lower concentrations and emissions. However, the 70% RH value reported by Takai et al. (1998) is based on the condensation of water on barley or wheat grains, and it is not known how representative this value is for other types of PM (e.g., different types of feed, manure, and feathers/skin). House exhaust daily RH values at NAEMS layer houses varied from 31.8% to 86.2%, with ADM at different sites varying from 48.7% at IN2H to 68.4% at NC2B (see Appendix D). It should be noted that determining the influence of airflow, temperature and relative humidity on emissions is complex due the intrinsic relationship between these variables in a mechanically ventilated animal house. Accordingly, for layer houses, the continuously measured variables, house exhaust relative humidity, ambient relative humidity, house exhaust temperature, ambient temperature and airflow were selected for further investigation for PM₁₀, PM_{2.5} and TSP emissions from layer houses. As mentioned for layer-manure shed, manure-shed temperature and RH were not measured, therefore only the airflow, ambient temperature and ambient relative humidity variables were selected for further investigation. As stated in section 3.1, airflow could be related to wind speed (Joo et al. 2014), therefore wind speed was also selected for further investigation.

Management activities can also affect particulate matter emissions from layer houses. Flock emptying and replacement will increase the disturbance of hen feathers, although there will be less birds in the house during this period. Cleaning out of manure from the house will obviously result in increased manure disturbance and thus increase particulate matter emissions. Molting is a management activity that intrinsically involves feather loss, so that can also increase

particulate matter emissions, however, it is also a period where reduced or different feed is provided to the birds that could also influence emissions. Accordingly, the influence of management activities was investigated further for particulate matter emissions from layer houses by assigning a status to each day (see section 3.1 for more information on status assignments). Since, there is no manure storage in manure-belt houses, only the flock emptying and replacement, and molting management activities were investigated for this house type. As mentioned previously for the manure shed (section 3.1), the only management event that occurred during the NAEMS monitoring period was the cleanout of manure, which occurred twice for two single days. However, neither of the two manure removal days coincided with particulate matter measurements. House management activities that influence the amount and solid content of manure can also potentially affect particulate matter emissions from the manure shed once the manure arrives 5 days later. Therefore, the influence of management activities was selected for further investigation using the assigned two letter statuses described in section 3.1.

4.0 SITE COMPARISON, TRENDS, AND ANALYSIS

Before developing the EEMs, EPA evaluated NAEMS data for each pollutant to identify patterns and trends in the emissions data using a combination of summary statistics (mean, standard deviation, number of data values, median, minimum, maximum, coefficient of variation, and number of data values less than zero) and time series plots. Section 4.1 summarizes the emissions trends from the sites, while Appendix D contains the tables of summary statistics. Appendix E presents the time series plots of the site-specific emissions, environmental and production parameters, and manure data collected under NAEMS.

Based on the analysis described in Section 3.0, EPA identified the key environmental and manure parameters that potentially affect emissions from egg-layer houses and associated manure sheds. Parameters of particular interest included inventory, average hen weight, live animal weight, hen age, house conditions (exhaust temperature, exhaust relative humidity, and airflow), ambient temperature, ambient relative humidity, manure moisture, manure total ammoniacal nitrogen (TAN), manure pH, and manure total Kjeldahl Nitrogen (TKN). For the manure shed, additional inventory and live animal weight parameters were considered that represented a five day lag to account for the average time is it takes for manure to move to the manure shed.

The next step of the analysis was to look at the key environmental and manure parameters compared to emissions trends. The exploratory data analysis was conducted to confirm that the variables were selected based on the following criteria: (1) data analysis in this study and in the literature suggested that these variables had an influence on emissions; (2) the variables should be easy to measure; and (3) the variables were already in the daily average NAEMS data and were available for most days of monitored emissions. This selection criteria issue particularly applies to the manure parameters, such as moisture content and TAN concentration, which were infrequent due to the intensive collection and analysis methods. Additional time could be taken to develop an appropriate methodology for interpolating between the few data points available for these parameters in the dataset. However, these parameters are difficult to acquire as they require chemical analysis from a laboratory. The exploratory data analysis was also used to explore whether additional parameters, such as hen age, could be included to explain trends. A summary of this analysis for environmental parameters is discussed in Section 4.2, and a summary of the manure parameters is presented in Section 4.3. Again, Appendix D contains summary statistics, Appendix E contains the relevant time series plots, and Appendix F contains least squares regression analysis between the identified parameters and emissions.

4.1 Emissions Data

4.1.1 High Rise Houses

Appendix D, Table D-1 presents the summary statistics for daily average emissions of NH₃ for the high rise sites. Daily averge NH₃ emissions ranged from 32.27 kg d⁻¹ to 248.18 kg d⁻¹. The table indicates the emissions are proportional to inventory. That means the houses with the fewest birds, CA2B, have the lowest average emissions (33.11 and 32.27 kg d⁻¹) and the houses with the largest number of birds, IN2H, have the highest average emissions (224.75 and 248.18 kg d⁻¹). Appendix E, Figure E-1 shows that the emissions can be quite variable at each site, as reiterated by standard deviations that can be as much as half the average value. However, the figure also demonstrates that the houses at the same site are not always in sync with each other, with emissions from one house peaking when the second was much lower. This asynchronous behavior makes it hard to discern any temporal patterns in the data due to seasonal effects. It suggests the emission from the emission from individual houses are also dependent on house specific parameters, such as bird weight and ventilation rate, that are not uniform across all houses at the site. The plot of IN2H suggests a peak in emissions following the new year, but that is not consistent across sites. There were only 4 negative values in the NH₃ dataset, two days for both IN2H H6 and H7.

H₂S summary statistics are presented in Appendix D, Table D-2. Average daily H₂S emissions ranged from 39.96 g d⁻¹at CA2B H6 to 301.54 g d⁻¹ at IN2H H6. Like the NH₃ emissions, the houses with the most birds (IN2H) have the highest emissions. The emissions from the IN2H houses were typically 5 to 7 times higher than the houses at other sites. Appendix E, Figure E-2 shows H₂S emissions are prone to isolated high daily averages. This can be seen in in the NC2B graph in particular as values typically stay below 100 grams per day until the second half of 2008 when a few isolated days jump to emissions greater than 200 grams per day. Appendix E, Figure E-2 also highlights the H₂S dataset had several negative average daily means across all the sites. Negative emission values can result from instrumentation drift between calibrations, instances where concentration measurements are near the minimum detection limit of the instrument, or instrument fluctuations due to ambient conditions (i.e., ambient concentrations greater than the house concentrations). Furthermore, Appendix E, Figure E-2 shows houses at the same site are not always in sync and peaks at one house do not necessarily mean the other house will peak as well. The best example is CA2B in early 2008, when H5 is experiencing near minimum values but H6 is experiencing maximum values. Appendix E, Figure E-2 does suggest that H₂S emissions peak in the summer. Appendix D, Table D-2 provides counts of the number of days with negative emission (N<0), and notes CA2B H6 had the least with 1 day while IN2H H7 had the most with 24.

For PM₁₀, the summary statistics (Appendix D, Table D-3) show the average daily emission ranges from 959.86 g d⁻¹ at CA2B H6 to 4,897.59 g d⁻¹ at IN2H H7. The table indicates the houses with the most birds again have the highest average daily emissions. However, the difference is not as drastic as seen with H₂S or NH₃. With the gaseous pollutants, gaseous emissions from IN2H ranged from 4 to 7 times higher than other houses. For PM₁₀, the IN2H emissions dip slightly, to 2-5 times higher than the other houses. Appendix E, Figure E-3 suggests that average daily emissions see an increase in the summer months, which looks consistent across sites and houses. In Appendix E, Figure E-3 there appears to be less asynchronous behavior between the houses, with the exception of IN2H in mid-2007. Most houses had less than 10 days with negative daily emissions (Appendix D, Table D-3), except for IN2H H6, which had 30 days.

The PM_{2.5} average daily emissions are more consistent across sites; compared to NH₃ and H₂S, which seemed to depend on the number of birds present. The average daily emissions for houses (Appendix D, Table D-4) are typically within a factor of two or less of each other. The exception is NC2B H3 at 144.89 g d⁻¹, which has an average daily emission 7 times lower than the site with the highest average daily emissions, CA2B H5 at 237.47 g d⁻¹. Appendix E, Figure E-4 further supports that PM_{2.5} emissions are generally more consistent, as values from houses at the same site typically have similar concentrations. The sparse temporal nature of the daily values makes it hard to determine if there is a seasonal trend to the data. The number of negative daily averages from the sites varied greatly. Both houses at IN2H had no negative values, while NC2B had 3 and 7 negative values at H3 and H4, respectively. The houses at CA2B had the most negative values with 29 and 25 at H5 and H6, respectively.

The summary statistics for TSP are available in Appendix D, Table D-5. These statistics indicate less of a disparity between houses, with the highest average daily emission value of 7,803.04 g d⁻¹ at site IN2H H6, approximately 2 to 3 times higher than the other houses. Appendix E, Figure E-5 does show some variation between houses at the same site. Similar to PM_{2.5}, the sparse temporal nature of the daily values makes it hard to determine if there is a seasonal trend to the data. The TSP dataset had only two negative values at NC2B H4.

4.1.2 Manure belt Houses

Appendix D, Table D-6 presents the summary statistics for daily average emissions of NH₃ for the manure belt houses. From the table, the average daily emissions are comparable between the two houses with values of 71.54 and 67.05 kg d⁻¹ at IN2B H8 and IN2B H9, respectively. Appendix E, Figure E-6 shows that the daily average emissions can be quite variable at each site, as reiterated by standard deviations that can be as much as half the average value. The figure also demonstrates that these houses are in better sync temporally than the high

rise houses, with peaks of NH₃ emissions occurring at approximately the same time. The plot shows a peak in emissions at the start of 2009, which coincides with the molting phase in both houses.

H₂S summary statistics are available in Appendix D, Table D-7 and again show comparable average daily emissions, with 492.07 and 471.07 gd-1 at IN2B H8 and IN2B H9, respectively. Appendix E, Figure E-7 generally supports this but shows the highest emission at each site occur at different times in the study. IN2B H8 saw its highest H₂S emission over the summer of 2008 while IN2B H9 saw the highest emissions in early 2009. The management of the houses were in sync, with both houses having new flocks placed toward the start of the monitoring period and management phases occurring at similar times. Emissions during these peak periods were comparable and seem within in normal operation. The dissociation of the peak emissions is likely due to a subtle management difference in the houses that was not logged or a house environmental factor, like temperature.

For PM₁₀, the summary statistics (Appendix D, Table D-8) are not as consistent as the gaseous pollutants. Average daily emissions for IN2B H8 were 3,038.68 gd⁻¹, while IN2B H9 had an averge daily emission of 6,076.17 gd⁻¹. Appendix E, Figure E-8 shows the collection of PM₁₀ at IN2B H8 was not as frequent as H9. The site report indicates that the TEOM at IN2B H8 was in repair for these periods due to various failures with the unit (Ni et al., 2010b). One of the repair periods for IN2B H8 occurred while IN2B H9 had high PM₁₀ emissions. It is possible that the absence of similarly high emission at IN2B H8 at this time is the reason for the higher average emissions at IN2B H9 seen in Appendix D, Table D-8. The table also shows more negative observations were recorded for PM₁₀ than with the gaseous pollutants, possibly due to data measurement limitations discussed in 4.1.4"As discussed in Section 2.1 and Appendix B, EPA developed a process to evaluate data that were likely affected by outdoor events, as recommended by the Science Advisory Board review of the initial 2012 models (SAB, 2013). Through this process, erroneous negative data were removed from the final dataset for model development. However, the analysis presented here and in Appendices D through F includes these negative values.

The summary statistics for PM_{2.5} (Appendix D, Table D-9) indicate that data collection for this pollutant proved challenging, as the mean for IN2B H8 is negative and over half of all the readings are negative. Appendix E, Figure E-9 shows that all the observations in the last collection period for IN2B H8 fell below zero. IN2B H9 had fewer negative observations and more observations in total, which yielded a positive average value. Valid negative readings can occur at very low concentrations near the method detection limit. Negative emissions can also occur when ambient concentrations are greater than the house concentrations. In these instances, the calculation to determine the emissions from the house subtract the ambient concentration. If

the concentration in the house is lower, an erroneous negative emission value will result. Section 4.1.4 explores additional concerns about the monitoring set up at IN2H that may contribute to the number of negative values. Similar to the PM_{10} data, EPA processed the $PM_{2.5}$ data further to exclude erroneous data from the final modeling dataset.

The TSP summary statistics (Appendix D, Table D-10) show differences in each house. IN2B H9 has an average daily emission of 21,870.99 gd⁻¹, which is more than twice IN2B H8, as well as a higher standard deviation, which is larger than the mean. Appendix E, Figure E-10 also shows a high variability in the H9 emissions, especially compared to H8.

4.1.3 Manure Shed

For the manure shed, the emissions for all pollutants were less than the emissions in the houses that supplied manure into the shed. The summary statistics for NH₃ (Appendix D, Table D-6) and H₂S (Appendix D, Table D -7) both show large standard deviations (6.52 kg d⁻¹) compared to the average daily mean (4.74 kg d⁻¹), which suggests highly variable emissions at the manure shed. The time series plots for NH₃ (Appendix E, Figure E-11) and H₂S (Appendix E, Figure E-12) shows some periods of more limited variability around the mean, particularly summer of 2009, along with instances of greater variability paired with spikes in emissions across the monitoring period. Notably, both NH₃ and H₂S both experience their maximum values near the same time in early July 2008. This appears to coincide with wetter manure in the shed. The site report (Ni et al., 2010) notes that on June 29, 2008 a manure pile in the shed was visibly wet. As noted in Section 3.1, higher moisture content results in higher emissions.

PM₁₀ (Appendix D, Table D-8; Appendix E, Figure E-13), PM_{2.5} (Appendix D, Table D-9; Appendix E, Figure E-14), and TSP (Appendix D, Table D-10; Appendix E, Figure E-15) observations were also variable, with standard deviations greater than the average across the monitoring period. The time series plots show more day to day variability that the plots for the gaseous species. The variability does not have an obvious pattern but could correlate with a shed environmental factor like temperature, or coincide with movement in the shed, which was not recorded frequently.

4.2 Environmental Parameters

4.2.1 High Rise Houses

The statistical summary of the environmental parameters associated with high rise laying houses are presented in Appendix D, Table D-11. The inventory varies widely across the sites, with CA2B having just over 32,000 birds in each house to IN2H with just over 218,000 birds in each house. Appendix E, Figure E-16 shows that the number of birds present in each house over

the course of NAEMS was fairly consistent. However, each house did have a restocking event during the course of the study, where the existing flock of birds was removed, the house remained empty for at least one day, and then a new flock of birds was placed in the house. Appendix F Figures F-1 through F-5 show the scatter plots of inventory versus each pollutant. A summary of the findings is provided in Table 4-1. In general, there is a positive relationship with inventory across all pollutants, which is consistent with literature.

Hen weight was fairly consistent across the houses with the average hen weight ranging from 1.44 to 1.66 kg. Weight varied the most at CA2B, as the standard deviations were 0.13 and 0.12 kg for H5 and H6, respectively. The remaining houses had standard deviations less than 0.09 kg. Appendix E, Figure E-17 shows this increased variability at CA2B, as compared to the other sites. With the consistent weight over the NAEMS period, the regression analysis (Appendix F, Figures F-6 through F-10, summary in Table 4-1) showed weak correlations between hen weight and all the pollutants.

Combining inventory with hen weight, Live Animal Weight (inventory * hen weight), can be predictive of emissions. With little variation in hen weight across houses in the study, the inventory drives the differences between the houses. That is, the site with the largest inventory (IN2H) has the highest live animal weight. Much like the inventory trend, Appendix E, Figure E-18 shows the live animal weight trend is relatively stable with some variation during the low weight times of hen replacement and molting. The regression analysis (Appendix F Figures F-11 through F-15, summary in Table 4-1) showed modest correlations between live animal weight and each pollutant, which were consistent with the inventory correlations.

EPA derived a hen age variable based on days since placed in the house, for those flocks placed during the study. As expected, the trends for hen age in Appendix E, Figure E-19 show a steady increase over the study. The regression analysis (Appendix F, Figures F-16 through F-20, summary in Table 4-1) only shows weak correlations between hen age and each pollutant. An age was not established for flocks already in the house when the monitoring started, and therefore limited the data available for this parameter.

Table 4-1. Bird specific parameters regression analysis for high rise houses.

Pollutant	Parameter	R ²	Strength	Figure
NH ₃	Inventory	0.6781	moderately strong	Appendix E, E-1
H ₂ S	Inventory	0.3377	modest	Appendix E, E-2
PM_{10}	Inventory	0.2583	modest	Appendix E, E-3
PM _{2.5}	Inventory	0.0008	Slight or weak	Appendix E, E-4
TSP	Inventory	0.2475	modest	Appendix E, E-5
NH ₃	Hen weight	0.1012	Slight or weak	Appendix E, E-6
H ₂ S	Hen weight	0.0489	Slight or weak	Appendix E, E-7
PM ₁₀	Hen weight	0.1969	Slight or weak	Appendix E, E-8
PM _{2.5}	Hen weight	0.1942	Slight or weak	Appendix E, E-9
TSP	Hen weight	0.2080	modest	Appendix E, E-10
NH ₃	Live animal weight	0.6869	moderately strong	Appendix E, E-11
H ₂ S	Live animal weight	0.3328	modest	Appendix E, E-12
PM ₁₀	Live animal weight	0.2269	modest	Appendix E, E-13
PM _{2.5}	Live animal weight	0.0020	Slight or weak	Appendix E, E-14
TSP	Live animal weight	0.2247	modest	Appendix E, E-15
NH ₃	Hen age	0.0253	Slight or weak	Appendix E, E-16
H ₂ S	Hen age	0.0008	Slight or weak	Appendix E, E-17
PM ₁₀	Hen age	0.0244	Slight or weak	Appendix E, E-18
PM _{2.5}	Hen age	0.0279	Slight or weak	Appendix E, E-19
TSP	Hen age	0.0017	Slight or weak	Appendix E, E-20

Appendix D, Table D-11 shows all the houses maintained a similar range of relative humidities across the study, with average daily values ranging from 48.72% at IN2H H6 to 68.40% at NC2B H3. The trends in house relative humidity shown in Appendix E, Figure E-20 appear to have some seasonality, with values increasing over the winter months and then decreasing after the New Year to a low in spring. Values pick up from spring, but there is not a consistent peak in the summer across sites. Relative humidities at the houses are more variable over the summer before settling back into a decline for winter. Values do exceed 70% at CA2B and NC2B, which could limit particulate matter emissions, as noted in Section 3.2. Regression analysis (Appendix F, Figures F-26 through FE-30, summary in Table 4-2) shows a weak relationship with house relative humidity and pollutant emissions.

The mean daily house temperature (Appendix D, Table D-11) is very consistent across the sites, with less than a 2 degree variation. Appendix E, Figure E-21 shows that the temperature is fairly constant for the year, with some lower temperatures coinciding with the time when hens were removed from the house. With the controlled temperatures in the house, the regression analysis (Appendix F Figures F-21 through F-25, summary in Table 4-2) showed only a weak relationship between house temperature and each pollutant.

Unlike house relative humidity and temperature, airflow varied between sites and houses. Table C-12 shows the average daily airflow rate was proportional with the inventory, meaning that the house with higher inventories had high average air flow rates. Appendix E, Figure E-22 shows a strong seasonal pattern at each site, with air flow rate peaking in the summer in an effort to keep the houses within the desired temperature range for the birds. The regression analysis (Appendix F, Figures F-31 through F-35, summary in Table 4-2) only indicates a weak linear relationship between airflow and any of the pollutants.

Table 4-2. House specific parameters regression analysis for high rise houses.

Pollutant	Parameter	R2	Strength	Figure
NH ₃	Exhaust temperature	0.0343	Slight or weak	Appendix E, E-21
H ₂ S	Exhaust temperature	0.0367	Slight or weak	Appendix E, E-22
PM_{10}	Exhaust temperature	0.0278	Slight or weak	Appendix E, E-23
PM _{2.5}	Exhaust temperature	0.4068	moderate	Appendix E, E-24
TSP	Exhaust temperature	0.0143	Slight or weak	Appendix E, E-25
NH ₃	House relative humidity	0.1701	Slight or weak	Appendix E, E-26
H ₂ S	House relative humidity	0.0514	Slight or weak	Appendix E, E-27
PM ₁₀	House relative humidity	0.0852	Slight or weak	Appendix E, E-28
PM _{2.5}	House relative humidity	0.0410	Slight or weak	Appendix E, E-29
TSP	House relative humidity	0.0650	Slight or weak	Appendix E, E-30
NH ₃	Airflow	0.0463	Slight or weak	Appendix E, E-31
H ₂ S	Airflow	0.3029	modest	Appendix E, E-32
PM ₁₀	Airflow	0.1737	Slight or weak	Appendix E, E-33
PM _{2.5}	Airflow	0.1331	Slight or weak	Appendix E, E-34
TSP	Airflow	0.0057	Slight or weak	Appendix E, E-35

The statistical summary of the ambient parameters for the high rise sites is presented in Appendix D, Table D-14. The table shows that while the sites had different mean ambient relative humidities, they were subject to the same range of values across the study. Appendix E, Figure E-23 shows some seasonality to the measurements, but these patterns vary between the sites. CA2B and IN2H have peaks at the start of the year, with lows midyear. The values for NC2B are more scattered, with high values occurring all year. If we look at the minimum values (i.e., the bottom edge of the scatter), it appears as though the lowest values occur at the start of the year with low values less likely for the summer months. The regression analysis (Appendix F Figures F-41 through F-45, summarized in Table 4-3) showed ambient relative humidity correlation with each pollutant emissions were comparable to the relationship with the house relative humidity.

For ambient temperature, the average daily temperature is lowest at IN2H followed by NC2B, and CA2B. The sites did have variation in the range of temperatures covered, as CA2B was not exposed to freezing temperatures, but IN2H and NC2B were. The temporal trend in

ambient temperature is as expected, with Appendix E, Figure E-24 showing peaks in the July timeframe and lows after the new year. The regression analysis (Appendix F Figures F-36 through F-40, summarized in Table 4-3) showed ambient temperature had a similar relationship to pollutant emissions as house temperature.

Table 4-3. Ambient parameters regression analysis for high rise houses.

Pollutant	Parameter	R ²	Strength	Figure
NH ₃	Ambient temperature	0.1192	Slight or weak	Appendix E, E-36
H ₂ S	Ambient temperature	0.0341	Slight or weak	Appendix E, E-37
PM ₁₀	Ambient temperature	0.0165	Slight or weak	Appendix E, E-38
$PM_{2.5}$	Ambient temperature	0.3637	modest	Appendix E, E-39
TSP	Ambient temperature	0.0959	Slight or weak	Appendix E, E-40
NH₃	Ambient relative humidity	0.0684	Slight or weak	Appendix E, E-41
H ₂ S	Ambient relative humidity	0.0088	Slight or weak	Appendix E, E-42
PM ₁₀	Ambient relative humidity	0.0018	Slight or weak	Appendix E, E-43
PM _{2.5}	Ambient relative humidity	0.2103	modest	Appendix E, E-44
TSP	Ambient relative humidity	0.0051	Slight or weak	Appendix E, E-45

4.2.2 Manure Belt Houses

The statistical summary of the environmental parameters associated with manure belt houses are presented in Appendix D, Table D-12. Time series plots of each parameter are available in Appendix E, Figures E-25 through E-33.

The inventory was fairly consistent between the two houses, with a population of approximately 250,000 birds in each house. Appendix E, Figure E-25 shows that the number of birds present over the course of NAEMS was fairly consistent. The exception is the restocking event at IN2B H9, where the existing flock of birds were removed, the house remained empty for at least one day, and then a new flock of birds was placed in the house. The flock replacement for IN2B H8 occurred just before the start of the study. Appendix F Figures F-63 through F-67, summarized in Table 4-4, show the scatter plots of inventory versus each pollutant. In general, there is a weak positive relationship with inventory across all pollutants, which is consistent with literature. The relationship is weaker than what was seen for high rise houses as the manure is being removed from the house daily, as opposed to accumulating over an extended period of time.

Hen weight is also consistent between the two houses, with a typical weight of 1.4 kg. Appendix E, Figure E-26 shows that hen weight over the course of NAEMS was fairly consistent, except for the dip during the molting process, which occurred midway through the study. Appendix F, Figures F-68 through F-72 show the scatter plots of hen weight versus each pollutant. In general, there is a slight or weak positive relationship with hen weight across all

pollutants. Similar to inventory and hen weight, live animal weight is consistent between the houses (Table C-13). Live animal weight follows a similar trend to hen weight, which is fairly constant except for a dip during the molting period (Appendix E, Figure E-27). Appendix F, Figures F-73 through F-67 are regression plots, summarized in Table 4-4, which suggest a weak positive relationship between live animal weight and each pollutant.

Hen age was also explored (Appendix E, Figure E-28) as a parameter. Appendix F Figures F-78 through F-82, summarized in Table 4-4, show the scatter plots of hen age versus each pollutant. In general, there is a weak negative relationship across all pollutants.

Table 4-4. Bird specific parameters regression analysis for manure belt houses.

Pollutant	Parameter	R ²	Strength	Figure
H ₂ S	Inventory	0.0870	Slight or weak	Appendix E, E-64
NH₃	Inventory	0.0524	Slight or weak	Appendix E, E-63
PM ₁₀	Inventory	0.0241	Slight or weak	Appendix E, E-65
PM _{2.5}	Inventory	0.3679	modest	Appendix E, E-66
TSP	Inventory	0.0012	Slight or weak	Appendix E, E-67
H ₂ S	Hen weight	0.0161	Slight or weak	Appendix E, E-69
NH₃	Hen weight	0.0461	Slight or weak	Appendix E, E-68
PM ₁₀	Hen weight	0.0009	Slight or weak	Appendix E, E-70
PM _{2.5}	Hen weight	0.1552	Slight or weak	Appendix E, E-71
TSP	Hen weight	0.0087	Slight or weak	Appendix E, E-72
H ₂ S	Live animal weight	0.0819	Slight or weak	Appendix E, E-74
NH₃	Live animal weight	0.0833	Slight or weak	Appendix E, E-73
PM ₁₀	Live animal weight	0.0207	Slight or weak	Appendix E, E-75
PM _{2.5}	Live animal weight	0.0357	Slight or weak	Appendix E, E-76
TSP	Live animal weight	0.0096	Slight or weak	Appendix E, E-77
H ₂ S	Hen age	0.1609	Slight or weak	Appendix E, E-79
NH ₃	Hen age	0.0405	Slight or weak	Appendix E, E-78
PM ₁₀	Hen age	0.1006	Slight or weak	Appendix E, E-80
PM _{2.5}	Hen age	0.0593	Slight or weak	Appendix E, E-81
TSP	Hen age	0.0626	Slight or weak	Appendix E, E-82

House relative humidity was slightly higher in IN2B H9 for most of the study. The summary statistics for relative humidity are 3-6% higher in IN2B H9 than IN2B H8 and the time series in Appendix E, Figure E-29 is higher for IN2B H9 than IN2B H8 almost every day. The other house parameters, temperature (Appendix E, Figure E-30) and airflow (Appendix E, Figure E-31), were fairly consistent between the houses. The only exception was a drop in temperature in IN2B H9 while the flock was being replaced. The regression analysis of house relative humidity (Appendix F, Figures F-88 through F-92, summarized in Table 4-5) showed a weak positive relationship for the gaseous species and a slight or weak negative relationship for PM₁₀, which was consistent with expectations. The plots suggested PM_{2.5} and house relative humidity

had a neutral/no relationship and TSP had a weak positive relationship. The limited number of observations for each pollutant are the likely cause for the inconsistent relationship with PM₁₀. Both the house exhaust temperature (Appendix F, Figures F-83 through F-87, summarized in Table 4-5) and airflow (Appendix F, Figures F-93 through F-97, summarized in Table 4-5) regressions showed slight linear relationships with the pollutants.

The statistical summary of the ambient parameters for the manure belt site is presented in Appendix D, Table D-12. Both the ambient relative humidity (Appendix E, Figure E-32) and temperature (Appendix E, Figure E-33) follow the typical seasonal patterns expected in Indiana. The linear regression analysis plots of ambient parameters (Appendix F, Figure F-98 through F-107, summarized in Table 4-6) show weak relationships that are not consistent across the gaseous pollutants or particulate matter species.

Table 4-5. House specific parameters regression analysis for manure belt houses.

Pollutant	Parameter	R ²	Strength	Figure
H ₂ S	Exhaust temperature	0.0716	Slight or weak	Appendix E, E-84
NH ₃	Exhaust temperature	0.00004	Slight or weak	Appendix E, E-83
PM ₁₀	Exhaust temperature	0.0033	Slight or weak	Appendix E, E-85
PM _{2.5}	Exhaust temperature	0.1947	Slight or weak	Appendix E, E-86
TSP	Exhaust temperature	0.0201	Slight or weak	Appendix E, E-87
H ₂ S	House relative humidity	0.0398	Slight or weak	Appendix E, E-89
NH₃	House relative humidity	0.1474	Slight or weak	Appendix E, E-88
PM ₁₀	House relative humidity	0.0031	Slight or weak	Appendix E, E-90
PM _{2.5}	House relative humidity	0.0006	Slight or weak	Appendix E, E-91
TSP	House relative humidity	0.0302	Slight or weak	Appendix E, E-92
H ₂ S	Airflow	0.0450	Slight or weak	Appendix E, E-94
NH₃	Airflow	0.0078	Slight or weak	Appendix E, E-93
PM ₁₀	Airflow	0.0016	Slight or weak	Appendix E, E-95
PM _{2.5}	Airflow	0.2047	Modest	Appendix E, E-96
TSP	Airflow	0.1518	Slight or weak	Appendix E, E-97

Table 4-6. Ambient parameter regression analysis for manure belt houses.

Pollutant	Parameter	R ²	Strength	Figure
H ₂ S	Ambient relative humidity	0.0010	Slight or weak	Appendix E, E-104
NH ₃	Ambient relative humidity	0.0883	Slight or weak	Appendix E, E-103
PM ₁₀	Ambient relative humidity	0.0008	Slight or weak	Appendix E, E-105
PM _{2.5}	Ambient relative humidity	0.0863	Slight or weak	Appendix E, E-106
TSP	Ambient relative humidity	0.0242	Slight or weak	Appendix E, E-107
H ₂ S	Ambient temperature	0.0162	Slight or weak	Appendix E, E-99
NH ₃	Ambient temperature	0.0964	Slight or weak	Appendix E, E-98
PM ₁₀	Ambient temperature	0.0010	Slight or weak	Appendix E, E-100
PM _{2.5}	Ambient temperature	0.1353	Slight or weak	Appendix E, E-101
TSP	Ambient temperature	0.0733	Slight or weak	Appendix E, E-102

4.2.3 Manure Shed

The statistical summary of the environmental parameters associated with the monitored manure shed are presented in Appendix D, Table D-12. For inventory (Appendix E, Figure E-34) and live animal weight (Appendix E, Figure E-37) the combined value from both houses were examined, as well as a version that represented a five-day lag from the emissions (Appendix E, Figures E-35 and E-38). Due to the scale of the plots, the five-day shift is indistinguishable. The summary statistics and plots for the inventory show it was relatively constant for the study, except when the flock was replaced at IN2B H9 in late 2008. Plots for live animal weight are similar, except there is a little more variation across the year. Appendix F, Figures F-120 through F-124 (summarized in Table 4-7) show the scatter plots of inventory, both lagged and not, versus each pollutant. Appendix F, Figures F-130 through F-134 show the scatter plots of live animal weight, both lagged and not, versus each pollutant. Table 4-7 summarizes the results for both sets of parameters. The analysis shows weak to modest linear relationship with inventory and the emission of each pollutant.

Average hen weight saw fluctuations over the study period as new birds were added and during the molting phase (Appendix E, Figure E-36), which contributed to most of the variability in live animal weight. Appendix F, Figures F-125 through F-129, summarized in Table 4-7, show the scatter plots of average hen weight versus each pollutant, which showed a slight linear relationship. Average hen age (Appendix E, Figure E-39) was also examined. The value represented the average age across the two houses contributing to the manure shed. Appendix F, Figures F-135 through F-139 show the scatter plots of average hen age versus each pollutant, which showed a slight linear relationship for all pollutants except PM_{2.5}, which had a moderate negative linear relationship.

Table 4-7. Bird specific parameters regression analysis for manure sheds.

Pollutant	Parameter	R ²	Strength	Figure
NH ₃	Inventory	0.0152	Slight or weak	Appendix E, E-120
H ₂ S	Inventory	0.0158	Slight or weak	Appendix E, E-121
PM ₁₀	Inventory	0.3074	modest	Appendix E, E-122
PM _{2.5}	Inventory	0.3074	modest	Appendix E, E-123
TSP	Inventory	0.0216	Slight or weak	Appendix E, E-124
NH ₃	Inventory, 5 day lag	0.0147	Slight or weak	Appendix E, E-120
H ₂ S	Inventory, 5 day lag	0.0120	Slight or weak	Appendix E, E-121
PM ₁₀	Inventory, 5 day lag	0.3074	modest	Appendix E, E-122
PM _{2.5}	Inventory, 5 day lag	0.3074	modest	Appendix E, E-123
TSP	Inventory, 5 day lag	0.0206	Slight or weak	Appendix E, E-124
NH ₃	Hen weight	0.0003	Slight or weak	Appendix E, E-125
H ₂ S	Hen weight	0.0278	Slight or weak	Appendix E, E-126
PM ₁₀	Hen weight	0.0214	Slight or weak	Appendix E, E-127
PM _{2.5}	Hen weight	0.1815	Slight or weak	Appendix E, E-128
TSP	Hen weight	0.0404	Slight or weak	Appendix E, E-129
NH₃	Live animal weight	0.0026	Slight or weak	Appendix E, E-130
H ₂ S	Live animal weight	0.0014	Slight or weak	Appendix E, E-131
PM ₁₀	Live animal weight	0.1730	Slight or weak	Appendix E, E-132
PM _{2.5}	Live animal weight	0.2852	modest	Appendix E, E-133
TSP	Live animal weight	0.0838	Slight or weak	Appendix E, E-134
NH ₃	Live animal weight, 5 day lag	0.0000	Slight or weak	Appendix E, E-130
H ₂ S	Live animal weight, 5 day lag	0.0102	Slight or weak	Appendix E, E-131
PM ₁₀	Live animal weight, 5 day lag	0.0164	Slight or weak	Appendix E, E-132
PM _{2.5}	Live animal weight, 5 day lag	0.2464	modest	Appendix E, E-133
TSP	Live animal weight, 5 day lag	0.0954	Slight or weak	Appendix E, E-134
NH₃	Hen age	0.0206	Slight or weak	Appendix E, E-135
H ₂ S	Hen age	0.0530	Slight or weak	Appendix E, E-136
PM ₁₀	Hen age	0.0006	Slight or weak	Appendix E, E-137
PM _{2.5}	Hen age	0.4424	moderate	Appendix E, E-138
TSP	Hen age	0.0717	Slight or weak	Appendix E, E-139

Reviewing the typical house parameters, the manure shed was naturally ventilated and maintained a temperature and humidity that approximated the ambient conditions. Since the manure shed is naturally ventilated, the airflow was estimated using wind velocity measurements from five 2-D sonic anemometers and two impeller anemometers (Ni et al. 2010b). Since the airflow (Appendix E, Figure E-40) is based on ambient wind flow through the house and is not dependent on temperature, it does not follow any seasonal trends. The scatter plots (Appendix F, Figures F-140 through F-144, summarized in Table 4-8) show at least a modest positive relationship with gaseous emissions and a weak positive relationship with particulate matter.

The statistical summary of the ambient parameters for the manure shed is presented in Appendix D, Table D-12. Both the ambient relative humidity (Appendix E, Figure E-41) and temperature (Appendix E, Figure E-42) follow the typical seasonal patterns expected in Indiana. The linear regression analysis plots (Appendix F, Figures F-145 through F-154, summarized in Table 4-8) show weak relationships that are not consistent across the gaseous pollutants or particulate matter species.

The wind speed was examined for manure shed, as it may be related to airflow through the shed. Wind speeds (Appendix D, Table D-12) ranged from 0.04 to 8.54 ms⁻¹ over the monitoring period, with an average of 0.73 ms⁻¹. Appendix E, Figure E-43 reiterates that most wind speeds are below 1 ms⁻¹, with occasional spikes likely related to synoptic events. The linear regression analysis plots (Appendix F, Figure F-155 through F-159, summarized in Table 4-8) show weak positive relationships across all pollutants.

Table 4-8. House and ambient parameter regression analysis for manure sheds.

Pollutant	Parameter	R ²	Strength	Figure
NH ₃	Airflow	0.4745	moderate	Appendix E, E-140
H ₂ S	Airflow	0.3917	modest	Appendix E, E-141
PM ₁₀	Airflow	0.1877	Slight or weak	Appendix E, E-142
PM _{2.5}	Airflow	0.6428	moderately strong	Appendix E, E-143
TSP	Airflow	0.0689	Slight or weak	Appendix E, E-144
NH₃	Ambient temperature	0.0328	Slight or weak	Appendix E, E-145
H ₂ S	Ambient temperature	0.0813	Slight or weak	Appendix E, E-146
PM ₁₀	Ambient temperature	0.0030	Slight or weak	Appendix E, E-147
PM _{2.5}	Ambient temperature	0.0040	Slight or weak	Appendix E, E-148
TSP	Ambient temperature		Slight or weak	Appendix E, E-149
NH₃	Ambient relative humidity	0.0057	Slight or weak	Appendix E, E-150
H ₂ S	Ambient relative humidity	0.0200	Slight or weak	Appendix E, E-151
PM ₁₀	Ambient relative humidity	0.0065	Slight or weak	Appendix E, E-152
PM _{2.5}	Ambient relative humidity	0.1697	Slight or weak	Appendix E, E-153
TSP	Ambient relative humidity	0.0066	Slight or weak	Appendix E, E-154
NH₃	Wind Speed	0.0017	Slight or weak	Appendix E, E-155
H ₂ S	Wind Speed	0.0022	Slight or weak	Appendix E, E-156
PM ₁₀	Wind Speed	0.0058	Slight or weak	Appendix E, E-157
PM _{2.5}	Wind Speed	0.0086	Slight or weak	Appendix E, E-158
TSP	Wind Speed	0.0001	Slight or weak	Appendix E, E-159

4.3 Manure Parameters

4.3.1 High Rise Houses

Appendix D, Table D-14 summarizes manure parameters for the high rise houses. For manure age, as determined from reports of house clean outs, the time the manure was left in the

house varied between sites and the houses at each site. CA2B cleaned out the stored manure more frequently than NC2B or IN2H. Appendix E, Figure E-44 shows the episodic cleaning at CA2B (seven times), and less frequent clean outs at IN2H (one time) and NC2B (three time). Appendix F Figures F-46 through F-50 show the scatter plots of manure age versus each pollutant, which are summarized in Table 4-9. The analysis shows only a weak linear relationship with manure age and the emission of each pollutant.

The average pH at the sites (Appendix D, Table D-15) ranged from 7.60 (CA2B H6) to 8.50 (IN2H H7). All readings fell within the range of 6.40 (CA2B H6) to 8.79 (IN2H H7). When plotted (Appendix E, Figure E-45), the sparse nature of the measurements makes it difficult to discern any trends. The regression analysis (Appendix F Figures F-51 through F-54) showed pH had a weak linear relationship to the emission of NH₃, H₂S, and PM₁₀. For PM_{2.5} and TSP emissions, there was not sufficient measurement data to conduct a linear regression analysis.

For the percent solids composition, average values were higher at CA2B, with IN2H and NC2B having slightly lower values. Again, the sparse nature of the readings makes it difficult to discern any trends or consistent temporal patterns in the measurements (Appendix E, Figure E-46). The regression analysis (Appendix F Figures F-55 through F-58) showed percent solids composition had a weak linear relationship to the emission of NH₃, H₂S, and PM₁₀. Again, there were insufficient measurement data to conduct a regression analysis for PM_{2.5} and TSP emissions.

For the percent total ammoniacal nitrogen (TAN), no sites had observations above 1%. IN2H saw the highest values followed by NC2B and CA2B. As with the other manure parameters, the sparse nature of the readings makes it difficult to discern any trends or consistent temporal patterns in the measurements (Appendix E, Figure E-47). The regression analysis (Appendix F Figures F-59 through F-62) showed TAN had a weak linear relationship to the emission of NH₃, H₂S, and PM₁₀. Again, there were insufficient measurement data to conduct a regression analysis for PM_{2.5} and TSP emissions.

Table 4-9. Manure parameter regression analysis for high rise houses.

Pollutant	Parameter	R ²	Strength	Figure
NH ₃	Manure age	0.1874	Slight or weak	Appendix E, E-46
H ₂ S	Manure age	0.0037	Slight or weak	Appendix E, E-47
PM_{10}	Manure age	0.1625	Slight or weak	Appendix E, E-48
PM _{2.5}	Manure age	0.1126	Slight or weak	Appendix E, E-49
TSP	Manure age	0.0616	Slight or weak	Appendix E, E-50
NH ₃	рН	0.2607	modest	Appendix E, E-51
H ₂ S	рН	0.0963	Slight or weak	Appendix E, E-52
PM ₁₀	рН	0.1496	Slight or weak	Appendix E, E-53
PM _{2.5}	рН		а	
TSP	рН		а	Appendix E, E-54
NH ₃	Solids	0.1597	Slight or weak	Appendix E, E-55
H ₂ S	Solids	0.0114	Slight or weak	Appendix E, E-56
PM ₁₀	Solids	0.1154	Slight or weak	Appendix E, E-57
PM _{2.5}	Solids		a	
TSP	Solids		a	Appendix E, E-58
NH ₃	TAN	0.4703	moderate	Appendix E, E-59
H ₂ S	TAN	0.0796	Slight or weak	Appendix E, E-60
PM ₁₀	TAN	0.1273	Slight or weak	Appendix E, E-61
PM _{2.5}	TAN		a	
TSP	TAN		а	Appendix E, E-62

^a EPA did not have sufficient measurement data from NAEMS to conduct a linear regression analysis

4.3.2 Manure belt Houses

Appendix D, Table D-15 summarizes manure parameters for the manure belt houses. For pH, the average at the site (Appendix D, Table D-16) ranged from a minimum of 7.08 (H9) to a maximum of 8.53 (H9). The average pH was slightly higher at H8 (7.98) than for H9 (7.77). When plotted (Appendix E, Figure E-48), the sparse nature of the measurements makes it difficult to discern any seasonal trends. The regression analysis (Appendix F, Figures F-108 through F-111), , summarized in Table 4-10, showed pH had a weak linear relationship to the emission of NH₃, H₂S, PM₁₀, and TSP. For PM_{2.5}, there was not sufficient measurement data to conduct a linear regression analysis as the manure collection events did not coincide with any PM_{2.5} monitoring days.

For the percent solids composition, the values between the two houses were relatively consistent. Again, the sparse nature of the readings makes it difficult to discern any trends or consistent temporal patterns in the measurements (Appendix E, Figure E-49). The regression analysis (Appendix F, Figures F-112 through F-115, summary in Table 4-10) showed percent solids composition had a weak linear relationship to the emission of NH₃, H₂S, and TSP. There was a modest relationship with PM₁₀. Again, there were insufficient measurement data to

conduct a regression analysis for $PM_{2.5}$ emissions since the manure sampling did not coincide with days with $PM_{2.5}$ emission measurements.

For the percent total ammoniacal nitrogen (TAN), no sites had observations above 1%. As with the other manure parameters, the sparse nature of the readings makes it difficult to discern any trends or consistent temporal patterns in the measurements (Appendix E, Figure E-50). The regression analysis (Appendix F, Figures F-116 through F-119, summary in Table 4-10) showed TAN had a weak linear relationship to the emission of NH₃, H₂S, and PM₁₀. There was a modest relationship with TSP. Again, there were insufficient measurement data to conduct a regression analysis for PM_{2.5} emissions.

Table 4-10. Manure parameter regression analysis for manure belt houses.

Pollutant	Parameter	R ²	Strength	Figure
H ₂ S	рН	0.0783	Slight or weak	Appendix E, E-109
NH₃	рН	0.0231	Slight or weak	Appendix E, E-108
PM ₁₀	рН	0.1230	Slight or weak	Appendix E, E-110
PM _{2.5}	рН		а	
TSP	рН	0.02	Slight or weak	Appendix E, E-111
H ₂ S	Solids	0.0122	Slight or weak	Appendix E, E-113
NH ₃	Solids	0.1110	Slight or weak	Appendix E, E-112
PM ₁₀	Solids	0.2291	modest	Appendix E, E-114
PM _{2.5}	Solids		а	
TSP	Solids	0.057	Slight or weak	Appendix E, E-115
H ₂ S	TAN	0.1739	Slight or weak	Appendix E, E-117
NH ₃	TAN	0.0115	Slight or weak	Appendix E, E-116
PM ₁₀	TAN	0.0119	Slight or weak	Appendix E, E-118
PM _{2.5}	TAN		а	
TSP	TAN	0.219	modest	Appendix E, E-119

^a EPA did not have sufficient measurement data from NAEMS to conduct a linear regression analysis.

4.3.3 Manure Shed

Appendix D, Table D-15 summarizes manure parameters for the manure shed. Compared to the manure samples taken from the belt house, the manure shed samples have similar pH values. The manure shed samples have a higher solids content and a lower TAN, both of which are due to the manure drying process which reduces moisture.

For the manure shed, manure age was determined based on the clean out dates reported in the site report. This is an approximate age, as the contents of the shed may not have been completely emptied at each removal event. Appendix E, Figure E-51 shows one removal event reported on June 5, 2009. Appendix F Figures F-160 through F-164, summarized in Table 4-11, show the scatter plots of manure age versus each pollutant. The analysis shows only a weak

positive linear relationship with manure age and the gaseous pollutants, and weak negative or neutral relationship with the particulate matter species.

The pH values for the manure shed ranged from 7.31 to 8.76 (Appendix D, Table D-15). The percent solids composition for the manure shed ranged from 49.90 to 86.05%, TAN values shed ranged from 0.26 % to 0.80%, and the percent TKN values ranged from 2.38 to 4.69% (Appendix D, Table D-16). The infrequent manure sample collection results in sparse scatter plots of the manure parameters (Appendix E, Figure E-52 through E-55), and makes it difficult to draw definitive conclusions about any seasonal trends in the parameters.

There was not sufficient measurement data to conduct a linear regression analysis between PM_{2.5} or TSP and the manure parameters. The regression analysis with pH (Appendix F, Figures F-165 through F-167, summary in Table 4-11) showed a moderate positive linear relationship to the emission of NH₃, and H₂S, and only a weak relationship with PM₁₀. Both solids composition (Appendix F, Figures F-168 through F-170) and TAN (Appendix F, Figures F-171 through F-173), showed a slight linear relationship to the emission of NH₃, H₂S. For PM₁₀, solids composition had a weak relationship. However, there was a strong positive relationship between TAN and PM₁₀. The strong relationship with PM₁₀ is based on only three observations and thus should not be taken as a definitive relationship. TKN (Appendix F, Figures F-174 through F-176, summary in Table 4-11) had a moderate positive linear relationship to NH₃ and a weak positive relationship with H₂S. TKN also showed a moderately strong negative relationship with PM₁₀., which was again only based on three observations and should not be considered definitive.

Table 4-11. Manure parameter regression analysis for manure belt houses.

Pollutant	Parameter	R ²	Figure	Strength
NH ₃	Manure age	0.0277	Slight or weak	Appendix E, E-160
H ₂ S	Manure age	0.0251	Slight or weak	Appendix E, E-161
PM ₁₀	Manure age	0.0010	Slight or weak	Appendix E, E-162
PM _{2.5}	Manure age	0.0080	Slight or weak	Appendix E, E-163
TSP	Manure age	0.0904	Slight or weak	Appendix E, E-164
NH ₃	рН	0.4836	moderate	Appendix E, E-165
H ₂ S	рН	0.7927	moderately strong	Appendix E, E-166
PM ₁₀	рН	0.0869	Slight or weak	Appendix E, E-167
PM _{2.5}	рН		a	
TSP	рН		a	
NH ₃	Solids	0.0066	Slight or weak	Appendix E, E-168
H ₂ S	Solids	0.0406	Slight or weak	Appendix E, E-169
PM ₁₀	Solids	0.1666	Slight or weak	Appendix E, E-170
PM _{2.5}	Solids		a	
TSP	Solids		a	
NH ₃	TAN	0.1008	Slight or weak	Appendix E, E-171
H ₂ S	TAN	0.0067	Slight or weak	Appendix E, E-172
PM ₁₀	TAN	0.9321	strong	Appendix E, E-173
PM _{2.5}	TAN		a	
TSP	TAN		a	
NH ₃	TKN	0.2271	modest	Appendix E, E-174
H ₂ S	TKN	0.1150	Slight or weak	Appendix E, E-175
PM ₁₀	TKN	0.7744	moderately strong	Appendix E, E-176
PM _{2.5}	TKN		a	
TSP	TKN		a	

^a EPA did not have sufficient measurement data from NAEMS to conduct a linear regression analysis.

5.0 DEVELOPMENT AND SELECTION OF MODELS FOR DAILY EMISSIONS

5.1 High Rise Operations

The literature review (Section 3) and exploratory data analysis (Section 4) suggested that EPA should consider ambient temperature, exhaust temperature, ambient relative humidity, exhaust relative humidity, management phase, manure age, hen age, inventory, and live animal weight in the development of the emission models for the layer high rise houses. House airflow, or ventilation rate, can have a substantial influence on the emission rate of gaseous pollutants, but was not included in the parameter list as it may not be easily obtained at all farms. Since ventilation rate is essentially driven by the temperature (i.e., the higher ambient temperature the higher the ventilation rate), the ambient temperature provides an indication of airflow in the models tested.

The various combinations of these parameters were used in test models. For NH₃ and H₂S, 10 different combinations were tested as potential models (Table 5-1). There were 15 models (Table 5-2) tested for particulate matter emissions, which had more variations using the relative humidity parameters.

Models G-6, G-10, P-13, and P-14 are slightly different due to the inclusion of the management phase categories as a parameter. These models are useful as they can investigate the potential effect of management activities on emissions, However, there is limited data for some of the management statuses (e.g., transition). EPA considers these models as experimental since an appropriate methodology for their evaluation and application has not been finalized. The models have been included in the tables to note all the options EPA explored, but were not considered as potential models at this time.

Table 5-1. Parameter combinations tested as models for NH₃ and H₂S emissions.

Model	Parameters
G-1	Intercept, Live animal weight, Ambient temperature
G-2	Intercept, Live animal weight, Exhaust temperature
G-3	Intercept, Inventory, Ambient temperature
G-4	Intercept, Inventory, Exhaust temperature
G-5	Intercept, Inventory, Hen age, Ambient temperature
G-6	Intercept, Inventory, Ambient temperature, Management phase (manure cleanouts (C),
G-0	flock emptying and replacement (E), full flock (F), molting (M), and transition (T))
G-7	Intercept, Inventory, Manure age, Ambient temperature
G-8	Intercept, Inventory, Ambient temperature, Ambient relative humidity
G-9	Intercept, Inventory, Ambient temperature, Exhaust relative humidity
G-10	Intercept, Inventory, Ambient temperature, Ambient relative humidity, Management phase
G-10	(C,E,F,M,T)

Table 5-2. Parameter combinations tested as models for PM₁₀, PM_{2.5}, and TSP emissions.

Model	Parameters
P-1	Intercept, Inventory
P-2	Intercept, Inventory, Ambient relative humidity
P-3	Intercept, Inventory, Exhaust relative humidity
P-4	Intercept, Inventory, Ambient relative humidity, Ambient temperature
P-5	Intercept, Inventory, Ambient temperature, Exhaust relative humidity
P-6	Intercept, Inventory, Ambient relative humidity, Exhaust temperature
P-7	Intercept, Inventory, Exhaust temperature, Exhaust relative humidity
P-8	Intercept, Live animal weight, Ambient temperature, Ambient relative humidity
P-9	Intercept, Live animal weight, Ambient temperature, Exhaust relative humidity
P-10	Intercept, Live animal weight, Ambient relative humidity, Exhaust temperature
P-11	Intercept, Live animal weight, Exhaust temperature, Exhaust relative humidity
P-12	Intercept, Hen age, Inventory, Ambient relative humidity
P-13	Intercept, Inventory, Ambient relative humidity, Management phase (C,E,F,M,T)
P-14	Intercept, Inventory, Ambient temperature, Ambient relative humidity, Management phase
P-14	(C,E,F,M,T)
P-15	Intercept, Inventory, Manure age, Ambient relative humidity

For both NH₃ (Appendix G, Table G-1) and H₂S (Appendix G, Table G-3), models G-5, G-6, and G-10 had terms that were not statistically significant (p > 0.05) and were removed from further consideration. The model fit (-2 log likelihood, AIC, AICc, and BIC) and evaluation statistics (ME, NME, MB, NMB) for NH₃ (Appendix G, Table G-2) and H₂S (Appendix G, Table G-4) indicate the remaining models had comparable performance, which suggested using ambient parameters was as effective as models that included house specific parameters. As noted in the main report, the model selection process also looked at how easily obtainable the parameters are as not to create an undue burden on the operators. Generally, ambient parameters were preferred since ambient meteorological data is actively recorded across the country and representative site data is accessible through the NCEI website. To further easy any burden, the EPA plans to provide a tool that automatically populates relevant ambient parameters for any given location instead of requiring producers to measure and record environmental parameters either inside or outside of the house to further reduce the burden of use on the producer.

Therefore, considering ambient temperature is a suitable proxy for house airflow and representative ambient temperature data is accessible, the EPA concluded that a model using ambient temperature and relative humidity would be preferable to one with exhaust temperature and relative humidity. Of the remaining models that used ambient parameters (G-1, G-3, G-7, and G-8), EPA selected model G-8 (including the parameters: intercept, inventory, ambient temperature, ambient relative humidity) for further analysis for both NH₃ and H₂S as it had the

best normalized mean bias of the remaining models. The final form of these models is presented in Table 5-3.

For PM₁₀ (Appendix G, Table G-5), models P-12, P-13, and P-14 had terms that were not statistically significant (p > 0.05) and were removed from further consideration. The model fit and evaluation statistics for PM₁₀ (Appendix G, Table G-6) indicate the remaining models were comparable, which suggested using ambient parameters was as effective as house parameters. Therefore, EPA considered the potential ease of data collection and concluded that ambient temperature and relative humidity would be preferable to one with exhaust temperature and relative humidity. Of the remaining models that used ambient parameters (P-1, P-3, P-4, P-8, and P-15), EPA selected model P-4 (including the parameters: intercept, inventory, ambient relative humidity, ambient temperature) for further analysis as it had the lowest mean error and one of the lowest normalized mean bias of the remaining models. The full form of the model is presented in Table 5-3.

As noted in Section 6.4 of the main report, the particulate matter model selection starts with the PM₁₀ models, as there are more emissions data. The PM₁₀ version of the models had between 1,579 and 2,712 records available depending on the completeness of the various predictive parameters. For PM_{2.5} and TSP the number of records available ranged between 77 – 160 for PM_{2.5} and 149 - 238 for TSP. This is substantially less data that was available for PM₁₀ and does not cover the breadth of conditions that the PM₁₀ data does. Therefore, the models generated with these smaller datasets were examined mainly for consistency with the PM₁₀ results to build confidence in using the same model form for all the particulate matter species. The biggest difference from PM₁₀ is more of the models have insignificant terms for both PM_{2.5} and TSP. For PM_{2.5} (Appendix G, Table G-7) only four models are comprised of significant parameters, and TSP (Appendix G, Table G-9) has only three significant models. Despite the insignificance of the parameters for most of the models, the relationships were consistent with the PM₁₀ models and literature. The model performance statistics for PM_{2.5} (Appendix G, Table G-8) and TSP (Appendix G, Table G-10) were fairly consistent, except for mean bias. Model P-4 had reasonable performance for both PM_{2.5} and TSP and would be consistent with the PM₁₀ formulation that was developed from a much larger dataset. Therefore, EPA selected model P-4 for both PM_{2.5} and TSP to conduct further evaluation and analysis as an emission estimation method. The full forms of the models are presented in Table 5-3.

Table 5-3. Selected daily models for high rise layer houses.

Pollutant	Formula	Equation Number
NH₃	$ln(NH_3) = 2.6598 + 0.0059 * Inventory + 0.0387 * Amb_T + 0.0018 * Amb_{RH}$	Equation 1
H₂S	$ln(H_2S) = 2.7231 + 0.0098 * Inventory + 0.0210 * Amb_T + 0.0038 * Amb_{RH}$	Equation 2
PM ₁₀	$ln(PM_{10}) = 6.8702 + 0.0077 * Inventory + 0.0145 * Amb_T - 0.0030 * Amb_{RH}$	Equation 3
PM _{2.5}	$ln(PM_{2.5}) = 4.6219 + 0.0080 * Inventory + 0.0510 * Amb_T - 0.0181 * Amb_{RH}$	Equation 4
TSP	$ln(TSP) = 7.5995 + 0.0079 * Inventory + 0.0137 * Amb_T - 0.0058 * Amb_{RH}$	Equation 5

5.2 Manure Belt Operations

The literature review (Section 3) and exploratory data analysis (Section 4) suggested that, of the data collected during NAEM ambient temperature, exhaust temperature, ambient relative humidity, exhaust relative humidity, management phase, hen age, inventory, and live animal weight should be considered in the development of emission models for the layer manure belt houses. The various combinations of these parameters were used in test models for the pollutants of interest. For NH₃ and H₂S, 12 different combinations were tested as potential models (Table 5-4). There were 16 models (Table 5-5) tested for particulate matter emissions, which had more variations using the relative humidity parameters.

Like the high rise models, models G-6, G-10, P-13, and P-14 include the management phase categories as a parameter. As noted in Section 3, the management activities and general movement in the house can have an impact on emissions, particularly particulate matter emissions. However, EPA is considering these models as experimental since an appropriate methodology for their evaluation and application has not been finalized. The models have been included in the tables to note all the options EPA explored, but were not considered as potential models at this time.

Table 5-4. Parameter combinations tested as models for manure belt house NH₃ and H₂S emissions.

Model	Parameters
G-1	Intercept, Live animal weight, Ambient temperature
G-2	Intercept, Live animal weight, Exhaust temperature
G-3	Intercept, Inventory, Ambient temperature
G-4	Intercept, Inventory, Exhaust temperature
G-5	Intercept, Inventory, Hen age, Ambient temperature
G-6	Intercept, Inventory, Ambient temperature, Management phase (flock emptying and replacement (E),
G-0	full flock (F), molting (M), and transition (T))
G-7	Intercept, Inventory, Ambient temperature, Ambient relative humidity
G-8	Intercept, Inventory, Ambient temperature, Exhaust relative humidity
G-9	Intercept, Inventory, Ambient temperature, Ambient relative humidity, Management phase (E,F,M,T)
G-10	Intercept, Inventory
G-11	Intercept, Inventory, Exhaust temperature, Ambient relative humidity
G-12	Intercept, Inventory, Exhaust temperature, Exhaust relative humidity

Table 5-5. Parameter combinations tested as models for manure belt house PM₁₀, PM_{2.5}, and TSP emissions.

Model	Parameters
P-1	Intercept, Inventory
P-2	Intercept, Inventory, Ambient relative humidity
P-3	Intercept, Inventory, Exhaust relative humidity
P-4	Intercept, Inventory, Ambient relative humidity, Ambient temperature
P-5	Intercept, Inventory, Ambient temperature, Exhaust relative humidity
P-6	Intercept, Inventory, Ambient relative humidity, Exhaust temperature
P-7	Intercept, Inventory, Exhaust temperature, Exhaust relative humidity
P-8	Intercept, Live animal weight, Ambient temperature, Ambient relative humidity
P-9	Intercept, Live animal weight, Ambient temperature, Exhaust relative humidity
P-10	Intercept, Live animal weight, Ambient relative humidity, Exhaust temperature
P-11	Intercept, Live animal weight, Exhaust temperature, Exhaust relative humidity
P-12	Intercept, Hen age, Inventory, Ambient relative humidity
P-13	Intercept, Inventory, Ambient relative humidity, Management phase (C,E,F,M,T)
P-14	Intercept, Inventory, Ambient temperature, Ambient relative humidity, Management phase (C,E,F,M,T)
P-15	Intercept, Inventory, Ambient temperature
P-16	Intercept, Inventory, Exhaust temperature

For NH₃ (Appendix G, Table G-15), models G-2, G-4, G-5, G-11, and G-12 had terms that were not statistically significant (p > 0.05) and were removed from further consideration. The model fit (-2 log likelihood, AIC, AICc, and BIC) and evaluation statistics (ME, NME, MB, NMB) for NH₃ (Appendix G, Table G-16) indicate the remaining models were comparable, which suggested using ambient parameters would incorporate regional differences due to climate. It also suggests that the ambient parameters were as effective as house parameters in estimating the effects of climate differences. Therefore, considering the potential ease of data collection, EPA concluded that ambient temperature and relative humidity would be preferable

to one with exhaust temperature and relative humidity, while considering the environmental impacts on the emissions. Of the remaining models (G-1, G-3, and G-7), EPA selected model G-7 (including the parameters: intercept, inventory, ambient temperature, ambient relative humidity) for further analysis for NH₃. The final form of this model is presented in Table 5-6.

For H₂S (Appendix G, Table G-17), model G-5 had terms that were not statistically significant and was removed from further consideration. The model fit (-2 log likelihood, AIC, AICc, and BIC) and evaluation statistics (ME, NME, MB, NMB) for H₂S (Appendix G, Table G-18) indicate the remaining models were comparable. This suggested using ambient parameters in the models was as effective as house parameters when evaluating temperature and humidity effects on emissions. Therefore, EPA considered the potential ease of data collection and concluded that a model ambient temperature and relative humidity would be preferable to one with exhaust temperature and relative humidity. Of the remaining models that used ambient parameters, EPA selected model G-7 (including the parameters: intercept, inventory, ambient temperature, ambient relative humidity) for further analysis for H₂S as it had the one of the lowest normalized mean bias of the remaining models, and was consistent with the parameters selected for the NH₃ model, limiting the data collection burden. The selected model is presented in Table 5-6.

For PM₁₀ (Appendix G, Table G-19), only models P-1 and P-16 were comprised of terms that were statistically significant (p < 0.05). EPA thoroughly reviewed the data to determine potential reasons for the lack of significant models. The ability of predictor variables to represent the chemical, physical and biological processes that control emissions will impact the performance of statistical models. In terms of PM emissions from manure belt houses, the two largest variables that will influence emissions is 1) the amount of source material (i.e. excreted manure, feathers, and feed) in the house, which is related to inventory; and 2) the amount of disturbance of the source materials, which is related to layer, human, and management activity. For manure belt houses, sufficient data was not collected during NAEMS to develop a variable that represents the second factor of activity inside the manure belt house. Not adequately capturing the variance of this key parameter makes it hard to determine the effect of other less influential parameters (i.e. relative humidity and temperature). The effect of not having a variable that represents house activity varies for each of the particulate matter models, depending on how much the activity varies at a house/barn and then how much activity influences emissions. Activity level in the house does not have as strong an effect on gas emissions, which is why more of those models had significant parameters.

In addition, for a model to determine the influence of a parameters it has to vary enough for its influence can be determined above the effect of all the other predictive parameters. For the manure belt house models there is only one site, consisting of two houses, which have a fairly steady inventory apart from one flock replacement event at H9. This means the influence of inventory on emissions cannot be easily quantified. The lack of variance of the inventory will affect both gas and particulate matter emissions. However, it will have a more pronounced effect on the PM₁₀ models since there are only six daily emission values during the flock replacement event, while NH₃ and H₂S have 20 and 21 days, respectively.

The model fit evaluation statistics for PM₁₀ (Appendix G, Table G-20) indicate model P-1 and P-16 performed similarly. Therefore, EPA selected model P-1 (including the parameter: inventory) for further analysis and consideration. The full form of the model is presented in Table 5-3.

As previously noted, the particulate matter model selection starts with the PM_{10} models, as there is more emission data available. The PM_{10} version of the models had between 460 and 566 days worth of data, depending on the completeness of the various predictive parameters. For $PM_{2.5}$ and TSP, the number of records available ranged between 26-34 for $PM_{2.5}$ and 66-69 for TSP. This is substantially less data that was available for PM_{10} and does not cover the same breadth of conditions that the PM_{10} data does. Therefore, the models generated with these smaller datasets were examined mainly for consistency with the PM_{10} results, to build confidence in using the same model form for all the particulate matter species.

Model P-1 (inventory including the parameter: inventory) have insignificant terms, unlike PM₁₀. The relationship inventory is consistent across the particulate matter size fractions. The PM_{2.5} model also has a negative intercept value. The negative intercept is likely due to the fact that values for 30 out of the 56 available days are negative. The statistics for PM_{2.5} (Appendix G, Table G-22) were fairly poor across all models, as the NME and NMB were greater than 100% for most of the models. The TSP model performance statistics (Appendix G, Table G-24) were nominally better than the PM_{2.5} models, as NME improved to 78%. The improvement in statistics is likely due to the increased amount of daily data available for model development compared to PM_{2.5}. Overall, model P-1 (including the parameter: inventory) was selected for both PM_{2.5} and TSP, as it would be consistent with the parameters selected for the PM₁₀ model. The full forms of the models are presented in Table 5-3.

Table 5-6. Selected daily models for manure belt layer houses.

Pollutant	Formula	Equation Number
NH ₃	$ln(NH_3) = 2.4392 + 0.0047 * Inventory + 0.0294 * Amb_T + 0.0019 * Amb_{RH}$	Equation 6
H ₂ S	$ln(H_2S) = 3.7391 + 0.0073 * Inventory + 0.0222 * Amb_T + 0.0048 * Amb_{RH}$	Equation 7
PM ₁₀	$ln(PM_{10}) = 6.631005 + 0.007205 * Inventory$	Equation 8
PM _{2.5}	$ln(PM_{2.5}) = -127.4489 + 0.534577 * Inventory$	Equation 9
TSP	ln(TSP) = 6.936206 + 0.00987 * Inventory	Equation 10

5.3 Manure Sheds

The literature review (Section 3) and exploratory data analysis (Section 4) suggested that EPA should consider ambient temperature, ambient relative humidity, wind speed, airflow, management phase, manure age, average hen age, inventory, and live animal weight in the development of the emission models for manure sheds associated with layer manure belt houses. As a reminder, the average hen age represented the average age from the two houses (IN2B H8 and H9) whose manure was stored in the shed. Similarly, the inventory and live animal weight represent the combined values for both houses that stored manure in the shed.

In addition to these parameters, EPA also used additional inventory and live animal weight that represents a value lagged by 5 days to account for the time it takes for the manure to be transported from the house to the shed via the conveyor belts. Again, the values are the combined value from both houses supplying manure to the shed.

The various combinations of these parameters were used in test models for the pollutants of interest. For NH₃ and H₂S, 20 different combinations were tested as potential models (Table 5-7). Thirteen models were tested for particulate matter emissions (Table 5-8).

EPA has included tested models that incorporated an indication of management phase into models G-9, G-18, G-19, and P-9. The management phase parameter for manure sheds incorporates the phase from both houses supplying manure to the shed, as well as any noted cleanout times in the manure shed itself. EPA considers these models experimental at this time, as validation and testing methods are still being vetted. They are included in this report to show all the options explored and may be pursued in a future version of the models.

Table 5-7. Parameter combinations tested as models for layer manure shed NH₃ and H₂S emissions.

Model	Parameter
G-1	Intercept, Inventory, Ambient temperature
G-2	Intercept, Inventory (5 day lag), Ambient temperature
G-3	Intercept, Live animal weight, Ambient temperature
G-4	Intercept, Live animal weight (5 day lag), Ambient temperature
G-5	Intercept, Inventory (5 day lag), Ambient temperature, Ambient relative humidity
G-6	Intercept, Inventory (5 day lag), Ambient temperature, Wind speed
G-7	Intercept, Inventory (5 day lag), Ambient temperature, Ambient relative humidity, Wind speed
G-8	Intercept, Inventory (5 day lag), Ambient temperature, Average hen age
	Intercept, Inventory (5 day lag), Ambient temperature, Management phase (Manure shed cleanout,
G-9	House 8 full & House 9 molting (CFF), House 8 full & House 9 empty (FE); House 8 full & House 9 full
G-9	(FF); House 8 full & House 9 molting (FM); House 8 full & House 9 transitioning (FT); and House 8
	molting & House 9 full (MF))
G-10	Intercept, Inventory (5 day lag), Ambient temperature, Manure age
G-11	Intercept, Inventory (5 day lag), Wind speed
G-12	Intercept, Inventory (5 day lag), Manure age
G-13	Intercept, Inventory (5 day lag), Wind speed, Manure age
G-14	Intercept, Inventory (5 day lag), Ambient relative humidity, Manure age
G-15	Intercept, Inventory (5 day lag), Airflow
G-16	Intercept, Ambient temperature, Airflow
G-17	Intercept, Airflow, manure age
G-18	Intercept, Inventory (5 day lag), Manure age, Management phase (CFF, FE, FF, FM, FT, and MF)
G-19	Intercept, Ambient Temperature, Manure age, Management phase (CFF, FE, FF, FM, FT, and MF)
G-20	Intercept, Inventory (5 day lag)

Table 5-8. Parameter combinations tested as models for layer manure shed PM_{10} , $PM_{2.5}$, and TSP emissions.

Model	Parameter
P-1	Intercept, Inventory, Airflow
P-2	Intercept, Inventory (5 day lag), Airflow
P-3	Intercept, Live animal weight, Airflow
P-4	Intercept, Live animal weight (5 day lag), Airflow
P-5	Intercept, Inventory (5 day lag), Wind speed
P-6	Intercept, Inventory (5 day lag), Ambient temperature
P-7	Intercept, Inventory (5 day lag), Ambient relative humidity
P-8	Intercept, Inventory (5 day lag), Average hen age
P-9	Intercept, Inventory (5 day lag), Management phase (CFF, FE, FF, FM, FT, and MF)
P-10	Intercept, Inventory (5 day lag), Manure age
P-11	Intercept, Inventory (5 day lag)
P-12	Intercept, Live animal weight (5 day lag)
P-13	Intercept, Airflow

For NH₃ (Appendix G, Table G-27), only models G-16 (includes parameters: intercept, ambient temperature, airflow) and G-17 (includes parameters: intercept, airflow, manure age) were entirely comprised of terms that were statistically significant (p < 0.05). The H₂S analysis (Appendix G, Table G-29) had similar results with models G-2 (includes parameters: intercept, inventory (5 day lag), ambient temperature), G-16, and G-17 receiving further consideration. The model fit (-2 log likelihood, AIC, AICc, and BIC) and evaluation statistics (ME, NME, MB, NMB) for NH₃ (Appendix G, Table G-28) and H₂S (Appendix G, Table G-30) indicates these remaining models were comparable. Models G-16 and G-17 both contained airflow, which is not be an easy parameter for a producer to calculate, as it requires hourly wind measurements in several openings of the structure (SOP A10, 2009). Depending on the structure, the number of anemometers needed may be cost prohibitive for operators.

Models G-16 and G-17 do not include a parameter that indicates the number of birds or size of the operation, which had previously been used as a proxy for the volume of manure produced. The volume of manure produced is a considerable factor in the emissions, as emissions are released from the surface of the manure. The issue in using inventory as a proxy for volume is that there is a lack of variation of inventory in this study. As mentioned earlier, for a model to determine the influence of a parameter it has to vary enough for that its influence to be determined above the effect of other predictive parameters. For the manure shed models there is only one site for the dataset. The two houses suppling the manure shed have a fairly steady inventory apart from one flock replacement event at one house. This means the influence of inventory, as a proxy for manure volume, on emissions cannot be easily identified. The lack of variance of the inventory will affect both gas and particulate matter emissions modeling efforts and can be seen as a limitation in the dataset.

Acknowledging the need for to indicate the size of the operation, and therefore an estimate of the volume of manure produced, the EPA selected the model G-2 (includes parameters: intercept, inventory (5 day lag), ambient temperature) for both NH₃ and H₂S. The established relationship between inventory and emissions trumps the insignificance finding in the NH₃ model tests. It should be noted that the ambient temperature coefficient is negative for the NH₃ and H₂S models. A possible explanation for this, is that the higher temperatures are drying out the manure resulting in less NH₃ and H₂S generation. The final forms of the NH₃ and H₂S models are presented in Table 5-9.

For PM₁₀ (Appendix F, Tables F-31, and F-32), PM_{2.5} (Appendix F, Tables F-33 and F-34), and TSP (Appendix F, Tables F-35 and F-36), only model P-13 (includes parameters: intercept, airflow) was comprised of significant parameters. As with the models for NH₃ and H₂S, it does not include a parameter to indicate the number of birds or size of the operation. The primary mechanism to generate particulate matter emissions is the disruption of the manure pile,

either by agitation of the pile by manure dropping off the belt or human activity within the shed (i.e., manure removal). Surface area would be a better indicator of exposed manure that could generate emissions. However, this was not estimated during the study, aside from the square footage of the shed. In other models, inventory has served as a proxy for the volume of manure produced, which would affect the surface area. However, the modeling dataset lacks enough variability in inventory values to yield a statistically significant relationship with inventory or live animal weight.

Again, EPA was faced with the choice to either adhere to the established protocol of selecting from models with only significant parameters or deviate from it to select a model with relationship established by literature. Acknowledging the need for to indicate the size of the operation, and therefore an estimate of the volume of manure produced, the EPA elected to select a model that included an indication of size. Of the model that included size and did not have airflow, EPA selected model P-11 (includes parameters: intercept, inventory (5 day lag)) for further consideration for PM₁₀, PM_{2.5}, and TSP. The final form of these models are presented in Table 5-9.

Table 5-9. Selected daily models for layer manure sheds.

Pollutant	Formula	Equation Number
NH ₃	$ln(NH_3) = -0.194945 + 0.003927 * Inventory (5 day lag) - 0.013752 * Amb_T$	Equation 11
H ₂ S	$ln(H_2S) = 1.295775 + 0.004976 * Inventory (5 day lag) - 0.024164 * Amb_T$	Equation 12
PM ₁₀	$ln(PM_{10}) = 4.5366 + 0.000732 * Inventory (5 day lag)$	Equation 13
PM _{2.5}	$ln(PM_{2.5}) = -30.57734 + 0.067599 * Inventory (5 day lag)$	Equation 14
TSP	ln(TSP) = 4.041666 + 0.002286 * Inventory (5 day lag)	Equation 15

6.0 MODEL COEFFICIENT EVALUATION

To ensure reliable prediction of the emissions, the model coefficients were evaluated with the jackknife method (Christensen et al., 2016; Leeden et al., 2007), which examined the cumulative effect on coefficient estimates of multiple "minus-one" runs. The jackknife approach called for removing one of the independent sample units from the dataset. For NAEMS, the individual houses at each site and the monitored sheds are the mutually exclusive independent sample units. EPA then determined the associated parameter estimates for the selected model based on this dataset. This was repeated for each of the sample units. These results were then compared to the model coefficients based on the full dataset (full model). For each jackknife model, the ME, NME, MB, and NMB were calculated, based on the equations outlined in Section 6 of the main report, to facilitate comparison.

EPA also prepared plots showing the variation in coefficients and standard errors for the selected models and compared to each of the jackknife models. EPA interpreted these plots similar to the Tukey confidence interval plots in that, if the result for the jackknife model overlapped the results for the full model (i.e., the area highlighted in gray on the figures), then the model coefficients are not inconsistent with one another. If the omission of one monitoring unit (e.g., a house or shed) resulted in a coefficient that was outside ± 1 standard error of the full model, the sample unit was reviewed to determine if a specific characteristic of that unit (e.g., animal placement strategy, manure handling system) might have caused the inconsistency. If the difference could not be ascribed to an operational characteristic of the unit, the data were reviewed for outliers that could be trimmed, and other potential remediation measures considered.

6.1 High Rise Layer House Models

6.1.1 NH₃ Model Evaluation

Table 6-1 shows the variation in coefficients and standard errors for the selected model ("None") and each of the jackknife models. The model coefficients from the jackknife approach were comparable across the withheld sets (Table 6-1) and remained significant across all models. The plots in Figure 6-1 show that the results for all jackknife models overlap the full model estimate \pm 1 standard error, except IN2H H7, which was just outside of this range for the ambient temperature and inventory. In comparison to the full model, that is where the site removed is "None", the maximum percent differences for parameter estimates across the six models were 11%, 12%, 5%, and 48% for intercept, ambient relative humidity, ambient temperature, and inventory, respectively. Across all models, the difference in NME and NMB (Table 6-2) in comparison to the selected model were moderate, with NME values differing by less than 6.15% and NMB by less than 4.38%.

Table 6-1. Model coefficients developed using the jackknife approach for NH₃ emissions from high rise houses.

			Standard	
Site out	Effect	Estimate	Error	p-value
None	Intercept	2.659821	0.22567	<.0001
None	Ambient Relative Humidity	0.001761	0.00031	<.0001
None	Ambient Temperature	0.038714	0.00097	<.0001
None	Inventory	0.00589	0.00126	<.0001
CA2BH5	Intercept	2.693697	0.02272	<.0001
CA2BH5	Ambient Relative Humidity	0.001914	0.00032	<.0001
CA2BH5	Ambient Temperature	0.037616	0.001	<.0001
CA2BH5	Inventory	0.005876	0.00071	<.0001
CA2BH6	Intercept	2.687925	0.10464	<.0001
CA2BH6	Ambient Relative Humidity	0.001901	0.0003	<.0001
CA2BH6	Ambient Temperature	0.037399	0.00095	<.0001
CA2BH6	Inventory	0.005869	0.00082	<.0001
IN2HH6	Intercept	2.651921	0.2604	<.0001
IN2HH6	Ambient Relative Humidity	0.001555	0.00033	<.0001
IN2HH6	Ambient Temperature	0.042286	0.0011	<.0001
IN2HH6	Inventory	0.00523	0.0016	0.0012
IN2HH7	Intercept	2.380192	0.1567	<.0001
IN2HH7	Ambient Relative Humidity	0.001594	0.00034	<.0001
IN2HH7	Ambient Temperature	0.040827	0.00112	<.0001
IN2HH7	Inventory	0.008688	0.00125	<.0001
NC2BH3	Intercept	2.656733	0.44505	<.0001
NC2BH3	Ambient Relative Humidity	0.001793	0.00038	<.0001
NC2BH3	Ambient Temperature	0.03862	0.00116	<.0001
NC2BH3	Inventory	0.005869	0.00131	<.0001
NC2BH4	Intercept	2.764139	0.3186	<.0001
NC2BH4	Ambient Relative Humidity	0.0017	0.00035	<.0001
NC2BH4	Ambient Temperature	0.03673	0.00107	<.0001
NC2BH4	Inventory	0.005059	0.00153	0.001

Table 6-2. Model fit statistics for the high rise house NH₃ jackknife.

Site out	n	LNME ^a (%)	NME ^b (%)	ME ^b (kg day ⁻¹)	MB ^b (kg day ⁻¹)	NMB ^b (%)	Corr
None	3562	16.851	59.12	60.818	0.165	0.161	0.516
CA2BH5	3035	16.042	58.735	67.574	1.988	1.728	0.489
CA2BH6	3016	14.159	56.459	65.635	1.524	1.311	0.493
IN2HH6	3025	18.044	63.521	51.826	1.72	2.108	0.45
IN2HH7	3023	15.499	52.975	41.132	3.527	4.543	0.628
NC2BH3	2852	18.734	61.241	69.816	0.233	0.204	0.516
NC2BH4	2859	19.066	62.574	71.287	-0.254	-0.223	0.484

^a Based on transformed data (i.e., ln(NH₃)).

^b Based on back-transformed data.




Figure 6-1. Comparison of variation in coefficients and standard errors for NH₃ high rise house

Variation in coefficients and standard errors (blue closed circle and \pm SE bar) for each jackknife model with the selected NH₃ manure belt house model coefficient ("None", gray band for \pm SE) for each model parameter.

6.1.2 H₂S Model Evaluation

Table 6-3 shows the variation in coefficients and standard errors for the selected model ("None") and each of the jackknife models. The model coefficients from the jackknife approach were comparable across the withheld sets (Table 6-3) and remained significant across all models. The plots in Figure 6-2 show that the results for all jackknife models overlap the full model estimate \pm 1 standard error, except for NC2B H4 for intercept and inventory, and NC2B H3 for ambient temperature and relative humidity. In comparison to the full model, the maximum percent differences for parameter estimates across the six models were 9%, 38%, 15%, and 43% for intercept, ambient temperature, inventory, and ambient relative humidity, respectively. Across all models, the difference in NME and NMB (Table 6-4) in comparison to the selected model were moderate, with NME values differing by less than 3.31% and NMB by less than 7.77%.

Table 6-3. Model coefficients developed using the jackknife approach for H₂S emissions from high rise houses.

Site out	Effect	Estimate	Standard Error	p-value
NONE	Intercept	2.723104	0.07259	<.0001
	Ambient Temperature	0.020988	0.00163	<.0001
	Inventory	0.009798	0.00049	<.0001
	Ambient Relative Humidity	0.003752	0.00053	<.0001
CA2BH5	Intercept	2.610429	0.07377	<.0001
	Ambient Temperature	0.020363	0.0017	<.0001
	Inventory	0.010621	0.00049	<.0001
	Ambient Relative Humidity	0.003796	0.00055	<.0001
CA2BH6	Intercept	2.671677	0.07674	<.0001
	Ambient Temperature	0.020253	0.00178	<.0001
	Inventory	0.010282	0.00051	<.0001
	Ambient Relative Humidity	0.003665	0.00058	<.0001
IN2HH6	Intercept	2.854195	0.08094	<.0001
	Ambient Temperature	0.018927	0.00168	<.0001
	Inventory	0.009165	0.00066	<.0001
	Ambient Relative Humidity	0.003257	0.00054	<.0001
IN2HH7	Intercept	2.739702	0.07635	<.0001
	Ambient Temperature	0.019068	0.00167	<.0001
	Inventory	0.01021	0.00057	<.0001
	Ambient Relative Humidity	0.00349	0.00053	<.0001
NC2BH3	Intercept	2.573999	0.08696	<.0001
	Ambient Temperature	0.028865	0.00202	<.0001
	Inventory	0.009316	0.0005	<.0001
	Ambient Relative Humidity	0.005353	0.00071	<.0001
NC2BH4	Intercept	2.970026	0.09319	<.0001
	Ambient Temperature	0.020607	0.00196	<.0001
	Inventory	0.008355	0.00054	<.0001
	Ambient Relative Humidity	0.003926	0.00061	<.0001

Table 6-4. Model fit statistics for the high rise house H₂S jackknife.

Site out	n	LNME ^a (%)	NME ^b (%)	ME ^b (g day ⁻¹)	MB ^b (g day ⁻¹)	NMB ^b (%)	Corr
None	3291	9.962	52.695	58.631	-1.931	-1.735	0.721
CA2BH5	2733	9.537	52.389	65.238	-1.327	-1.066	0.715
CA2BH6	2715	9.973	52.54	66.589	-1.649	-1.301	0.71
IN2HH6	2908	9.629	52.417	44.002	-1.9	-2.263	0.71
IN2HH7	2901	9.482	49.388	43.181	0.999	1.143	0.739
NC2BH3	2595	10.335	53.332	66.998	-2.476	-1.971	0.726
NC2BH4	2603	10.334	54.317	67.339	-11.78	-9.503	0.714

^a Based on transformed data (i.e., ln(H₂S)).

^b Based on back-transformed data.




Figure 6-2. Comparison of variation in coefficients and standard errors for H₂S high rise house model.

Variation in coefficients and standard errors (blue closed circle and \pm SE bar) for each jackknife model with the selected H₂S manure belt house model coefficient ("None", gray band for \pm SE) for each model parameter.

6.1.3 PM₁₀ Model Evaluation

Table 6-5 shows the variation in coefficients and standard errors for the selected model ("None") and each of the jackknife models. The model coefficients from the jackknife approach were comparable across the withheld sets (Table 6-5) and remained significant across all models. The plots in Figure 6-3 show that the results for all jackknife models overlap the full model estimate ± 1 standard error, except for IN2H H7 for inventory, and NC2B H3 and CA2B H6, which are just outside the range for ambient temperature. In comparison to the full model, the maximum percent differences for parameter estimates across the six models were 2%, 27%, 19%, and 24% for intercept, inventory, ambient relative humidity, and ambient temperature, respectively. Across all models, the difference in NME and NMB (Table 6-6) in comparison to the selected model were moderate, with NME values differing by less than 7.96% and NMB by less than 15.68%.

Table 6-5. Model coefficients developed using the jackknife approach for PM₁₀ emissions from high rise houses.

Site out	Effect	Estimate	Standard Error	p-value
None	Intercept	6.870178	0.06748	<.0001
	Inventory	0.007684	0.00054	<.0001
	Ambient Relative Humidity	-0.003022	0.00051	<.0001
	Ambient Temperature	0.014477	0.00153	<.0001
CA2BH5	Intercept	6.846607	0.07861	<.0001
	Inventory	0.008111	0.00063	<.0001
	Ambient Relative Humidity	-0.003125	0.00054	<.0001
	Ambient Temperature	0.012766	0.00162	<.0001
CA2BH6	Intercept	6.958724	0.07918	<.0001
	Inventory	0.007213	0.00062	<.0001
	Ambient Relative Humidity	-0.002751	0.00054	<.0001
	Ambient Temperature	0.011153	0.00162	<.0001
IN2HH6	Intercept	6.807521	0.06686	<.0001
	Inventory	0.008409	0.00058	<.0001
	Ambient Relative Humidity	-0.003222	0.00052	<.0001
	Ambient Temperature	0.016333	0.00156	<.0001
IN2HH7	Intercept	6.742102	0.07799	<.0001
	Inventory	0.009771	0.00079	<.0001
	Ambient Relative Humidity	-0.003175	0.00052	<.0001
	Ambient Temperature	0.014566	0.0016	<.0001
NC2BH3	Intercept	6.868256	0.08028	<.0001
	Inventory	0.007341	0.00057	<.0001
	Ambient Relative Humidity	-0.002505	0.00062	<.0001
	Ambient Temperature	0.018019	0.00191	<.0001
NC2BH4	Intercept	6.87016	0.07509	<.0001
	Inventory	0.006952	0.00053	<.0001
	Ambient Relative Humidity	-0.003055	0.00066	<.0001
	Ambient Temperature	0.015921	0.00185	<.0001

Table 6-6. Model fit statistics for the high rise house PM₁₀ jackknife.

Site out	n	LNME ^a (%)	NME ^b (%)	ME ^b (g day ⁻¹)	MB ^b (g day ⁻¹)	NMB ^b (%)	Corr
None	2623	4.977	50.241	1218.1	111.63	4.604	0.617
CA2BH5	2184	5.099	51.3	1360	164.59	6.208	0.594
CA2BH6	2125	5.038	50.092	1382.4	107.75	3.905	0.572
IN2HH6	2258	4.671	48.397	1037.8	99.102	4.622	0.66
IN2HH7	2234	4.979	58.197	1141.4	397.88	20.288	0.59
NC2BH3	2246	5.2	50.708	1305.7	81.332	3.159	0.617
NC2BH4	2068	5.058	50.456	1253.5	-12.53	-0.504	0.633

^a Based on transformed data (i.e., In(PM₁₀)).

^b Based on back-transformed data.




Figure 6-3. Comparison of variation in coefficients and standard errors for PM_{10} high rise house model.

Variation in coefficients and standard errors (blue closed circle and \pm SE bar) for each jackknife model with the selected PM₁₀ manure belt house model coefficient ("None", gray band for \pm SE) for each model parameter.

6.1.4 PM_{2.5} Model Evaluation

Table 6-7 shows the variation in coefficients and standard errors for the selected model ("None") and each of the jackknife models. The model coefficients from the jackknife approach were comparable across the withheld sets (Table 6-7) and remained significant across all models. The plots in Figure 6-4 show that the results for all jackknife models overlap the full model estimate \pm 1 standard error, with NC2B H4 just falling into the range for ambient temperature. In comparison to the full model, the maximum percent differences for parameter estimates across the six models were 4%, 29%, 21%, and 48% for intercept, inventory, ambient relative humidity, and ambient temperature, respectively. Across all models, the difference in NME and NMB (Table 6-8) in comparison to the selected model were moderate, with NME values differing by less than 11.14% and NMB by less than 11.36%.

Table 6-7. Model coefficients developed using the jackknife approach for $PM_{2.5}$ emissions from high rise houses.

Site out	Effect	Estimate	Standard Error	p-value
None	Intercept	4.621874	0.46141	<.0001
	Inventory	0.008039	0.00336	0.0295
	Ambient Relative Humidity	-0.018133	0.00367	<.0001
	Ambient Temperature	0.051013	0.01149	<.0001
CA2BH5	Intercept	4.653248	0.38683	0.0003
	Inventory	0.007859	0.00222	0.0006
	Ambient Relative Humidity	-0.017236	0.00354	<.0001
	Ambient Temperature	0.042139	0.01067	0.0001
CA2BH6	Intercept	4.592269	0.56282	<.0001
	Inventory	0.007681	0.00393	0.0724
	Ambient Relative Humidity	-0.015357	0.00418	0.0005
	Ambient Temperature	0.037024	0.01334	0.0067
IN2HH6	Intercept	4.663189	0.44356	<.0001
	Inventory	0.006925	0.00359	0.0848
	Ambient Relative Humidity	-0.017512	0.00376	<.0001
	Ambient Temperature	0.049026	0.01204	<.0001
IN2HH7	Intercept	4.442859	0.48259	<.0001
	Inventory	0.010404	0.00387	0.0177
	Ambient Relative Humidity	-0.018169	0.00384	<.0001
	Ambient Temperature	0.054367	0.01256	<.0001
NC2BH3	Intercept	4.677209	0.43766	<.0001
	Inventory	0.00805	0.00258	0.0081
	Ambient Relative Humidity	-0.019376	0.00398	<.0001
	Ambient Temperature	0.054242	0.01191	<.0001
NC2BH4	Intercept	4.557335	0.50869	<.0001
	Inventory	0.007702	0.00267	0.0298
	Ambient Relative Humidity	-0.021994	0.00561	0.0002
_	Ambient Temperature	0.075354	0.0143	<.0001

Table 6-8. Model fit statistics for the high rise house PM_{2.5} jackknife.

Site out	n	LNME ^a (%)	NME ^b (%)	ME ^b (g day-1)	MB ^b (g day-1)	NMB ^b (%)	Corr
None	142	16.51	78.095	164.4	-44.37	-21.08	0.636
CA2BH5	117	13.6	76.908	132.53	-16.74	-9.716	0.504
CA2BH6	110	16.688	89.236	179.76	-22.82	-11.33	0.377
IN2HH6	126	17.276	74.491	157.15	-58.24	-27.61	0.785
IN2HH7	133	16.636	78.008	169.95	-42.58	-19.55	0.542
NC2BH3	124	17.206	74.787	174.13	-50.59	-21.73	0.677
NC2BH4	100	16.083	68.557	155.69	-63.25	-27.85	0.853

^a Based on transformed data (i.e., ln(PM_{2.5})).

^b Based on back-transformed data.




Figure 6-4. Comparison of variation in coefficients and standard errors for PM_{2.5} high rise house model.

Variation in coefficients and standard errors (blue closed circle and ± SE bar) for each jackknife model with the selected PM_{2.5} manure belt house model coefficient ("None", gray band for ± SE) for each model parameter.

6.1.5 TSP Model Evaluation

Table 6-9 shows the variation in coefficients and standard errors for the selected model ("None") and each of the jackknife models. The model coefficients from the jackknife approach were comparable across the withheld sets (Table 6-9) and remained significant across all models. In comparison to the full model, the maximum percent differences for parameter estimates across the six models were 3%, 32%, 30%, and 121% for intercept, inventory, ambient relative humidity, and ambient temperature, respectively. However, the plots in Figure 6-5 show that the results for all jackknife models overlap the full model estimate ± 1 standard error. Across all models, the difference in NME and NMB (Table 6-10) in comparison to the selected model were moderate, with NME values differing by less than 3.65% and NMB by less than 2.71%.

Table 6-9. Model coefficients developed using the jackknife approach for TSP emissions from high rise houses.

Site out	Effect	Estimate	Standard Error	p-value
None	Intercept	7.599452	0.30204	<.0001
	Inventory	0.007927	0.0012	<.0001
	Ambient Relative Humidity	-0.005795	0.00282	0.0423
	Ambient Temperature	0.01367	0.00924	0.1417
CA2BH5	Intercept	7.582551	0.29409	<.0001
	Inventory	0.00778	0.00144	<.0001
	Ambient Relative Humidity	-0.005215	0.00254	0.0412
	Ambient Temperature	0.012892	0.00808	0.1125
CA2BH6	Intercept	7.421154	0.35744	<.0001
	Inventory	0.009214	0.00159	<.0001
	Ambient Relative Humidity	-0.006977	0.00315	0.0296
	Ambient Temperature	0.018426	0.01036	0.0784
IN2HH6	Intercept	7.843589	0.32542	<.0001
	Inventory	0.005379	0.00196	0.0181
	Ambient Relative Humidity	-0.00759	0.00293	0.011
	Ambient Temperature	0.012946	0.00972	0.1855
IN2HH7	Intercept	7.594248	0.30187	<.0001
	Inventory	0.007871	0.00119	<.0001
	Ambient Relative Humidity	-0.004642	0.00285	0.1067
	Ambient Temperature	0.009979	0.00956	0.2991
NC2BH3	Intercept	7.38713	0.40201	<.0001
	Inventory	0.009109	0.00143	<.0001
	Ambient Relative Humidity	-0.006726	0.00388	0.0859
	Ambient Temperature	0.030217	0.01211	0.0145
NC2BH4	Intercept	7.747179	0.31464	<.0001
	Inventory	0.006533	0.00109	<.0001
	Ambient Relative Humidity	-0.004076	0.00305	0.1845
	Ambient Temperature	0.004942	0.00998	0.6215

Table 6-10. Model fit statistics for the high rise house TSP jackknife.

Site out	n	LNME ^a (%)	NME ^b (%)	ME ^b (g day ⁻¹)	MB ^b (g day ⁻¹)	NMB ^b (%)	Corr
None	221	7.467	40.869	1641.5	64.602	1.608	0.524
CA2BH5	194	7.198	39.65	1675.2	74.729	1.769	0.508
CA2BH6	189	8.084	42.694	1805.2	163.03	3.856	0.505
IN2HH6	206	8.128	41.585	1545.1	-32.51	-0.875	0.409
IN2HH7	204	7.317	37.223	1449	-42.89	-1.102	0.605
NC2BH3	176	8.241	43.467	1810.6	166.63	4	0.519
NC2BH4	136	5.894	40.763	1578.9	43.373	1.12	0.602

^a Based on transformed data (i.e., In(TSP)).

^b Based on back-transformed data.




Figure 6-5. Comparison of variation in coefficients and standard errors for TSP high rise house model.

Variation in coefficients and standard errors (blue closed circle and \pm SE bar) for each jackknife model with the selected TSP manure belt house model coefficient ("None", gray band for \pm SE) for each model parameter.

6.2 Manure belt Layer House Models

6.2.1 NH₃ Model Evaluation

The model coefficients from the jackknife approach were comparable across the withheld sets (Table 6-11), though a few parameters were insignificant in the withheld models. However, the plots in Figure 6-6 show that the results for all jackknife models overlap the full model estimate \pm 1 standard error, often with the average value falling within \pm 1 standard error. Comparing the full model to the withheld models, the maximum percent differences for parameter estimates across the two models were 17%, 28%, 5%, and 5% for intercept, inventory, ambient temperature, and ambient relative humidity, respectively. Across all models, the difference in NME and NMB (Table 6-12) in comparison to the selected model were minor, with NME values differing by less than 4.39% and NMB by less than 1.50%.

Table 6-11. Model coefficients developed using the jackknife approach for NH₃ emissions from manure belt houses.

Site out	Effect	Estimate	Standard Error	p-value
None	Intercept	2.439187	0.38084	<.0001
	Inventory	0.004716	0.00148	0.0015
	Ambient temperature	0.029431	0.0021	<.0001
	Ambient relative humidity	0.001858	0.0008	0.0211
IN2BH8	Intercept	2.370221	0.42525	<.0001
	Inventory	0.004633	0.0016	0.004
	Ambient temperature	0.030883	0.00319	<.0001
	Ambient relative humidity	0.001972	0.0012	0.1001
IN2BH9	Intercept	2.851005	2.69364	0.3089
	Inventory	0.003414	0.01076	0.756
	Ambient temperature	0.027856	0.00275	<.0001
	Ambient relative humidity	0.00176	0.00107	0.1006

Table 6-12. Model fit statistics for the manure belt house NH₃ jackknife.

		LNME	NME ^b	ME ^b	MB ^b	NMB ^b	
Site out	n	(%)	(%)	(kg day ⁻¹)	(kg day ⁻¹)	(%)	Corr
None	1159	12.551	58.798	40.195	9.866	14.432	-0.199
IN2BH8	583	13.255	63.191	41.67	10.508	15.934	-0.185
IN2BH9	576	11.646	54.568	38.639	9.286	13.114	-0.242




Figure 6-6. Comparison of variation in coefficients and standard errors for NH₃ manure belt house model.

Variation in coefficients and standard errors (blue closed circle and \pm SE bar) for each jackknife model with the selected NH₃ manure belt house model coefficient ("None", gray band for \pm SE) for each model parameter.

6.2.2 H₂S Model Evaluation

The model coefficients from the jackknife approach were comparable across the withheld sets (Table 6-13) and were significant, except for IN2B H9. As with NH₃, the plots in Figure 6-7 show that the results for all jackknife models overlap the full model estimate \pm 1 standard error. The maximum percent differences for parameter estimates compared to the full model were 32%, 69%, 18%, and 12% for intercept, inventory, ambient temperature, and ambient relative humidity, respectively. Across all models, the difference in NME and NMB (Table 6-14) in comparison to the selected model were moderate, with NME values differing by less than 8.45% and NMB by less than 2.65%.

Table 6-13. Model coefficients developed using the jackknife approach for H₂S emissions from manure belt houses.

Site out	Effect	Estimate	Standard Error	p-value
NONE	Intercept	3.739104	0.34302	<.0001
	Ambient temperature	0.022216	0.00179	<.0001
	Inventory	0.007345	0.00135	<.0001
	Ambient relative humidity	0.004788	0.00068	<.0001
IN2BH8	Intercept	3.678677	0.3546	<.0001
	Ambient temperature	0.026288	0.00279	<.0001
	Inventory	0.007002	0.00136	<.0001
	Ambient relative humidity	0.005352	0.00102	<.0001
IN2BH9	Intercept	2.545273	1.34145	0.0689
	Ambient temperature	0.019604	0.00225	<.0001
	Inventory	0.012431	0.00535	0.0283
	Ambient relative humidity	0.00436	0.00091	<.0001

Table 6-14. Model fit statistics for the manure belt house H₂S jackknife.

		LNME ^a	NME ^b	ME ^b	MB ^b	NMB ^b	
Site out	n	(%)	(%)	(g day ⁻¹)	(g day ⁻¹)	(%)	Corr
None	1185	6.18	38.385	185.96	7.55	1.559	0.338
IN2BH8	598	7.42	46.835	221.81	19.952	4.213	0.199
IN2BH9	587	4.894	31.064	153.93	1.635	0.33	0.503




Figure 6-7. Comparison of variation in coefficients and standard errors for H₂S manure belt house model.

Variation in coefficients and standard errors (blue closed circle and \pm SE bar) for each jackknife model with the selected H₂S manure belt house model coefficient ("None", gray band for \pm SE) for each model parameter.

6.2.3 PM₁₀ Model Evaluation

The model coefficients from the jackknife approach were comparable across the withheld sets (Table 6-15), except for IN2BH9. The intercept and inventory parameters for the IN2BH9 withheld model fall outside the full model estimate \pm 1 standard error band shown in Figure 6-8. The maximum percent differences for parameter estimates compared to the full model 122% and 435% for intercept and inventory, respectively. Across all models, the difference in NME and NMB (Table 6-16) in comparison to the selected model were moderate, with NME values differing by less than 8.30% and NMB by less than 0.77%.

Table 6-15. Model coefficients developed using the jackknife approach for PM₁₀ emissions from manure belt houses.

Site out	Effect	Estimate	Standard Error	p-value
None	Intercept	6.631005	0.74268	<.0001
	Inventory	0.007205	0.00304	0.0186
IN2BH8	Intercept	7.038744	0.75413	<.0001
	Inventory	0.006027	0.00309	0.0525
IN2BH9	Intercept	-1.475494	2.38503	0.5391
	Inventory	0.038528	0.00973	0.0002

Table 6-16. Model fit statistics for the manure belt house PM₁₀ jackknife.

Site out	n	LNME ^a (%)	NME ^b (%)	ME ^b (g day ⁻¹)	MB ^b (g day ⁻¹)	NMB ^b (%)	Corr
None	566	9.608	85.204	4619.6	-61.72	-1.138	0.187
IN2BH8	334	9.188	79.429	5386.5	-42.89	-0.632	0.135
IN2BH9	232	8.088	76.906	2664.2	-66.03	-1.906	0.323




Figure 6-8. Comparison of variation in coefficients and standard errors for PM_{10} manure belt house model.

Variation in coefficients and standard errors (blue closed circle and \pm SE bar) for each jackknife model with the selected PM₁₀ manure belt house model coefficient ("None", gray band for \pm SE) for each model parameter.

6.2.4 PM_{2.5} Model Evaluation

Table 6-17 shows the variation in coefficients and standard errors for the selected model ("None") and each of the jackknife models. The intercept and inventory parameters for the IN2BH8 withheld model fall outside the full model estimate \pm 1 standard error band shown in Figure 6-9. When compared to the full model, the coefficients vary up to 123% and 119% for intercept and inventory, respectively. This may be due to the model selection being based off of PM₁₀ data, and not PM_{2.5} data. However, this was necessary as there is a total of 34 days of PM_{2.5} data from both houses. The plots in Figure 6-9 show that the results for all jackknife models overlap the full model estimate \pm 1 standard error. Across all models, the difference in NME and NMB (Table 6-18) in comparison to the selected model were substantial, with NME values differing up to 94.40% and NMB by up to 110.07% from the full model.

Table 6-17. Model coefficients developed using the jackknife approach for PM_{2.5} emissions from manure belt houses.

Site out	Effect	Estimate	Standard Error	p-value
None	Intercept	-127.4489	61.0184	0.0681
	Inventory	0.534577	0.24656	0.0604
IN2BH8	Intercept	29.681669	68.2977	0.6864
	Inventory	-0.099217	0.27555	0.737
IN2BH9	Intercept	-152.4	41.9311	0.0073
	Inventory	0.635083	0.16867	0.0062

Table 6-18. Model fit statistics for the manure belt house PM_{2.5} jackknife.

		LNME ^a	NME ^b	ME ^b	MB ^b	NMB ^b	
Site out	n	(%)	(%)	(g day ⁻¹)	(g day ⁻¹)	(%)	Corr
None	34	24.719	158.33	485.85	322.42	105.07	0.337
IN2BH8	23	18.389	94.887	197.28	18.756	9.021	-0.138
IN2BH9	11	16.2	63.933	328.47	-25.7	-5.002	0.433




Figure 6-9. Comparison of variation in coefficients and standard errors for PM_{2.5} manure belt house model.

Variation in coefficients and standard errors (blue closed circle and \pm SE bar) for each jackknife model with the selected PM_{2.5} manure belt house model coefficient ("None", gray band for \pm SE) for each model parameter.

6.2.5 TSP Model Evaluation

The model coefficients from the jackknife approach were comparable across the withheld sets (Table 6-19), though the parameters were insignificant. When compared to the full model, the coefficients vary up to 45% and 123% for intercept and inventory, respectively. Again, this is largely due to the reduced number of days available for the TSP models. However, the plots in Figure 6-10 show that the results for all jackknife models overlap the full model estimate \pm 1 standard error. Across all models, the difference in NME and NMB (Table 6-20) in comparison to the selected model were moderate, with NME values differing by less than 24.70% and NMB by less than 0.69% from the full model.

Table 6-19. Model coefficients developed using the jackknife approach for TSP emissions from manure belt houses.

Site out	Effect	Estimate	Standard Error	p-value
None	Intercept	6.936206	8.87165	0.4404
	Inventory	0.00987	0.03594	0.7855
IN2BH8	Intercept	10.081357	11.6563	0.3995
	Inventory	-0.002242	0.04719	0.9627
IN2BH9	Intercept	7.026336	10.2175	0.5012
	Inventory	0.006512	0.04163	0.8776

Table 6-20. Model fit statistics for the manure belt house TSP jackknife.

		LNME	NMEb	MEb	MB ^b	NMBb	
Site out	n	(%)	(%)	(kg day ⁻¹)	(kg day ⁻¹)	(%)	Corr
None	69	9.95	78.285	11668	82.753	0.555	-0.044
IN2BH8	34	9.001	76.513	16734	-28.81	-0.132	0.166
IN2BH9	35	10.914	53.585	4359.8	65.823	0.809	-0.157




Figure 6-10. Comparison of variation in coefficients and standard errors for TSP manure belt house model.

Variation in coefficients and standard errors (blue closed circle and \pm SE bar) for each jackknife model with the selected TSP manure belt house model coefficient ("None", gray band for \pm SE) for each model parameter.

6.3 Manure Shed Models

For the manure shed model, we did not complete jackknife analysis because there was only one site in the dataset. We also did not pursue a model evaluation using a k-fold cross validation technique based on previous SAB comments (SAB, 2013) recommending against using this method to select data for temporally correlated data. Future EPA efforts will look into obtaining additional data that would allow for further model testing and evaluation and an improved emission model.

7.0 ANNUAL EMISSION ESTIMATES AND MODEL UNCERTAINTY

To estimate annual pollutant emissions, the results of the daily emission models are summed over the number of operating days per year. This approach requires values for the necessary ambient and house parameters. For an actual emissions estimate, the daily estimates are based on meteorology from nearby monitors and house occupancy and weight records for the year from the producer. Since the models were developed with all the available data, producers can specify downtime for cleaning or other reasons with an inventory value of zero. For farms with multiple houses, annual emissions are determined for individual houses and summed across houses to calculate total annual farm-scale emissions.

As noted in Section 6 of the main report, the model results are transformed values of the emission. To convert to the native emission units (e.g., kg or g), the back transformation equation (Equation from Section 6 of the main All Sector report) is applied using the values of \overline{E}_l and C provided in Table 7-1 for each emission model. Section 8 contains an example of this calculation.

Table 7-1. Back transformation parameters

Table 7-1. Back transformation parameters										
Animal Type	Pollutant	\overline{E}_i	С	Resulting units						
High Rise House	NH ₃	1.58238	0	kg						
High Rise House	H ₂ S	1.24359	15	g						
High Rise House	PM ₁₀	1.11745	494	g						
High Rise House	PM _{2.5}	1.51089	37	g						
High Rise House	TSP	1.11429	0	g						
Manure Belt House	H₂S	1.09812	39	g						
Manure Belt House	NH ₃	1.27315	0	kg						
Manure Belt House	PM ₁₀	1.45218	1045	g						
Manure Belt House	PM _{2.5}	2.97703	108	g						
Manure Belt House	TSP	1.34146	696	g						
Manure Shed	H₂S	1.36619	6.0	g						
Manure Shed	NH ₃	1.28615	1.3	kg						
Manure Shed	PM ₁₀	1.68902	54.0	g						
Manure Shed	PM _{2.5}	1.68697	0.0	g						
Manure Shed	TSP	2.01361	30.0	g						

EPA also developed an estimate of uncertainty for total annual emissions, characterized by the random error in the model prediction using an approach similar to Monte Carlo analysis. Under this approach, EPA developed the statistical properties of predicted annual emissions by replicating annual sums of daily emissions. EPA ran these simulations for several different intervals of a predictor variable that fell within the observed range. For example, high rise house inventory ranged from 0 to 230 thousand head. The simulations were then run for inventory

intervals of 5 thousand head (e.g., 0, 5, 10, 15). Table 7-2 list the predictor variable and the number of intervals used for the annual uncertainty simulations for each model.

Simulations were run 10,000 times for each day for each interval to create an average uncertainty associated with the annual emissions from a single house. EPA added a random residual to each day of the simulation to replicate the variability that would be seen in a real-world application of the model. For each of the intervals run, EPA calculated standard statistics (i.e., minimum, median, mean, maximum, range) and used these to calculate the uncertainty for a single source via Equation 39:

Single source uncertainty =
$$0.5 \times \left(\frac{Range}{Median\ annual\ emission}\right) \times 100$$
 Equation 16

EPA then plotted this single house uncertainty against its associated annual emissions. This plot was then fit with a curve to model annual percent uncertainty for a single source (i.e., house, house, lagoon, basin). For all uncertainty models, the curve took the form of:

$$Uncertainty (\%) = \frac{k}{Annual \ Emissions}$$
 Equation 17

Where k is a constant, listed in Table 7-2, and annual emissions are the total sum from the daily models. EPA has not calculated particulate matter annual uncertainty models for the manure belt house and manure shed models in order to allow more time to optimize the models. EPA will include the annual uncertainty models in the final report.

Table 7-2. Annual Uncertainty Model Details

			Number of		Emission			
Animal Type	Pollutant	Simulation variable	Simulations	k	Units			
High Rise	H ₂ S	Inventory	10,000	452,862	g			
High Rise	NH₃	Inventory	10,000	328,272	kg			
High Rise	PM ₁₀	Inventory	10,000	7,670,567	g			
High Rise	PM _{2.5}	Inventory	10,000	1,076,172	g			
High Rise	TSP	Inventory	10,000	8,361,161	go			
Manure Belt House	H ₂ S	Inventory	10,000	923,681	g			
Manure Belt House	NH₃	Inventory	10,000	18,7085	kg			
Manure Belt House	PM ₁₀		a					
Manure Belt House	PM _{2.5}		a					
Manure belt House	TSP		a					
Manure Shed	H ₂ S	Inventory (% day lag)	10,000	179,952	g			
Manure Shed	NH ₃	Inventory (% day lag)	10,000	23,920.2	kg			
Manure Shed	PM ₁₀		a					
Manure Shed	PM _{2.5}	a						
Manure Shed	TSP	a						

^a Annual models were not calculated to allow time to optimize the daily models.

Multiplying this percentage by the annual emissions calculated for the source provides the resulting uncertainty in the native emission units (e.g., kg or g), demonstrated in Equation 40.

Resulting Uncertainty =
$$\frac{Percent uncertainty \times Annual emissions}{100}$$
 Equation 18

To propagate the uncertainty across all sources at a farm, EPA combined the estimates of absolute uncertainty for each source according to:

$$Total\ farm\ uncertainty = \sqrt{(U_{B1})^2 + \dots + (U_{Bi})^2 + (U_{L1})^2 + \dots + (U_{Lj})^2} \qquad \text{Equation 19}$$

Where:

Total farm uncertainty = total uncertainty for the total emissions from all farm sources. UBi = the resulting uncertainty for houses, with i representing the total number of houses on the farm,

ULj = the resulting uncertainty for manure sheds, with j representing the total number of open sources on the farm.

EPA notes that the uncertainty framework described above reflects the random uncertainty (error) in the prediction of daily emissions calculated using the emission models, which includes the random uncertainty in the measurements used to develop the equation. This framework does not, however, consider systematic error (e.g., bias) in either NAEMS measurements or the emission model. Section 8 provides an example of how the daily, annual, and annual uncertainty calculations are completed.

8.0 MODEL APPLICATION AND ADDITIONAL TESTING

Key to the development of any model is the demonstration of the use and practical examples of how the model behaves and replicates independent data. This section provides a series of example calculations to demonstrate the application of the models (Section 8.1), the sensitivity of the models to their inputs (Section 8.2), a comparison of the models developed to literature (section 8.3), and a test of model performance against an independent data set (Section 8.4). Finally, this section wraps up with a discussion of data limitations that could be driving sensitivity or performance issues.

8.1 Model Application Example

The following sections demonstrate how the daily emission models from Section 5 and the annual uncertainty from Section 7 are used to calculate emissions for an example farm for each structure type. Details about the use of the emission models to demonstrate compliance with Clean Air Act (CAA) thresholds will be addressed in a forthcoming implementation document. This example is provided to walk through a calculation to demonstrate how the system of equations is intended to work.

Section 6.4 of the main report noted that, since the data were log transformed prior to developing the models, the result would need to be back transformed to represent emissions in units of grams or kilograms. To complete the back transformation, users need two parameters that are specific to each model 1) \overline{E}_l , the average residual between model-predicted and observed (or measured) emissions on the natural log scale and 2) C, which is a constant added to the data prior to the log transformation. The values for \overline{E}_l and C for each layer model is provided in Table 7-1.

Once the emission models are finalized, EPA will work with stakeholders to develop a tool to facilitate the calculation of house and open source emissions. For transparency and to help stakeholders better understand the process of calculating emissions, this section will walk through example calculations to estimate NH₃ emissions from a high rise layer house, a manure belt house, and a manure shed.

The examples in this section use a fictional farm located in Hancock County, Iowa on January 1, 2020. Iowa was chosen as it is a top five egg producing state according to the USDA Economic Research Service (https://www.ers.usda.gov/topics/animal-products/poultry-eggs/sector-at-a-glance/). The ambient weather data used in each equation can be obtained for free from several sources including the National Centers for Environmental Information (NCEI; https://www.ncdc.noaa.gov/cdo-web/). NCEI stores hourly and daily ambient data from various monitors located across the country that can be used for emission estimation. The Forest City

Municipal Airport, IA site (WBAN: 54940), a Local Climatological Data (LCD) Station located in Hancock County. It's data file provides the daily average vales of the key meteorological parameters needed for calculations.

Additionally, the high rise and manure belt models require the number of birds in the house. For this fictious farm, 100,000 birds are placed in each house. The equations use thousands of birds, so this value will be divided by 1,000 for use in the emission models. A summary of the input values for the example calculations is provided in Table 8-1.

Table 8-1. Daily calculation parameter values

Parameter	Value
Daily Average Ambient Temperature (°C)	-0.9
Daily Average Relative Humidity (%)	89
Inventory (birds)	100,000

8.1.1 High Rise Example

Referring back to Equation 1, in Section 5, the log transformed values are calculated as follows:

$$\ln(NH_3) = 2.6598 + 0.0059 * Inventory + 0.0387 * Amb_T + 0.0018 * Amb_{RH}$$

$$\ln(NH_3) = 2.6598 + 0.0059 * \left(\frac{100,000}{1,000}\right) + 0.0387 * -0.9 + 0.0018 * 89$$

$$\ln(NH_3) = 2.6598 + 0.5890 - 0.0348 + 0.1567$$

$$\ln(NH_3) = 3.3707$$

To back transform the results to NH₃ in kg, use Equation 7, from the main report. For a high rise house, \overline{E}_i is 1.58238 and C is 0.

$$NH_3 = e^{3.3707} \times 1.58238 - 0$$

This comes to 46.09 kg NH₃ for the day. This process is repeated for each day, then the daily emissions are added together to get an annual estimate of emissions. After considering the values for each day in 2020, the total annual emissions for the house was calculated at 25,997.17 kg. To calculate the uncertainty associated with this estimate, use Equation 17 with the value of k from Table 7-1. This results in an annual uncertainty of:

Uncertainty (%) =
$$\frac{3,282.72}{25,997.17}$$
 = 12.63%

This translates to an uncertainty of \pm 3,282.72 kg. Thus, the final annual estimate for this house is 25,997.17 kg \pm 3,282.72 kg. This calculation would be repeated for any other high rise houses on the site.

8.1.2 Manure Belt House Example

Referring back to Equation 6, in Section 5, the log transformed NH₃ emission values for manure belt houses are calculated as follows:

$$ln(NH_3) = 2.4392 + 0.0047 * Inventory + 0.0294 * Amb_T + 0.0019 * Amb_{RH}$$

 $ln(NH_3) = 2.4392 + 0.0047 * \left(\frac{100,000}{100}\right) + 0.0294 * -0.9 + 0.0019 * 89$
 $ln(NH_3) = 2.4392 + 0.4700 - 0.0265 + 0.1691$
 $ln(NH_3) = 3.05184$

To back transform the results to NH₃ in kg, use Equation 7 from the main report. For a manure belt house, \overline{E}_l is 1.27315 and C is 0.

$$NH_3 = e^{3.05184} \times 1.27315 - 0$$

This comes to 26.95 kg NH₃ for the day. This process is repeated for each day, then the daily emissions are added together to get an annual estimate of emissions. After considering the values for each day in 2020, the total annual emissions for the House was calculated at 13,402.68 kg. To calculate the uncertainty associated with this estimate, use Equation 17 with the value of k from Table 7-1. This results in an annual uncertainty of:

Uncertainty (%) =
$$\frac{187,085}{13,402.68}$$
 = 13.96%

This translates to an uncertainty of \pm 1,870.85 kg. Thus, the final annual estimate for this house is 13,402.68 kg \pm 1,870.85 kg. This calculation would be repeated for any other manure belt houses on the site.

8.1.3 Manure Shed Example

Similar to the set up in NAEMS, the hypothetical farm will have two houses, with constant inventories of 200,000 birds, that move manure into a shed. Referring back to Equation 11, in Section 5, the log transformed NH₃ emission values for the manure shed are calculated as follows:

$$ln(NH_3) = -0.194945 - 0.01375 * Amb_T + 0.00393 * Inventory (5 day lag)$$

$$ln(NH_3) = -0.194945 - 0.01375 * -0.9 + 0.003927 * \left(\frac{200,000}{1,000}\right)$$
$$ln(NH_3) = -0.19494 + 0.0124 + 0.7854$$
$$ln(NH_3) = 0.60283$$

To back transform the results to NH₃ in kg, use Equation 7. For a manure shed, \bar{E}_l is 1.2862 and C is 1.3.

$$NH_3 = e^{0.60283} \times 1.2862 - 1.3$$

This comes to 1.05 kg NH₃ for the day. This process is repeated for each day, then the daily emissions are added together to get an annual estimate of emissions. After considering the values for each day in 2020, the total annual emissions for the house was calculated at 288.71 kg. To calculate the uncertainty associated with this estimate, use Equation 17 with the value of k from Table 7-1. This results in an annual uncertainty of:

Uncertainty (%) =
$$\frac{23,920.20}{288.71}$$
 = 82.85%

This translates to an uncertainty of \pm 239.20 kg. Thus, the final annual estimate for this house is 288.71 kg \pm 239.20 kg. This calculation would be repeated for any other manure sheds on the site.

8.1.4 Combining Structures

To calculate farm total emissions, the emissions from each unit are added. As an example, consider a farm with two high rise houses with a capacity of 100,000 birds. These houses will have the same emission estimate for the year, 25,997.17 kg \pm 3,282.72 kg. The annual farm emission estimate is:

$$Farm\ Total\ Emissions = 25,997.17 + 25,997.17 = 51,994.34\ kg\ NH_3$$

To estimate the total farm uncertainty, use Equation 19:

Total Farm Uncertainty =
$$\sqrt{U_{house\ 1}^2 + U_{house\ 2}^2}$$

Total Farm Uncertainty = $\sqrt{(3,282.72)^2 + (3,282.72)^2}$
Total Farm Uncertainty = 4,642.47 kg

The final annual NH_3 estimate for the farm is $51,994.34 \pm 4,642.47$ kg. Once the emission models are finalized, EPA will work with stakeholders to develop a tool to facilitate the calculation of house and manure shed emissions, which will also include the necessary ambient parameter data.

8.2 Model Sensitivity Testing

To further test the models, EPA varied the model parameters to ensure the model results would vary based on these key parameters. Two different tests were conducted: 1) bird placement was increased while the meteorological parameters were held constant, and 2) bird placement was held constant while the meteorological parameters were replaced with the values for a warmer climate.

8.2.1 Sensitivity to Inventory

To test the sensitivity to the bird population, the initial placement was increased to 150,000 birds. Using the same meteorology from Section 8.1, the emissions for a high rise house on January 1, 2020 is as follows:

$$\ln(NH_3) = 2.6598 + 0.0059 * \left(\frac{150,000}{1,000}\right) + 0.0387 * -0.9 + 0.0018 * 89$$

$$\ln(NH_3) = 2.6598 + 0.8835 - 0.0348 + 0.1567$$

$$\ln(NH_3) = 3.6652$$

$$NH_3 = e^{3.6652} \times 1.58238 - 0$$

This comes to 61.81 kg NH₃ for the day. This is 15.7 kg (34%) more than a house with a bird population of 100,000 layers for the same day. This demonstrates the model has sensitivity to the number of animals in the house. When the annual emission for this house is calculated, the annual difference is 8,902.86 kg, which is a 34% increase.

Looking at the manure belt house, increasing the bird population to 150,000 birds for January 1, 2020 results in the following:

$$ln(NH_3) = 2.4392 + 0.0047 * \left(\frac{150,000}{100}\right) + 0.0222 * -0.9 + 0.0048 * 89$$

 $ln(NH_3) = 2.4392 + 0.7050 - 0.0265 + 0.1691$
 $ln(NH_3) = 3.2868$
 $NH_3 = e^{3.2868} \times 1.27315 - 0$

This comes to 34.09 kg NH₃ for the day. This is 7.17 kg more NH₃ than a house with a bird population of 100,000 layers for the same day. When the annual emissions for this house are calculated, the annual difference is 3,550.49 kg, a 26% change with a 50% change in inventory. This demonstrates the model has a sensitivity to the number of animals in the house.

Looking at the manure shed, increasing the number of birds contributing to the house to 300,000 birds for January 1, 2020 results in the following:

$$ln(NH_3) = -0.194945 - 0.01375 * -0.9 + 0.003927 * \left(\frac{300,000}{1,000}\right)$$

 $ln(NH_3) = -0.194945 + 0.0127 + 1.1781$
 $ln(NH_3) = 0.99553$
 $NH_3 = e^{0.99553} \times 1.2862 - 1.3$

This comes to 2.18 kg NH₃ for the day. This is 1.13 kg more NH₃ than a shed supplied by a bird population of 200,000 layers for the same day. When the annual emissions for this shed is calculated, the annual difference is 367.71 kg, a 127% change with a 50% change in inventory. This demonstrates the model is sensitivity, perhaps overly, to the number of animals in the house.

8.2.2 Sensitivity to Climate

To further test model sensitivity, specifically that climate differences were producing different emission results, EPA calculated the emissions for the same farm in two distinctly different climate regions. The first was the theoretical farm from the previous example (Section 12.1) that is in northern Iowa. The NH₃ emission for this same farm setup (i.e., one high rise house and one manure belt house that empties into a manure shed) were calculated using meteorology from Dalton, Georgia. These locations were chosen based on 2017 Census of agriculture data indicating areas of poultry and egg sales (Figure 8-1).




Figure 8-1. 2017 Census of Agriculture plot indicating areas of broiler sales. Blue circles indicate approximate locations of test meteorology from lowa (IA) and Georgia (GA).

For our test sites, the temperatures from the Iowa (IA) site were typically lower than the Georgia (GA) site (Figure 8-1). Average daily temperatures differences between Iowa and Georgia by as little as 4 °C to as much as 35°C across the year. On average, the temperatures in Iowa were 8.4 °C less than those in Georgia (Table 8-2). With respect to relative humidity, the IA and GA sites were fairly similar during the warmer months (Apr-Oct; Figure 8-2 and Table 8-3), however, during the cooler months (Nov-Mar), average humidity was 16.2% higher in IA in comparison to GA. Both locations have humidities that vary between 35% to 100% across the year.




Figure 8-2. Comparison on temperatures at test locations in Iowa (IA) and Georgia (GA)

Table 8-2. Summary of temperature at the two meteorological sites

Site	Statistic	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Overall
	Min	-18.9	-22.7	-2.2	-3.2	5.3	17.7	17.6	15.4	7.5	-4.8	-4.5	-16.1	-22.7
IA	Max	0.8	3.6	10.8	16.9	20.7	27.9	27.7	25.7	22.2	19.4	16.5	3.3	27.9
	Average	-5.7	-5.9	3.3	7.0	13.8	22.4	23.2	21.1	15.4	6.5	3.7	-2.9	8.5
	Min	-1.2	2.0	6.7	9.1	10.4	20.0	23.5	22.7	15.4	11.7	6.0	-2.1	-2.1
GA	Max	18.9	18.2	23.2	21.9	25.4	27.0	29.2	27.5	27.7	22.7	22.6	14.6	29.2
	Average	8.6	8.7	15.0	15.2	19.3	24.0	26.9	25.5	22.4	18.2	12.7	6.3	16.9




Figure 8-3. Comparison of relative humidities at test locations IA and GA

Table 8-3. Summary of relative humidity at the two meteorological sites

Site	Statistic	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Overall
	Min	72.3	70.3	60.4	41.6	39.3	44.6	69.6	67.4	55.6	52.9	63.2	61.7	39.3
IA	Max	100.0	98.7	100.0	96.4	99.6	90.8	94.2	89.6	100.0	97.7	100.0	100.0	100.0
	Average	91.2	85.1	84.9	68.6	70.9	66.0	79.2	79.0	76.8	71.5	83.8	84.9	78.5
	Min	40.4	49.4	36.3	35.5	41.9	46.2	55.2	63.1	46.7	55.3	46.1	49.2	35.5
GA	Max	91.0	90.7	90.5	85.2	84.7	84.1	80.8	88.2	90.5	91.1	86.8	90.0	91.1
	Average	68.6	72.1	69.3	60.0	66.3	71.1	69.7	77.1	72.7	74.2	68.2	70.8	70.1

8.2.2.1 High rise house

When the daily calculations are performed for the entire year for a high rise house with 100,000 birds, the Georgia site typically has a higher daily emission values for the gaseous pollutants than the Iowa site (Figure 8-3). Table 8-4 has the estimated annual emission of all the pollutants studied. The total annual NH₃ emissions estimate for the farm using meteorology from Georgia was 25,997 kg, a 7,911 kg increase from the same high rise house with meteorology from Iowa. A similar trend is seen across the other pollutants. This is consistent with the trends of lower temperatures yielding lower gas emissions and higher humidity yielding lower PM emissions seen during the data exploration in Section 4. Overall, this suggests that the emission models are robust enough to account for the climatic differences of the different growing regions in the results for high rise houses.




Figure 8-4. Comparison of daily emission at test high rise locations IA and GA.

Table 8-4. Total annual emission from the theoretical high rise houses in IA and GA.

	IA Emission	GA Emissions
Pollutant	(kg per year)	(kg per year)
NH ₃	25,997	33,909
H ₂ S	25	29
PM ₁₀	590	702
PM _{2.5}	45	77
TSP	1,308	1,520

8.2.2.2 Manure belt house

For a manure belt house, when the daily calculations are performed for the entire year for a high rise house with 100,000 birds, the Georgia site typically has greater daily emission value for the gaseous pollutants than the Iowa site (Figure 8-4). Particulate matter estimates are the

same between the locations as there were no ambient parameters included in the selected models. Table 8-5 has the annual emission estimates of all the pollutants studied. The total annual NH₃ emissions estimate for the farm using meteorology from Georgia was 16,466 kg, a 3,063 kg increase from the same manure belt house with meteorology from Iowa. A similar trend is seen with H₂S. This is consistent with the trend of lower temperatures yielding lower gas emissions seen during the data exploration in Section 4. Overall, this suggests that the emission models are robust enough to account for the climatic differences of the different growing regions in the results for manure belt houses for the gaseous pollutant.




Figure 8-5. Comparison of daily emission at a theoretical manure belt house in IA and GA.

Table 8-5. Total annual emission from a theoretical manure belt house in IA and GA.

	IA Emission	GA Emissions
Pollutant	(kg per year)	(kg per year)
NH ₃	13,403	16,466
H₂S	49	58
PM ₁₀	446	446
PM _{2.5}	-40	-40
TSP	1,101	1,101

8.2.2.3 Manure Shed

For a manure shed, when the daily calculations are performed for the entire year the Georgia site has lower daily emission values for the gaseous pollutants than the Iowa site (Figure 8-5). Particulate matter estimates are the same between the locations as there were no ambient parameters included in the selected models. Table 8-6 has the estimate annual emissions of all the pollutants studied. The total annual NH₃ emissions estimate for the farm using meteorology from Iowa was 289 kg, an 88 kg difference from the same high rise house with meteorology from Georgia. A similar trend is seen with H₂S. Emissions of NH₃ and H₂S are higher when ambient temperatures are lower due to the negative ambient temperature coefficients in the models. As mentioned, a possible explanation for this is that the higher temperatures are leading to a drying of the manure and thus less NH₃ and H₂S generation. Overall, this suggests that the

emission models are robust enough to account for the climatic differences of the different growing regions in the results for high rise houses.




Figure 8-6. Comparison of daily emission at a theoretical manure shed in IA and GA.

Table 8-6. Total annual emission from a theoretical manure shed in IA and GA.

	IA Emission	GA Emissions
Pollutant	(kg per year)	(kg per year)
NH ₃	289	201
H ₂ S	2	1
PM ₁₀	47	47
PM _{2.5}	0	0
TSP	55	55

8.2.3 Model Limitations

As noted in the 2013 SAB review, extrapolating to conditions beyond those represented in the model development dataset could produce unreasonable results. To test the limitations of the model, EPA conducted a series of emission calculations over a range of conditions that could be seen at a farm in the US. These emission calculations tested one parameter at a time, with the selected parameter varied by a constant value through the range. For example, ambient temperature was increased by 1 °C from the minimum value in the model development dataset up to the maximum value. While one parameter was tested, the remaining parameters were held constant at the average value determined in the model development dataset. The emission values for each individual test were plotted on graphs for further examination.

If the sensitivity analysis produces unreasonable emissions or emission trends under certain conditions, this may indicate the need to limit the range of conditions that the model should be applied. Examples of unreasonable emissions or emission trends include unreasonably high (or low) emissions in certain conditions, and/or large changes in relative sensitivity (i.e., changes in sensitivity analysis slope). The following sections outlines the analysis for each of the selected models and provides a rudimentary examination of the sensitivity analysis for conditions where there may be unreasonable emissions or emission trends. It should be noted that this

analysis does not account for interaction between multiple terms within an equation, which could further affect the results. For example, a manure belt house with higher ambient temperatures would be able to cover a larger range of inventory before producing negative NH₃ emissions. Conversely, a house with lower ambient temperatures would cover a smaller range of inventory before producing negative NH₃ emission values.

8.2.3.1 High rise

All of the high rise house models included inventory, ambient temperature, and ambient relative humidity. The ranges tested for each parameter are in Table 8-7, with the plotted results plotted in Figure 8-7 and Figure 8-8. For the all the variables, the emissions increase with increasing values, which is consistent with established relationships for all three parameters. For inventory and ambient temperature, there are some changes in relative sensitivity, but the changes are not extreme. Neither the NH₃ nor H₂S models produce negative emissions under average conditions, which is an indicator of unreasonable emissions. For PM₁₀, PM_{2.5}, and TSP (Figure 8-8), only the PM_{2.5} model produces negative values when ambient temperatures are very low under average conditions.

Table 8-7. Parameter ranges tested for the high rise house models.

Parameter	Upper limit	Lower limit	Average Value	Increment
Ambient temperature (°C)	32	-25	15.4	0.9
Ambient relative humidity (%)	100	27	66.2	1
Inventory (birds)	338,800	0	90,522	4,600

To further explore any limitations in the models, emissions were calculated for 416,250 combinations of the range of values specified in Table 8-7. Across this range of conditions, neither the NH₃, PM₁₀, nor TSP models produce negative emissions. The models for H₂S and PM_{2.5} will produce negative values in instances of negative ambient temperature. Specifically, for H₂S, when the ambient temperature falls below -15°C and house inventory is less than 20,000 birds the models can produce negative emissions values. For PM_{2.5}, the range of values that can produce negative values increases to temperatures less than 7°C and inventory is less than 200,000. The plots in Figure 8-9 show the maximum values of live animal weight and ambient temperature that produce negative emissions at the relative humidity specified on the x-axis, but not necessarily in combination.




Figure 8-7. High rise house limitation tests for gaseous pollutants.

Visualization of the results for NH_3 (top row) and H_2S (bottom row) with tests for inventory (left), ambient temperature (center) and relative humidity (right).




Figure 8-8. High rise house limitation tests for particulate matter. Visualization of the results for PM_{10} (top row), $PM_{2.5}$ (center row), and TSP (bottom row) with tests for inventory (left) and ambient temperature (center), and ambient relative humidity (right).




Figure 8-9. Maximum values at which the high rise house models yield negative emissions. Visualization of the results for PM_{10} (top left), $PM_{2.5}$ (top right), and TSP (bottom).

8.2.3.2 Manure Belt Houses

For the manure belt house, the NH₃ and H₂S equations included inventory, ambient temperature, and ambient relative humidity. The ranges tested for each parameter are in Table 8-8 with the plotted results in Figure 8-10. For the all the variables, the emissions increase with increasing values, which is consistent with established relationships for all three parameters. Neither the NH₃ nor H₂S models produce negative emissions under average conditions, which is an indicator of unreasonable emissions.

PM₁₀, PM_{2.5}, and TSP models use only inventory. The range and average values used for testing are listed in Table 8-8 and the results are plotted in Figure 8-11. Neither PM₁₀ nor TSP models produce negative emissions under average conditions, which is an indicator of unreasonable emissions. For PM_{2.5}, the model produces negative emission for inventory levels less than 248,000 birds for average exhaust temperatures, which is unreasonable. The figure for PM_{2.5} also shows a rapid change in relative sensitivity with the model producing very high emission values for inventories greater than 260,000 birds. Overall, this analysis suggests that for PM_{2.5}, the conditions applied in the sensitivity analysis have exceeded the range of conditions that the model should be applied. The testing suggests the model will likely only be suitable for house inventories similar to those at IN2B.

Table 8-8. Parameter ranges tested for the manure belt model.

Parameter	Upper limit	Lower limit	Average Value	Increment
Ambient temperature (°C)	32	-25	15.4	0.9
Ambient relative humidity (%)	100	27	66.2	1
Inventory (birds)	338,800	0	90,522	4,600




Figure 8-10. Manure belt limitation tests for gaseous pollutants.

Visualization of the results for NH₃ (top row) and H₂S (bottom row) with tests for inventory (left), ambient temperature (center) and relative humidity (right).




Figure 8-11. Manure belt limitation tests for particulate matter.

Visualization of the results of inventory tests for PM₁₀ (top left), PM_{2.5} (top right), and TSP (bottom row).

To further explore any limitations in the models, emissions were calculated for 416,250 combination of the range of values specified in Table 8-8. Across this range of conditions, neither the NH₃, PM₁₀, nor TSP models produce negative emissions the range of conditions tested. For H₂S the model will produce negative values in instances when the ambient temperature falls below -14.2°C and house inventory is less than 33,000 birds the models can produce negative emissions values. The plots in Figure 8-12 show the maximum values of live animal weight and ambient temperature that produce negative emissions at the relative humidity specified on the x-axis, but not necessarily in combination.




Figure 8-12. Maximum values at which the manure belt house models yield negative emissions. Visualization of the results for H₂S.

8.2.3.3 Manure Shed

For the manure shed, the NH₃ and H₂S equations included ambient temperature and inventory (lagged by 5 days). The ranges tested for each parameter are in Table 8-9, with the plotted results in Figure 8-13. The plots show emissions increase with increased inventory and emissions decrease with increasing temperature. The decrease in emissions with increasing temperature could be due to increased drying of the manure. Both the NH₃ and H₂S models produce negative emissions as temperatures increase over 10°C.

PM₁₀, PM_{2.5}, and TSP models only use inventory (lagged by 5 days) to predict emissions. The range and average values used for testing are listed in Table 8-9 and the results are plotted in Figure 8-14. For all pollutants, emissions increase as airflow increases. There is a substantial change in relative sensitivity when the inventory of the supplying houses is over 620,000 birds. None of the particulate matter models produced negative emission across the range tested.

Table 8-9. Parameter ranges tested for the manure shed model.

Parameter	Upper limit	Lower limit	Average Value	Increment
Ambient temperature (°C)	32	-25	15.4	0.9
Inventory	677,600	0	90,522	10,000




Figure 8-13. Manure shed limitation tests for gaseous pollutants. Visualization of the results for NH_3 (top row) and H_2S (bottom row) with tests for inventory (left) and airflow (right).




Figure 8-14. Manure shed limitation tests for particulate matter.

Visualization of the results for PM₁₀ (top left), PM_{2.5} (top right), and TSP (bottom) for tests of airflow.

To further explore any limitations in the models, emissions were calculated for 5,625 combination of the range of ambient temperature and inventory values specified in Table 8-9. For NH₃ and H₂S, the models have an inverse relationship with emissions and inventory. As a result, as temperatures increase, there is an increased level of inventory that will produce negative values. The plots in Figure 8-15 show the maximum values of inventory that produce negative emissions at the ambient temperature specified on the x-axis.




Figure 8-15. Maximum values at which the manure belt house models yield negative emissions. Visualization of the results for NH_3 (right) and H_2S (left).

8.3 Comparison to literature

To further validate the emission models developed under this effort, EPA compared the results for the emission models to the emissions calculated using emission factors found in literature. EPA scanned the literature for a variety of emission factors for this comparison. Wood et al. (2015) contained a review of emission factors for both layer house types for NH₃. Liang et al. (2005) provided additional factors for manure belt houses. EPA selected the two most recent factors not derived from NAEMS for comparison, which are summarized separately for each house type in their respective sections.

8.3.1 High Rise House

The factors selected for comparison are listed in Table 8-7. The original units provided in Wood et al. (2015) were g yr⁻¹ AU⁻¹, based on an animal unit (AU) of 500kg. Consistent with Liang (2005), an average bird weight of 1.5 kg was used to convert AU the head (hd). For a further comparison, the emission factor included in EPA's 2001 draft AP-42 chapter is included. The emission factor was converted from the original units of the document, which were lb yr⁻¹ AU⁻¹, where AU was equivalent to 1000 birds, to kg hd⁻¹ yr⁻¹.

Table 8-10. NH₃ Emission factors for high rise houses from literature

Source	lb yr ⁻¹ AU ⁻¹	g yr ⁻¹ AU ⁻¹	kg hd ⁻¹ yr ⁻¹
EPA (2001)	28.5 a		0.285
Heber et al. (2005)		278 a	0.304
Liang et al. (2005)		298 ª	0.326

^a as reported in source.

These emission factors were then applied to the theoretical high rise houses from the previous example calculations. Comparisons were made for an inventory of 100,000 birds and 150,000 birds for both a cold weather location (Iowa) and a warm weather location (Georgia). The results are presented in Table 8-8. For both inventory levels, the emission factors from

literature fall between the estimate produced by the emission models for the two climate extremes.

Table 8-11. Comparison of resulting high rise house NH₃ emission from various estimation methods.

		NH₃ Emissions (kg yr ⁻¹)					
Meteorology	Inventory	2021	EPA	Heber et al.	Liang et al.		
site	(hd)	emission models	(2001)	(2005)	(2005)		
lowa	100,000	25,997	28,500	30,441	32,631		
Georgia	100,000	33,909	28,500	30,441	32,631		
Iowa	150,000	34,900	42,750	45,661	48,946		
Georgia	150,000	45,521	42,750	45,661	48,946		

8.3.2 Manure belt House

The emission factors selected for manure belt house model comparison are listed in Table 8-12. The original units provided in Wood et al. (2015) and Liang et al. (2005) were g yr⁻¹ AU⁻¹, based on an animal unit (AU) of 500kg. An average bird weight of 1.58 kg was used to convert AU to head (hd). Since the manure belt house was not included in EPA's 2001 draft AP-42 chapter, a third refence was included to show the range of values provided in literature. To a degree, the emission factors vary based on the removal frequency and whether the manure drying was enhanced beyond what was offered by house ventilation.

Table 8-12. NH₃ Emission factors for manure belt houses from literature

Source	Manure Management Details	g d ⁻¹ AU ⁻¹ a	kg hd ⁻¹ yr ⁻¹	
Liang et al., 2005	Removed twice a week	30.8	0.036	
Morgan et al., 2014	Removed twice a week	19.5	0.022	
Nicholson et al., 2004	Removed weekly	96	0.111	

^a as reported in source.

These emission factors were then applied to the theoretical high rise houses from the previous example calculations. Comparisons were made for an inventory of 100,000 birds and 150,000 birds for both a cold weather location (Iowa) and a warm weather location (Georgia). The results are presented in Table 8-13. Overall, the emission models presented here produce greater emissions than what has previously been reported in literature. The emission model results for the cold weather site were fairly consistent with the Nicholson et al. (2014), which was also based in a cooler climate (United Kingdom).

Table 8-13. Comparison of resulting manure belt house NH₃ emission from various estimation methods.

		NH₃ Emissions (kg yr ⁻¹)					
Meteorology	Inventory	2021	Liang et al.	Morgan et	Nicholson et al.		
site	(hd)	emission models	(2005)	al. (2014)	(2014)		
lowa	100,000	13,403	3,552	2,249	11,073		
Georgia	100,000	16,466	3,552	2,249	11,073		
lowa	150,000	16,953	5,329	3,374	16,609		
Georgia	150,000	20,873	5,329	3,374	16,609		

8.3.3 Manure Shed

EPA searches did not find sources with emission factors for manure sheds.

8.4 Replication of Independent Measurements

A final test of the developed emission models is to compare the predicted emissions to observed values from an independent study. For this test EPA obtained data from the Air Pollutant Emissions from Confined Animal Buildings (APECAB) Project. The APECAB project was conducted from the fall of 2002 through 2004 (Jacobson et. al 2011; Heber et. al 2006). Similar to NAEMS, the goal was to collect a long-term (i.e., at least a year) air pollutant information from animal feeding operations buildings. The project collected emissions data, ambient meteorological, and building parameters. Since APECAB collect many of the same parameters as NAEMS, the emission models can be applied and then compared to the observed emissions.

The APECAB project included two caged hen layer houses in Indiana. EPA was able to obtain data for this site, which included the inventory and meteorological parameters needed to estimate emission from the houses using the developed emission models. These estimates were then compared to the observed values, when available, using the same model performance statistics noted in Section 6 of the main report. The statistics for all observations are presented in Table 8-11. These statistics suggest the current model under predicts all three pollutants to some degree. The model performance statistics were also calculated for each season (Table 8-12). The season statistics show better performance during the summer and autumn for NH₃, and during the autumn for H₂S. PM₁₀ performed similarly across all seasons.

Table 8-14. Model performance evaluation statistics for high rise houses

Pollutant	n	MB (kg)	ME (kg)	NMB (%)	NME (%)	r
NH ₃	544	-164.32	204.96	-51%	64%	-0.379
H ₂ S	578	-0.049	0.244	-13%	64%	0.628
PM ₁₀	560	-7542.73	7938.27	-53%	56%	0.366

Table 8-15. Model performance evaluation statistics by season

Pollutant	season	n	MB	ME	NMB	NME	r
NH3	spring (MAM)	166	-137.85	220.95	-46%	74%	-0.49
NH3	summer (JJA)	68	6.36	99.02	3%	40%	0.55
NH3	autumn (SON)	156	-73.79	81.15	-30%	34%	0.40
NH3	winter (DJF)	154	-359.93	359.93	-79%	79%	-0.25
H2S	spring (MAM)	166	-0.11	0.29	-25%	65%	0.59
H2S	summer (JJA)	94	-0.35	0.48	-45%	61%	0.31
H2S	autumn (SON)	154	0.04	0.12	12%	39%	0.65
H2S	winter (DJF)	164	0.10	0.19	65%	116%	0.26
PM10	spring (MAM)	162	-6503.98	6734.67	-49%	50%	0.12
PM10	summer (JJA)	97	-12010.28	12010.28	-59%	59%	-0.11
PM10	autumn (SON)	135	-8742.36	9012.29	-57%	59%	0.38
PM10	winter (DJF)	166	-4970.31	5859.97	-47%	55%	-0.19

Scatter plots were also developed to present the ordered pairs with observations on the x-axis and the model predicted values on y-axis. These plots are useful for indicating trends of either over, or under prediction across the range of values. The plots include the 1:1 line (solid line) and the 1:0.5 and 1:2 lines (dashed lines). Points that fall on the 1:1 were predicted correctly, and points that fall between the 1:0.5 and 1:2 are within a factor of two observations. Good model performance would be indicated by scatter contained within a factor of two of 1:1 line, that is between the 1:0.5 and 1:2 lines. Looking for scatter confined to within a factor of two of the observation has been used as a model performance metric in air quality modeling by EPA for some time (Chang & Hanna, 2004), and continues to be included in EPA's Atmospheric Model Evaluation Tool (Appel, et al 2011) which is the current model evaluation platform.

The scatter plots were developed by season and color coded to show the performance for each house. The NH₃ scatter plots (Figure 8-16) show that a vast majority of the predicted values fall within a factor of two of the observations during summer and autumn. The scatter plot for winter (lower right) shows the model underpredicts in all instances, particularly at house B13. The H₂S scatter plots (Figure 8-17) show that a vast majority of the predicted values fall within a factor of two of the observations during all seasons. The scatter is more pronounced in spring and summer, but overall, it is reasonable model performance for these sites. The PM₁₀ scatter plots (Figure 8-18) show reasonable model performance, with most of the predicted values falling within a factor of two of the observation for all seasons, except summer. The plots also show that most of the severe underprediction occurs at house B13. Additional plots and statistics are available in Appendix H.




Figure 8-16. Scatter plot of the observed NH_3 emissions at the APECAB IN high rise site versus the emission model estimates.




Figure 8-17. Scatter plot of the observed H_2S emissions at the APECAB IN high rise site versus the emission model estimates.




Figure 8-18. Scatter plot of the observed PM_{10} emissions at the APECAB IN high rise site versus the emission model estimates.

8.5 Data Concerns

In an effort to better characterize the model performance, EPA examined the data and data collection methods to identify areas that may have contributed to poorer model performance. The following section summarizes these areas for each egg-layer source.

8.5.1 High Rise

At NC2B, there are ventilation fans at both the manure pit level and the layer room level of the houses. During NAEMS, PM concentration measurements were made at the manure pit

level. However, Li et al. (2013), which is a peer-reviewed paper that summarizes and discusses PM emissions from NC2B, describes that PM measurements at the layer room level of house 4 were also made for a period of several months. Li et al. (2013) compared PM emissions using manure pit only measurements and combined manure pit and layer room measurements and found PM emissions using the combined measurements to be 23%, 28% and 39% higher for PM_{2.5}, PM₁₀ and TSP, respectively. Differences in emissions were related to ambient temperature as ambient temperatures greater than 20°C caused the layer room fans to turn on (Li et al. 2013). Accordingly, the NAEMS NC2B PM dataset may underestimate PM emissions, particularly during warmer periods.

8.5.2 Manure belt

The particulate matter models for manure belt were difficult to develop. As noted in the Section 5.2 discussion, there are two primary influences on particulate matter emissions: 1) the amount of source material (i.e., excreted manure, feathers, and feed) and 2.) disturbance of the source materials. With respect to the first mechanism, inventory or live animal weight has been used in the models to account for the amount of source material. For an indication of when and to what extent the source material is disturbed, there is not a good indicator available in the dataset. The lack of an estimate for the effect of this component of the on emissions makes it harder to detect if other parameters have a significant influence on emissions and thus can potentially result in models with a limited number of significant parameters.

Another factor adding to the challenge of the particulate matter models was the lack of variability in the data available to develop the models. NAEMS only included one manure belt site, with two houses. These two houses have a steady inventory for the two years, except for a flock replacement event at H9. This is a problem for model development because it is hard to tease out the influence of inventory on emissions if it is roughly constant. This was not as much of an issue for the gaseous pollutants because they had approximately 20 daily observations during the flock replacement, while PM₁₀ only had 6 daily observations.

A third factor adding to the challenge was concern about the quality of the PM₁₀ measurements. The particulate matter inlet measurement is from instrumentation that resides on top of the on-farm instrument shelter (OFIS). The manure belt site (IN2B) was part of a large farm that also provided high-rise monitoring data for NAEMS (IN2H). For the manure belt house emission calculations, the inlet concentration data from the companion site (IN2H) was used, which was located on top of the IN2H OFIS between H6 and H7 (Figure 8-9). The use of this inlet concentration assumes that the high-rise house inlet concentration is representative of the manure belt house inlet concentration. The inlet PM₁₀ measurements might be unrepresentative due to the influence of nearby exhaust fan locations and local (farm scale)

meteorological conditions. The diagram from the site report (Figure 8-10), shows the inlet monitor in raceway between the two houses. The figure also shows there are exhaust fans that point outward into that same raceway. If the inlet PM concentrations are more influenced by these exhaust fans than the air entering the manure belt houses, this would result in an overestimation of inlet concentrations for the manure belt houses, particularly during periods of higher exhaust temperature and ventilation rate. This would then result in a higher frequency of negative emissions during these periods of high exhaust temperature and ventilation rates. Not only is this an issue because negative emission values are generated, but it could lead to a negative relationship between PM₁₀ emissions and exhaust temperature. The exploratory data analysis plots for IN2B H8 (See Appendix F, Figures F-85), does indicate a negative relationship with exhaust temperature.




Figure 8-19. Overhead view of IN2H/IN2B, with in the PM inlet measurement location indicated by a triangle. (Ni et al. 2010a).

Triangle indicates PM inlet concentration measurement location for both IN2H and IN2B.




Figure 8-20. Overhead (top) and side (bottom) view of sensor measurements at IN2H (Ni et al. 2010a).

8.5.3 Manure Shed

There are concerns with the quality of the manure shed data due to the methodology used to determine building inlet concentrations, exhaust concentrations, airflow and thus emissions. Each wall or ridge of the manure shed can act as an inlet or exhaust depending on the direction of the wind, therefore it is important to accurately determine average concentrations and airflow. Accordingly, the NAEMS QAPP (Heber et al. 2008) provides a reasonable methodology for determining average concentrations, airflow and thus emissions from naturally ventilated buildings. This monitoring plan was applied at the dairy barns with multiple concentration and airflow measurements made on each wall and ridge (Figure 8-21).

For the naturally ventilated manure shed at IN2B, an alternative methodology was used which involved five 2-D sonic anemometers, two impeller anemometers and one concentration measurement (Figure 8-22), which was used as the exhaust concentration when winds were generally and steadily from the west.




Figure 8-21. Overhead (a.) and side (b.) view of sensor measurements at WA5A barn (Ramirez-Dorronsoro et al. 2010).




Figure 8-22. Overhead view of sensor measurements at IN2B (Ni et al. 2010b).

Inlet concentrations for the manure shed were not measured at the manure shed. Instead, the gas inlets from the manure belt houses were used for NH₃ and H₂S, and for. For PM, the PM inlet from the high-rise inlet at IN2H was used. The accuracy and thus error associated with this alternative methodology is not known, however, there are a number of concerns associated with this alternative methodology that likely increase the error associated with the emission measurements. One concern is related to the representativeness of using inlet concentrations that were not measured at the manure shed. It is unknown how representative the high-rise house inlet and manure-belt house inlet concentrations are of the manure shed; however, it is possible that these inlet measurements are unrepresentative due to the influence of nearby exhaust fan locations and local (farm scale) meteorological conditions.

A further concern is related to the small number of airflow measurements at the various walls and ridges of the manure shed, which may not take into account the spatial variability of airflow, which can be highly variable in livestock buildings (Ogink et al. 2013). There is a similar concern regarding how well the concentration measurements used represent the spatial variability of concentrations in animal buildings, which can also be highly variable. For example, a study by Lefcourt, (2002) identified that incorrect selection of sampling locations could results in errors in gas concentrations ranging from -50% to over 200%.

Another concern with the emission measurements is due to problems with the sensors used to determine when the manure shed doors were open (Heber, personal communication), which could result in error associated with airflow measurements and thus emissions. Furthermore, at the manure shed, airflow measurements were determined using 2-D sonic anemometers. However, Ogink et al. (2013) recommends that 3-D sonic anemometers be used to measure airflows, since the direction of airflow in an opening is varied and related to ambient wind conditions. The effect of using 2-D sonic anemometers on emission measurements is not known and is likely site dependent.

9.0 SUMMARY AND CONCLUSIONS

Consistent with the Air Compliance Agreement with the AFO industry, EPA has developed emission estimation methods for NH₃, H₂S, PM₁₀, PM_{2.5}, and TSP for confinement and manure storage sources at layer operations. These draft statistical models focus on parameters that have been identified in published peer-reviewed journals as having empirical relationships with emissions. These relationships were evaluated within the NAEMS dataset before selecting parameters for emission model development. EPA also considered which variables could be measured or obtained with minimal effort.

For high rise houses, inventory was identified as a key parameter and is used in all the models as a proxy for the volume of manure generated. Temperature parameters were also identified as important variables for NH₃ and H₂S emission rates across many of the confinement house emission models. Relative humidity parameters proved to be key for particulate matter prediction, as the higher moisture levels keep house materials from entraining into the air with mechanical disruptions. Confinement parameters specific to the house, like ventilation rate and exhaust temperature, showed promise as predictive parameters. However, these parameters are not routinely measured at farms and would therefore represent an increased burden to operators should they be required for emissions estimation. As such, all of the draft high rise emission models put forward for potential future use in this document apply parameters that are already routinely collected as part of the standard farm operation (e.g., inventory and animal weight) or are ambient meteorological parameters, which are freely available from public sources such National Center for Environmental Information (NCEI, https://gis.ncdc.noaa.gov/maps/), and can be easily incorporated into an emission estimation tool for users.

For manure belt houses, inventory was identified as a key parameter and is used in all the models as a proxy for the volume of manure generated. Temperature parameters were also identified as important variables for NH₃ and H₂S emission rates across many of the confinement house emission models. For particulate matter, most tested models contained parameters that, while found in literature to have a relationship with emissions, were statistically insignificant when tested. The established development process produced only two combinations of models that were composed entirely of statistically significant parameters. One of the two models included exhaust temperature, which is not necessarily retained by producers. The manure shed models had limited statistically significant models for all pollutants. Airflow was the key parameter in predicting emissions for both gaseous pollutants and particulate matter. However, airflow is not routinely measured for manure sheds, and can be particularly difficult to estimate for naturally ventilated structures. For the manure belt house and manure shed models, EPA considered overlooking the significance calculations and selecting models purely based on the

relationships established in literature. All candidate models tested appear in Appendix F for review and consideration during this comment period.

Overall, the method used to develop the emission models allows for the incorporation of additional emissions and monitoring datasets from other studies, should they become available after the release of the emission models. Revised emission models for any individual farm type could be issued once significant additional data becomes available. Similarly, if monitoring options for house or manure shed parameters become more widespread as automation options grow, future evaluations could assess whether emission models should be developed to include these parameters.

EPA recognizes the scientific and community desire for process-based models. The data collected during NAEMS and the emission models developed here lay the groundwork for developing these more process-related emission estimates. EPA supports the future development of process-based models which account for the entire animal feeding process. While the interim statistical models allow estimation of emissions from various categories of layer operations across the U.S., process-based models would allow producers to estimate the impacts of different best management practices to reduce air emissions, helping to incentivize change.

10.0 REFERENCES

Aarnink, A.J.A, & Ellen, H.H. (2007). Processes and factors affecting dust emissions from livestock production. In: *Proceedings of the DustConf 2007. How to Improve Air Quality*. Maastricht, The Netherlands.

Appel, K.W., Gilliam, R.C., Davis, N., Zubrow, A., and Howard, S.C. (2011). Overview of the Atmospheric Model Evaluation Tool (AMET) v1.1 for evaluating meteorological and air quality models, Environ. Modell. Softw., 26(4), 434-443.

Arogo, J., Zhang, R.H., Riskowski, G.L., Christianson, L.L., & Day, D.L. (1999). Mass transfer coefficient of ammonia in liquid swine manure and aqueous solutions. *Journal of Agricultural Engineering Research* 73, 77-86.

Cambra-López, M., Aarnink, A.J.A., Zhao, Y., Calvet, S., & Torres, A.G. (2010). Airborne particulate matter from livestock production systems: A review of an air pollution problem. *Environmental Pollution 158*, 1-17.

Cambra-López, M., Torres, A.G., Aarnink, A.J.A., & Ogink, N.W.M. (2011). Source analysis of fine and coarse particulate matter from livestock houses. *Atmospheric Environment* 45(3), 694-707.

Chang, J., Hanna, S. Air quality model performance evaluation. Meteorol Atmos Phys 87, 167–196 (2004). https://doi.org/10.1007/s00703-003-0070-7

Christensen J.R., Nash, M.S., Chaloud D., and Pitchford A. 2016. Spatial Distribution of Small Water Body Types Across Indiana Ecoregions. *Ecohydrology*. 9,122-137. Available at: https://cfpub.epa.gov/si/si public record report.cfm?dirEntryId=308263&Lab=NERL

Cortus, E.L., Lin X-J., Zhang R., and Heber A.J. (2010). National Air Emissions Monitoring Study: Emissions Data from Two Layer Houses in California - Site CA2B. Final Site Report (FSR). Purdue University, West Lafayette, Indiana, July 2. [Online]. Available at: https://archive.epa.gov/airquality/afo2012/web/pdf/ca2bsummaryreport.pdf. [Accessed Mar. 2017].

Feldman I., and Tibor T. 1996. ISO 14000: A Guide to the New Environmental Management Standards. Irwin Professional Pub.

Grant, R.H. (2008). Quality assurance project plan for the National Air Emissions Monitoring Study (open source emissions component), Ver. 4. Purdue University.

Grant, R.H., Boehm, M.T., Lawrence, A.F., & Heber, A.J. (2013). Ammonia emissions from anaerobic treatment lagoons at sow and finishing farms in Oklahoma. *Agricultural and Forest Meteorology 180*, 203-210.

Groot Koerkamp, P.W. (1994). Review on emissions of ammonia from housing systems for laying hens in relation to sources, processes, building design and manure handling. *Journal of Agricultural Engineering Research* 59, 73-87.

- Hanni S.M, and Bogan B.W. 2008. Manure Sampling Standard Operating Procedure (SOP) M1. Rev 1.0. 2008. Purdue Ag Air Quality Lab.
- Heber, A.J., J.Q. Ni, T.T. Lim, R. Chervil, P.C. Tao, Jacobson, Larry D, S.J. Hoff, Y. Zhang, J.A. Koziel, D.S. Beasley, J.M. Sweeten. 2005. Air pollutant emissions from two high-rise layer barns in Indiana Proc. Annual Conference and Exhibition of the Air and Waste Management Association, Air and Waste Management Association, Pittsburgh, PA (2005)
- Heber A.J., Ni, J-Q., Lim T.T., Tai, P.C.; Schmidt A.M, Koziel, J.A., Beasley, D.B., Hoff S.J., Nicolai, R.E., Jacobson, L.D., & Zhang, Y. 2006. Quality Assured Measurements of Animal Building Emissions: Gas Concentrations, Journal of the Air & Waste Management Association, 56:10, 1472-1483, DOI: 10.1080/10473289.2006.10465680
- Heber A.J., Ni, J-Q., Ramirez, J.C., Schrock, W., & Elkins, J. (2008). *Quality assurance project plan for the National Air Emissions Monitoring Study (barn component)*. Purdue University.
- Heber A.J., Ni J-Q., Bogan B.W., Cortus E.L., Kilic, I., Zhang R., and Wang-Li L. 2012. Evaluation and Analysis of NAEMS Layer Data. Industry Report. Purdue University, West Lafayette, Indiana.
- Jacobson, L.D.; Hetchler, B.P.; Akdeniz, N.; Hoff, S.J.; Heber, A.J.; Ni, J.Q.; Zhang, Y.; and Koziel, J.A. 2011. Air Pollutant Emissions from Confined Animal Buildings (APECAB) Project Summary. *Agricultural and Biosystems Engineering Publications*. 72. https://lib.dr.iastate.edu/abe_eng_pubs/72
- Joo, H.S., Ndegwa, P.M., Heber, A.J., Bogan, B.W., Ni, J.-Q., Cortus, R.L., & Ramirez-Dorronsoro, J.C. (2014). A direct method of measuring gaseous emissions from naturally ventilated dairy barns. *Atmospheric Environment* 86, 176-186.
- Leeden (Van Der), Meijer R.E. and Busing F.M. 2007. Resampling Multilevel Models. Ch. 11. *Handbook of Multilevel Analysis*. Springer, New York.
- Lefcourt, A.M., 2002. Some potential problems for measuring ammonia emissions from farm structures. Transactions of the ASAE, 48 (1), 367-372.
- Li Q-F., Wang-Li L., Bogan B.W, Wang K., Chai L., Ni J-Q., and Heber A.J. 2013. The National Air Emissions Monitoring Study's Southeast Layer Site: Part IV. Effects of Farm Management. *Trans. ASABE*. 56(3): 1199-1209.
- Liang, Y., Xin, H., Wheeler, E.F., Gates, R.S., Li, H., Zajaczkowski, J.S., Topper, P.A., Casey, K.D., Behrends, Burnham, D.J., & Zajaczkowski, F.J. (2005). Ammonia emissions from U.S. laying hen houses in Iowa and Pennsylvania. *Transactions of the ASAE* 48(5): 1927-1941.

Liang, W-Z. 2015. Ammonia Emission Model for High-Rise Layer Houses. Doctoral Dissertation. Department of Agricultural & Biological Engineering. Purdue University, Graduate School. West Lafayette, Indiana.

Lim, T. T., Heber, A. J., Ni, J. Q., Gallien, J. X., & Xin, H. (2003). Air quality measurements at a laying hen house: Particulate matter concentrations and emissions. In Air Pollution from Agricultural Operations-III (p. 249). American Society of Agricultural and Biological Engineers.

Lin X-J., Cortus E.L., Zhang R., Jiang S., and Heber A.J. 2012a. Ammonia, Hydrogen Sulfide, Carbon Dioxide and Particulate Matter Emissions from California High-Rise Layer Houses. *Atmos. Environ.* (46) 81-91

Lin X-J., Cortus E.L., Zhang R., Jiang S., Heber A.J., and Kilic I. 2012b. Thermal Environmental Control of High-Rise Layer Houses in California. *Trans. ASABE*. 55(5): 1909-1920.

Mackie R.I., Stroot P.G., & Varel V.H. (1998). Biochemical identification and biological origin of key odor components in livestock waste. *Journal of Animal Science* 76, 1331-1342.

Montes, F., Rotz, C. A., & Chaoui, H. (2009). Process modeling of ammonia volatilization from ammonium solution and manure surfaces: A review with recommended models. *Transactions of the ASABE* 52(5), 1707-1719.

Morgan, R.J., Wood, D.J., Van Heyst, B.J., 2014. The development of seasonal emission factors from a Canadian commercial laying hen facility. Atmos. Environ. 86, 1e8.

Ni, J. (1999). Mechanistic models of ammonia release from liquid manure: a review. *Journal of Agricultural Engineering Research* 72, 1-17.

Ni, J-Q., Heber A.J., Sutton A.L., & Kelly D.T. (2009). Mechanisms of gas releases from swine wastes. *Transactions of the ASABE* 52(6), 2013-2025.

Ni, J-Q., Diehl C.A., Lim T.T., Bogan B.W., Chen L., Chai L., and Heber A.J. 2010a. National Air Emissions Monitoring Study: Emissions Data from Two High-Rise Layer Houses in Indiana - Site IN2H. Final Site Report (FSR). Purdue University, West Lafayette, Indiana, July 30. [Online]. Available at:

https://archive.epa.gov/airquality/afo2012/web/pdf/in2hsummaryreport.pdf. [Accessed Mar. 2017].

Ni, J-Q., Diehl C.A., Chai L., Bogan B.W., Cortus E.L., Lim T.T., and Heber A.J. 2010b. National Air Emissions Monitoring Study: Emissions Data from Two Manure Belt Layer Houses in Indiana - Site IN2B. Final Report. Purdue University, West Lafayette, IN, July 2. Available at: https://archive.epa.gov/airquality/afo2012/web/pdf/in2bsummaryreport.pdf

Ni J-Q., Liu S., Diehl C.A., Lim T.T, Bogan B.W., Lide C., Chai L., Wang K., and Heber A.J. 2017a. Emission Factors and Characteristics of Ammonia, Hydrogen Sulfide, Carbon Dioxide, and Particulate Matter at Two High-Rise Layer Hen Houses. *Atmos. Environ.* 154: 260-273.

Ni, J.-Q., Diehl, C.A., Chai, L., Chen, Y., Heber, A.J., Lim, T.T., Bogan, B.W. 2017b. Factors and characteristics of ammonia, hydrogen sulfide, carbon dioxide, and particulate matter emissions from two manure-belt layer hen houses. *Atmospheric Environment* 156, 113-124.

Nicholson, F. A., B. J. Chambers, and A. W. Walker. 2004. Ammonia emissions from broiler litter and laying hen manure management systems. *Biosystems Eng.* 89(2): 175-185.

Ogink, N.W.M, Mosquera, J. Calvet, S., Zhang, 2013. Methods for measuring gas emissions from naturally ventilated livestock buildings: Developments over the last decade and perspectives for improvement. Biosystems Engineering 116, 297-308.

Ramirez-Dorronsoro, J.C., Joo, H.S., Ndegwa, P., Heber, A.J., 2010. National Air Emissions Monitoring Study: Data from Two Dairy Freestall Barns in Washington WA5B, Final Report. Purdue University, West Lafayette, IN, July 30.

Rumsey I.C., & Aneja V.P. (2014). Measurement and modeling of hydrogen sulfide lagoon emissions from a swine concentrated animal feeding operation. *Environmental Science and Technology* 48, 1609-1617.

Shepherd, T.A., Zhao, Y., Li, H., Stinn, J.P., Hayes, M.D., & Xin, H., (2015). Environmental assessment of three egg production systems – Part II. Ammonia, greenhouse gas and particulate matter emissions. *Poultry Science* 94, 534-543.

SOP A10. 2009. Calculating Airflow and Emissions From Naturally-Ventilated Barns. Standard Operating Procedure A10, v. 0.0. Purdue Ag Air Quality Lab.

Takai H., Pedersen S., Johnsen J.O., Metz J.H.M., Groot Koerkamp P.W.G., Uenk G.H., Phillips V.R., Holden M.R., Sneath R.W., Short, J.L., White, R.P., Hartung, J., Seedorf J., Schroder M., Linkert K.H., & Wathes C.M. (1998). Concentrations and emissions of airborne dust in livestock buildings in Northern Europe. *Journal of Agricultural Engineering Research* 70(1), 59-77.

Tong, X., Zhao, L., Heber, A.J., Ni, J.-Q. (2020). Mechanistic modelling of ammonia emission from laying hen manure at laboratory scale. *Biosystems Engineering* 192, 24-41.

USDA NASS, 2017 Census of Agriculture, Ag Census Web Maps. Available at: www.nass.usda.gov/Publications/AgCensus/2017/Online_Resources/Ag_Census_Web_Maps/Overview/.

U.S. EPA (U.S. Environmental Protection Agency). 2001a. Emissions from Animal Feeding Operations (Draft). EPA Contract No. 68-D6-0011.

U.S. EPA Science Advisory Board (SAB), 2013. SAB Review of Emissions-Estimating Methodologies for Broiler Animal Feeding Operations and for Lagoons and Basins at Swine and Dairy Animal Feeding Operations, EPA-SAB-13-003.

Wall E., Weersink A., and Swanton C. 2001. Agriculture and ISO 14000. *Food Policy*, 26: 35-48.

Wang K., Kilic I., Li, Wang L., Bogan, B.W., Ni J-Q., Chai L., and A.J. Heber. 2010. National Air Emissions Monitoring Study: Emissions Data from Two Tunnel-Ventilated Layer Houses in North Carolina - Site NC2B. Final Site Report (FSR). Purdue University, West Lafayette, Indiana, June 18. [Online]. Available at:

https://archive.epa.gov/airquality/afo2012/web/pdf/nc2bsummaryreport.pdf. [Accessed Feb. 2017].

Wang-Li L., Li Q-F., Wang K., Bogan B.W., Ni J-Q., Cortus E.L., and Heber A.J. 2013a. The National Air Emissions Monitoring Study's Southeast Layer Site: Part I. Site Characteristics and Monitoring Methodology. *Trans. ASABE*. 56(3): 1157-1171.

Wang-Li L., Li Q-F., Chai L., Cortus E.L., Wang K., Kilic I., Bogan B.W., Ni J-Q., and Heber A.J. 2013b. The National Air Emissions Monitoring Study's Southeast Layer Site: Part III. Ammonia Concentrations and Emissions. *Trans. ASABE*. 56(3): 1185-1197.

Wang K-Y., Li Q-F., Wang-Li L., Cortus E.L., Bogan B.W., Kilic I., Liang W-Z., Xiao C.H., Chai L.L., Ni J-Q., and Heber A.J. 2016. The National Air Emissions Monitoring Study's Southeast Layer Site: Part V. Hydrogen Sulfide and Volatile Organic Compounds. *Trans. ASABE*. 59(2): 681-693.

Weaver W.D., & Meijerhof, R. (1991). The effect of different levels of relative humidity and air movement on litter conditions, ammonia levels, growth, and carcass quality for broiler chickens. *Poultry Science* 70, 746-755.

Wood et al., 2015. D. Wood, S. Cowherd, B. Van Heyst A summary of ammonia emission factors and quality criteria for commercial poultry production in North America. Atmos. Environ., 115 (2015), pp. 236-245

Wu-Haan, W., Powers, W. J., Angel, C. R., Hale III, C. E., & Applegate, T. J. (2007). Effect of an acidifying diet combined with zeolite and slight protein reduction on air emissions from laying hens of different ages. Poultry Science, 86(1), 182-190.

Xin, H., Gates, R.S., Green, A.R., Mitloehner, F.M., Moore Jr, P.A., & Wathes, C.M. (2011). Environmental impacts and sustainability of egg production systems. *Poultry Science* 90, 263-277.

Zhao, Y., Xin, H., Shepherd, T.A., Haye, M.D., Stinn, J.P., & Li, H. (2013). Thermal environment, ammonia concentrations and ammonia emissions of aviary houses with white laying hens. *Transactions of the ASABE* 56(3), 1145-1156.

Appendix A - Particulate Matter Sampling Schedu	le

List of Tables

Table A-1. PM Sampling Schedule CA2B	A-3
Table A-2. PM Sampling Schedule IN2H	A-4
Table A-3. PM Sampling Schedule IN2B (houses and manure shed)	A-5
Table A-4. PM Sampling Schedule NC2B	A-6

Table A-1. PM Sampling Schedule CA2B

Start Date	End Date	Duration (days)				
Start Date	End Date	PM ₁₀	TSP	PM _{2.5}		
10/17/07	12/14/07	58.6	NS	NS		
12/14/07	12/21/07	NS	7.0	NS		
12/21/07	01/07/08	16.9	NS	NS		
01/07/08	01/23/08	NS	NS	15.9		
01/23/08	03/14/08	51.0	NS	NS		
03/14/08	03/21/08	NS	7.0	NS		
03/21/08	05/20/08	60.0	NS	NS		
05/20/08	05/27/08	NS	6.9	NS		
05/27/08	07/07/08	40.9	NS	NS		
07/07/08	07/22/08	NS	NS	14.9		
07/22/08	09/19/08	59.0	NS	NS		
09/19/08	09/29/08	NS	10.1	NS		
09/29/08	01/09/09	102.1	NS	NS		
01/09/09	01/28/09	NS	NS	18.9		
01/28/09	04/14/09	75.9	NS	NS		
04/14/09	04/22/09	NS	7.9	NS		
04/22/09	07/09/09	78.1	NS	NS		
07/09/09	07/10/09	NS	0.9	NS		
07/10/09	07/29/09	18.9	NS	NS		
07/29/09	08/11/09	NS	12.9	NS		
08/11/09	10/16/09	65.5	NS	NS		

NS = Not sampled.

Table A-2. PM Sampling Schedule IN2H

		Duration (days) H6			Dura	tion (da	ys) H7
Start Date	Stop Date	PM ₁₀	TSP	PM _{2.5}	PM ₁₀	TSP	PM _{2.5}
05/09/07	07/03/07	54	NS	NS	NS	NS	NS
05/09/07	07/12/07	NS	NS	NS	63	NS	NS
07/03/07	07/12/07	NS	NS	9	NS	NS	NS
07/12/07	07/18/07	NS	NS	NS	NS	NS	6
07/12/07	09/13/07	61	NS	NS	NS	NS	NS
07/18/07	09/13/07	NS	NS	NS	55	NS	NS
09/13/07	09/19/07	NS	NS	6	NS	NS	6
09/19/07	11/08/07	49	NS	NS	49	NS	NS
11/08/07	11/20/07	NS	NS	12	NS	NS	12
11/20/07	01/01/08	41	NS	NS	41	NS	NS
01/01/08	01/31/08	30	NS	NS	NS	NS	NS
01/01/08	04/04/08	NS	NS	NS	93	NS	NS
01/31/08	02/08/08	NS	8	NS	NS	8	NS
02/08/08	04/04/08	56	NS	NS	NS	NS	NS
04/04/08	04/11/08	NS	7	NS	NS	7	NS
04/11/08	04/18/08	NS	NS	7	NS	NS	7
04/18/08	06/19/08	61	NS	NS	NS	NS	NS
04/18/08	12/12/08	NS	NS	NS	235	NS	NS
06/19/08	06/27/08	NS	8	NS	NS	NS	NS
06/27/08	12/12/08	165	NS	NS	NS	NS	NS
12/12/08	12/18/08	NS	6	NS	NS	6	NS
12/18/08	01/01/09	13	NS	NS	13	NS	NS
01/01/09	02/06/09	35	NS	NS	35	NS	NS
02/06/09	02/13/09	NS	7	NS	NS	7	NS
02/13/09	04/09/09	NS	NS	56	NS	NS	56
04/09/09	04/16/09	NS	7	NS	NS	7	NS
04/16/09	06/04/09	48	NS	NS	48	NS	NS
06/04/09	06/12/09	NS	8	NS	NS	8	NS
06/12/09	06/26/09	NS	NS	14	NS	NS	14
06/26/09	07/30/09	34	NS	NS	34	NS	NS

NS = Not sampled.

Table A-3. PM Sampling Schedule IN2B (houses and manure shed)

Start Data	Ford Data	Duration (days)				
Start Date	End Date	PM ₁₀	TSP	PM _{2.5}		
1/1/08	1/30/08	29	NS	NS		
1/30/08	2/8/08	NS	8	NS		
2/8/08	4/4/08	56	NS	NS		
2/13/09	3/6/09	NS	NS	23		
4/4/08	4/11/08	NS	7	NS		
4/11/08	4/18/08	NS	NS	7		
4/18/08	6/19/08	61	NS	NS		
6/12/09	6/26/09	NS	NS	14		
6/19/08	6/27/08	NS	8	NS		
6/26/08	8/14/08	48	NS	NS		
6/27/08	10/3/08	96	NS	NS		
10/3/08	10/9/08	NS	6	NS		
10/9/08	12/12/08	63	NS	NS		
12/12/08	12/18/08	NS	6	NS		
12/18/08	2/6/09	48	NS	NS		
2/6/09	2/13/09	NS	7	NS		
3/4/09	4/9/09	35	NS	NS		
4/9/09	4/17/09	NS	8	NS		
4/17/09	6/4/09	47	NS	NS		
6/4/09	6/12/09	NS	8	NS		
8/14/09	8/20/09	NS	6	NS		
8/20/09	10/9/09	49	NS	NS		
10/9/09	10/16/09	NS	7	NS		
10/16/09	1/16/10	90	NS	NS		

NS = Not sampled.

Table A-4. PM Sampling Schedule NC2B

Start Data	Stop Data	Duration (days)			
Start Date	Stop Date	PM ₁₀	TSP	PM _{2.5}	
09/24/07	01/16/08	114.5	NS	NS	
01/16/08	02/04/08	NS	NS	18.9	
02/04/08	03/26/08	51.1	NS	NS	
03/26/08	04/04/08	NS	9.1	NS	
04/04/08	05/12/08	37.9	NS	NS	
05/12/08	05/28/08	NS	16.0	NS	
05/28/08	08/07/08	71.1	NS	NS	
08/07/08	08/21/08	NS	14.0	NS	
08/21/08	08/21/08	NS	20.9**	NS	
08/21/08	09/11/08	20.9*	NS	NS	
09/11/08	10/17/08	36.0	NS	NS	
10/17/08	10/23/08	NS	5.8	NS	
10/23/08	10/24/08	0.9	NS	NS	
10/24/08	10/30/08	NS	NS	6.0	
10/30/08	01/09/09	71.2	NS	NS	
01/09/09	01/15/09	NS	5.8	NS	
01/15/09	02/26/09	42.0	NS	NS	
02/26/09	02/27/09	1 [§]	NS	NS	
02/27/09	03/04/09	NS	4.9 [§]	NS	
03/04/09	04/02/09	28.9 ^{§§}	NS	NS	
04/02/09	04/10/09	NS	8.0	NS	
04/10/09	06/04/09	55.1	NS	NS	
06/04/09	06/11/09	NS	7.0	NS	
06/11/09	07/24/09	43.0	NS	NS	
07/24/09	08/06/09	NS	NS	12.8	
08/06/09	08/06/09	NS	NS	10.9**	
08/06/09	08/07/09	1.0†	NS	NS	
08/07/09	08/17/09	10.2*	NS	NS	
08/17/09	08/20/09	2.9	NS	NS	
08/20/09	08/27/09	NS	6.8	NS	
08/27/09	09/15/09	19.2	NS	NS	
09/15/09	09/15/09	20.9*	NS	NS	
09/15/09	09/22/09	NS	6.8**	NS	
09/22/09	10/06/09	14.2**	NS	NS	

NS = Not sampled.; *All except inlet; **Only inlet

†Only H4 upstairs; ‡All except H3 §H4 TEOM collocated with H3 §§H3 TEOM relocated to H4 upstairs **Appendix B - Data Processing**

List of Tables
Table B-1. Summary of the effect of applying the negative emission cut-off to layer high rise data B-4
Table B-2. Summary of the effect of applying the negative emission cut-off to layer manure belt house data. B-4
Table B-3. Summary of the effect of applying the negative emission cut-off to layer manure shed data.B-4

1.0 NEGATIVE EMISSION VALUE ASSESSMENT METHODOLOGY

Negative calculated emission values can occur in NAEMS datasets due to a range of different scenarios as described in the SAB review of the 2012 EEMs developed by EPA (U.S. EPA SAB, 2013). A summary of these scenarios and whether SAB recommended the data should be retained or removed is provided below:

- 1. A calculation bias may occur when measured values are at or close to the detection limit, or negative. This scenario should result in small negative values, which should be retained.
- 2. In NAEMS, the background and source measurements were measured either intermittently (twice a day for gas), or continuously without correction for lag time in the barn (PM data), thus leading to a bias either up or down, introducing the potential for negative emission values. Negative emission values should be retained because this bias could occur in either the positive or negative direction.
- 3. Outdoor events may affect background and barn concentrations. For example, if there was activity outside an animal barn which resulted in increased pollutant concentration (e.g., manure cleanout of another barn)), the measured background values would create a negative bias. Alternatively, a positive bias could occur if meteorological conditions caused the barn exhaust air to return into the barn, thus affecting measured barn concentrations.

To avoid bias from the true value, the SAB suggests keeping calculated values from scenario 1 and 2 and removing values identified to be caused by scenario 3, however the NAEMS did not record outdoor events that may affect background concentration (scenario 3), therefore it could not be determined if negative emissions were caused by scenario 2 or 3. It is likely that scenarios 1 and 2 result in smaller negative (closer to zero) emissions than scenario 3. Therefore, a methodology was developed to remove large negative emissions likely associated with scenario 3. In the NAEMS QAPP, the gas and PM barn emission uncertainty were determined to be $\pm 27\%$ and $\pm 32\%$ for mechanically ventilated barns and $\pm 50\%$ and $\pm 53\%$ for naturally ventilated barns (Heber et al. 2008). Cut-offs for valid negative data were therefore determined for each pollutant by multiplying the emission uncertainty by the median of the positive measured emission values.

Table B-1. Summary of the effect of applying the negative emission cut-off to layer high rise data.

	Median		Negative	# Of Negative Emission Values		
	Positive		Emission	Before	Removed	After
	Emission	Uncertainty	Cut-Off	Cut-Off	Due To	Cut-Off
Pollutant	(kg d ⁻¹ / g d ⁻¹) ^a	(%)	(kg d ⁻¹ / g d ⁻¹) ^a	Applied	Cut-Off	Applied
NH ₃	55.19	27	-14.90	4	0	4
H ₂ S	51.78	27	-13.98	89	43	46
PM ₁₀	1724.00	32	-551.68	54	22	32
PM _{2.5}	113.50	32	-36.32	65	50	15
TSP	3614.80	32	-1156.74	0	0	0

^a NH₃ emissions in units of kg day⁻¹, all other pollutants in units of g day⁻¹

Table B-2. Summary of the effect of applying the negative emission cut-off to layer manure belt house data.

	Median		Negative	# Of Nega	# Of Negative Emission Valu			
	Positive		Emission	Before	Removed	After		
	Emission	Uncertainty	Cut-Off	Cut-Off	Due To	Cut-Off		
Pollutant	(kg d ⁻¹ / g d ⁻¹) ^a	(%)	(kg d ⁻¹ / g d ⁻¹) ^a	Applied	Cut-Off	Applied		
NH ₃	59.67	27	-16.11	0	0	0		
H ₂ S	440.89	27	-119.04	8	0	8		
PM ₁₀	3429.35	32	-1097.39	88	46	42		
PM _{2.5}	343.2	32	-109.82	30	22	8		
TSP	8769.45	32	-2806.22	1	0	1		

^a NH₃ emissions in units of kg day⁻¹, all other pollutants in units of g day⁻¹

Table B-3. Summary of the effect of applying the negative emission cut-off to layer manure shed data.

	Median		Negative	# Of Negative Emission Values		
	Positive		Emission	Before	Removed	After
	Emission	Uncertainty	Cut-Off	Cut-Off	Due To	Cut-Off
Pollutant	(kg d ⁻¹ / g d ⁻¹) ^a	(%)	(kg d ⁻¹ / g d ⁻¹) ^a	Applied	Cut-Off	Applied
NH ₃	2.82	50	-1.41	0	0	0
H ₂ S	19.28	50	-9.64	5	1	4
PM ₁₀	103.16	53	-54.67	85	40	45
PM _{2.5}	15.41	53	-8.16	0	0	0
TSP	191.78	53	-101.64	3	1	2

 $^{^{\}rm a}$ NH $_{\rm 3}$ emissions in units of kg day $^{\rm -1}$, all other pollutants in units of g day $^{\rm -1}$

2.0 REFERENCES

Heber A.J., Ni J-Q., Ramirez J.C., Schrock W., and Elkins J. 2008. Quality assurance project plan for the National Air Emissions Monitoring Study (barn component). Purdue University, West Lafayette, IN.

U.S. EPA Science Advisory Board (SAB), 2013. SAB Review of Emissions-Estimating Methodologies for Broiler Animal Feeding Operations and for Lagoons and Basins at Swine and Dairy Animal Feeding Operations, EPA-SAB-13-003.

Appendix C - Data Completeness

List of Tables

Table C-1. Number of ADM for high rise house NH3 emissions at varying percentages of data completeness	C-4
Table C-2. Number of ADM for high rise house H ₂ S emissions at varying percentages of data completeness	C-4
Table C-3. Number of ADM for high rise house PM ₁₀ emissions at varying percentages of data completeness	C-5
Table C-4. Number ADM for high rise house PM _{2.5} emissions at varying percentages of data completeness	C-5
Table C-5. Number of ADM for high rise house TSP emissions at varying percentages of data completeness	C-6
Table C-6. Number of manure belt house ADM for NH ₃ at varying percentages of data completeness	C-6
Table C-7. Number of manure belt house ADM for H ₂ S at varying percentages of data completeness	C-7
Table C-8. Number of manure belt house ADM for PM ₁₀ at varying percentages of data completeness	C-7
Table C-9. Number of manure belt house ADM for PM _{2.5} at varying percentages of data completeness	C-7
Table C-10. Number of manure belt house ADM for TSP at varying percentages of data completeness	C-7
Table C-11. Number of manure shed ADM for each pollutant at varying percentages of data completeness	C-8

Data Completeness Criteria for the Revised Data Set

The appropriate data completeness criteria to use in a study depends on the size of the dataset and the accuracy needed. A study by Grant et al. (2013), in which NH₃ emissions were modeled from swine lagoons based on NAEMS data, investigated data completeness and associated accuracy. The swine lagoon NH₃ emissions dataset had limited data availability at a data completeness of 75%. Grant et al. (2013) explored how much the data completeness criteria could be relaxed but still result in data with acceptable error. The study suggested an error of $\pm 25\%$ to be acceptable and determined that a daily data completeness of 52% (or 25 out of 48 30-minute periods) gave less than $\pm 25\%$ error (see Figure B-1). Using this relaxed daily completeness criteria resulted in a substantial increase in the size of the dataset.

Based on Figure B-1 from the Grant et al. (2013) study, it can be observed that a daily completeness criterion of 75% (36 out of 48 30-minute periods) would give an error of approximately 10%. If it is assumed that the relationship between data completeness and error from the Grant et al. (2013) study is representative of other NAEMS datasets, the effect of relaxed data completeness criteria can be investigated for other NAEMS sources.

The following sections examine the effect of a reduced data completeness criterion on the number of valid average daily means (ADM) for both the layer barns and manure shed, based on additional analysis completed by Heber that examined the effect of different completeness criteria by comparing the number of valid ADM.

EPA reviewed this data for the egg-layer sites and retained the 75% completeness criterion for all sites. The full analysis can be found in Appendix B.




Figure C-1. Ratio of mean predicted emissions for portion of day with valid emissions measurements to mean predicted emissions for the complete day at the finishing (A) and sow (B) farm. Error plotted against number of valid 30-minute measurements (from Grant et al.

Data Completeness Review and Conclusions for the High Rise Dataset

The number of average daily means (ADM) for NH₃ emissions at varying percentages of data completeness for the revised data set are shown in Table B-1. For the layer high rise data set in this study, decreasing the daily completeness criteria from 75% to 50% would increase the number of valid days by 230 (6.4%), but based on the Grant et al. (2013) study there would be an approximate 15% increase in error. Therefore, based on this analysis, a daily completeness criterion of 75% was chosen for the NH₃ layer high rise revised data set. This value matches the data completeness criteria used in the 2010 NAEMS data sets (Grant et al. 2008; Heber et al. 2008).

Table C-1. Number of ADM for high rise house NH3 emissions at varying percentages of data completeness.

% Valid												
Data	0	10	20	30	40	50	60	70	75	80	90	100
IN2H-H6	680	677	671	663	648	628	608	579	564	558	335	335
IN2H-H7	687	684	679	668	655	638	616	588	570	558	374	374
CA2B-H5	606	606	601	601	596	591	581	578	570	567	564	470
CA2B-H6	628	625	618	618	613	607	602	596	590	585	580	412
NC2B-H3	690	690	683	682	678	673	665	645	638	629	614	410
NC2B-H4	684	684	675	674	667	664	656	647	639	629	616	392
Total	3,975	3,966	3,927	3,906	3,857	3,801	3,728	3,633	3,571	3,526	3,083	2,393

For H₂S, the number of ADM at varying percentages of data completeness for the revised data set are shown in Table B-2. For the high rise data set, decreasing the daily completeness criteria from 75% to 50% would increase the number of valid days by 195 (5.8%), but based on the Grant et al. (2013) study there would be an approximate 15% increase in error. Since the small increase in the number of ADM values does not justify the 15% increase in error, a daily completeness criterion of 75% was chosen for the revised H₂S high rise data set.

Table C-2. Number of ADM for high rise house H₂S emissions at varying percentages of data completeness.

% Valid												
Data	0	10	20	30	40	50	60	70	75	80	90	100
IN2H-H6	482	479	474	470	460	447	434	413	401	398	248	248
IN2H-H7	489	486	482	475	470	458	445	426	410	401	264	264
CA2B-H5	646	635	630	630	627	622	610	607	601	597	594	495
CA2B-H6	676	654	648	648	644	638	631	625	620	614	610	466
NC2B-H3	713	713	709	708	702	698	688	670	662	656	650	444
NC2B-H4	706	706	701	700	693	689	680	669	663	654	650	423
Total	3,712	3,673	3,644	3,631	3,596	3,552	3,488	3,410	3,357	3,320	3,016	2,340

For PM, the number of ADM at varying percentages of data completeness for the revised data set are shown in Table B-3Error! Reference source not found., B-4 and B-5 for PM₁₀, PM_{2.5} and TSP, respectively. For the high rise site data set, decreasing the daily completeness criteria from 75% to 50% would increase the number of valid days by 151 (5.4 %) for PM₁₀, 17 (10.5%) for PM_{2.5} and 11 (5.0%) for TSP, respectively. Again, the small increase in the number of ADM values does not justify the 15% increase in error. Therefore, a daily completeness criterion of 75% was chosen for the all the PM species for high rise data set. This value also matches the data completeness criteria used in the 2010 NAEMS data sets (Heber et al. 2008).

Table C-3. Number of ADM for high rise house PM₁₀ emissions at varying percentages of data completeness.

% Valid Data	0	10	20	30	40	50	60	70	75	80	90	100
IN2H-H6	489	484	479	472	462	451	439	423	417	404	200	200
IN2H-H7	479	476	472	465	458	450	443	425	421	409	222	222
CA2B-H5	492	492	487	487	482	467	457	454	451	450	439	347
CA2B-H6	593	592	581	581	575	559	539	531	527	525	501	306
NC2B-H3	450	450	448	446	442	429	423	411	410	406	391	283
NC2B-H4	603	603	601	596	587	572	565	555	551	547	532	328
Total	3,106	3,097	3,068	3,047	3,006	2,928	2,866	2,799	2,777	2,741	2,285	1,686

Table C-4. Number ADM for high rise house PM_{2.5} emissions at varying percentages of data completeness.

% Valid Data	0	10	20	30	40	50	60	70	75	80	90	100
IN2H-H6	24	23	22	19	19	19	18	17	16	15	11	11
IN2H-H7	18	17	17	14	14	14	12	11	9	9	7	7
CA2B-H5	48	48	46	46	44	43	40	40	40	40	38	29
CA2B-H6	53	53	52	52	48	47	45	44	43	43	41	29
NC2B-H3	26	26	26	25	23	22	21	21	21	21	21	8
NC2B-H4	40	40	40	38	36	34	33	33	33	33	33	18
Total	209	207	203	194	184	179	169	166	162	161	151	102

Table C-5. Number of ADM for high rise house TSP emissions at varying percentages of data completeness.

% Valid Data	0	10	20	30	40	50	60	70	75	80	90	100
IN2H-H6	27	27	27	23	23	22	19	19	19	19	14	14
IN2H-H7	29	29	28	25	24	24	22	21	21	21	15	15
CA2B-H5	48	48	46	46	44	38	36	36	36	36	35	29
CA2B-H6	43	42	40	40	37	32	32	32	32	30	29	20
NC2B-H3	51	51	51	49	45	42	41	41	41	41	40	33
NC2B-H4	90	90	90	85	79	72	70	70	70	70	70	31
Total	288	287	282	268	252	230	220	219	219	217	203	142

Data Completeness Review and Conclusions for the Manure belt house Dataset

The number of ADM for NH₃ emissions at varying percentages of data completeness for the revised data set are shown in Table B-6. For the layer site data set, decreasing the daily completeness criteria from 75% to 50% would increase the number of valid days by 95 (7.6%), but based on the Grant et al. (2013) study there would be an approximate 15% increase in error. Since the small increase in the number of ADM values does not justify the 15% increase in error, a daily completeness criterion of 75% was chosen for the revised NH₃ belted battery data set.

Table C-6. Number of manure belt house ADM for NH₃ at varying percentages of data completeness.

% Valid												
Data	0	10	20	30	40	50	60	70	75	80	90	100
IN2B H8	691	690	685	680	676	667	654	639	621	612	603	454
IN2B H9	695	694	690	687	684	678	663	646	629	618	609	462
Total	1,386	1,384	1,375	1,367	1,360	1,345	1,317	1,285	1,250	1,230	1,212	916

For H₂S, the number of ADM at varying percentages of data completeness for the revised data set are shown in Table B-7. For the belted battery data set, decreasing the daily completeness criteria from 75% to 50% would increase the number of valid days by 90 (7.1%), but based on the Grant et al. (2013) study there would be an approximate 15% increase in error. Since the small increase in the number of ADM values does not justify the 15% increase in error, a daily completeness criterion of 75% was chosen for the revised H₂S belted battery data set.

Table C-7. Number of manure belt house ADM for H₂S at varying percentages of data completeness.

% Valid												
Data	0	10	20	30	40	50	60	70	75	80	90	100
IN2B H8	701	701	697	692	688	677	662	649	631	622	614	460
IN2B H9	705	705	702	700	697	689	675	658	645	633	623	473
Total	1,406	1,406	1,399	1,392	1,385	1,366	1,337	1,307	1,276	1,255	1,237	933

For PM₁₀, the number of ADM at varying percentages of data completeness for the revised data set are shown in Table B-8. For the manure belt house data set, decreasing the daily completeness criteria from 75% to 50% would increase the number of valid days by 69 (9.8 %). The number of ADM for PM_{2.5} are presented in Table B-9 and show the number of valid ADM would increase by 6 (10.7%). TSP (Table B-10) had an increase of 15 days (21.7%) when shifting to 50% completeness criteria. Again, the small increase in the number of ADM values does not justify the 15% increase in error. Therefore, a daily completeness criterion of 75% was chosen for all the PM species for the belted battery data set.

Table C-8. Number of manure belt house ADM for PM₁₀ at varying percentages of data completeness.

% Valid Data	0	10	20	30	40	50	60	70	75	80	90	100
IN2B H8	433	429	422	409	393	381	367	354	346	339	300	160
IN2B H9	441	438	430	420	403	395	384	372	361	351	309	195
Total	874	867	852	829	796	776	751	726	707	690	609	355

Table C-9. Number of manure belt house ADM for PM_{2.5} at varying percentages of data completeness.

% Valid Data	0	10	20	30	40	50	60	70	75	80	90	100
IN2B H8	30	29	29	29	29	28	25	25	25	25	24	16
IN2B H9	37	37	37	37	36	34	31	31	31	31	30	20
Total	67	66	66	66	65	62	56	56	56	56	54	36

Table C-10. Number of manure belt house ADM for TSP at varying percentages of data completeness.

% Valid Data	0	10	20	30	40	50	60	70	75	80	90	100
IN2B H8	56	54	52	47	45	41	40	36	35	35	34	20
IN2B H9	56	54	51	47	45	43	41	36	34	34	33	22
Total	112	108	103	94	90	84	81	72	69	69	67	42

Data Completeness Review and Conclusions for Manure Shed Dataset

For the manure shed dataset (Table B-11), reducing the completeness criteria to 50% results in an additional 26 days (5%) for NH₃, 23 days (4.3%) for H₂S, 29 days (9.4%) for PM₁₀, 3 days (10%) for PM_{2.5}, and 4 days (16.7%) for TSP. These modest gains in the number of ADM available do not justify the estimated 15% increase in error.

Table C-11. Number of manure shed ADM for each pollutant at varying percentages of data completeness.

% Valid Data	0	10	20	30	40	50	60	70	75	80	90	100
NH ₃	588	561	554	552	545	544	531	526	518	506	502	394
H ₂ S	586	570	567	566	560	557	547	543	534	520	515	406
PM ₁₀	368	367	365	355	342	336	321	312	307	295	267	183
PM _{2.5}	37	37	37	36	35	33	31	31	30	30	29	21
TSP	38	38	35	33	29	28	27	25	24	24	24	18

Appendix D - Summary Statistics

List of Tables

Table D-1. Summary statistics for NH ₃ emissions (kg d ⁻¹) from high rise layer sites.	D-3
Table D-2. Summary statistics for NH ₃ emissions (kg hd ⁻¹ d ⁻¹) from high rise layer sites	D-3
Table D-3. Summary statistics for H ₂ S emissions (g hd ⁻¹ d ⁻¹) from high rise layer sites	D-3
Table D-4. Summary statistics for H ₂ S emissions (g hd ⁻¹ d ⁻¹) from high rise layer sites	D-3
Table D-5. Summary statistics for PM ₁₀ emissions (g d ⁻¹) from high rise layer sites	D-4
Table D-6. Summary statistics for PM ₁₀ emissions (g hd ⁻¹ d ⁻¹) from high rise layer sites	
Table D-7. Summary statistics for PM _{2.5} emissions (g d ⁻¹) from high rise layer sites	
Table D-8. Summary statistics for PM _{2.5} emissions (g hd ⁻¹ d ⁻¹) from high rise layer sites	
Table D-9. Summary statistics for TSP emissions (g d ⁻¹) from high rise layer sites	
Table D-10. Summary statistics for TSP emissions (g hd ⁻¹ d ⁻¹) from high rise layer sites.	D-5
Table D-11. Summary statistics for NH ₃ emissions (kg d ⁻¹) from manure belt houses and manure shed	D-6
Table D-12. Summary statistics for NH ₃ emissions (kg hd ⁻¹ d ⁻¹) from manure belt houses and manure shed	D-6
Table D-13. Summary statistics for H ₂ S emissions (g d ⁻¹) from manure belt houses and manure shed	D-6
Table D-14. Summary statistics for H ₂ S emissions (g hd ⁻¹ d ⁻¹) from manure belt houses and manure shed	D-7
Table D-15. Summary statistics for PM ₁₀ emissions (g d ⁻¹) from manure belt houses and manure shed	
Table D-16. Summary statistics for PM ₁₀ emissions (g hd ⁻¹ d ⁻¹) from manure belt houses and manure shed	D-7
Table D-17. Summary statistics for PM _{2.5} emissions (g d ⁻¹) from manure belt houses and manure shed	
Table D-18. Summary statistics for PM _{2.5} emissions (g hd ⁻¹ d ⁻¹) from manure belt houses and manure shed	
Table D-19. Summary statistics for TSP emissions (g d-1) from manure belt houses and manure shed	
Table D-20. Summary statistics for TSP emissions (g hd ⁻¹ d ⁻¹) from manure belt houses and manure shed	
Table D-21. Summary statistics of environmental and production parameters at high rise layer sites	
Table D-22. Summary statistics of environmental and production parameters at manure belt houses and manual	re shed
Table D-23. Summary statistics of ambient meteorological parameters at high rise layer sites	
Table D-24. Summary statistics of manure parameters at high rise layer sites	
Table D-25. Summary statistics of manure parameters at manure belt houses and manure shed	D-14

Table D-1. Summary statistics for NH₃ emissions (kg d⁻¹) from high rise layer sites.

Statistic	CA2BH5	CA2BH6	IN2HH6	IN2HH7	NC2BH3	NC2BH4
Mean	33.11	32.27	224.75	248.18	58.09	57.46
St. Dev	17.4	29.8	89.58	102.05	20.85	24.36
N	570	590	564	570	715	715
Median	34.11	20.26	222.17	246.18	56.54	55.19
Min	2.13	1.96	-89.38	-130.32	8.65	1.18
Max	82.96	134.52	614.24	931.89	137.25	171.3
CV(%)	52.558	92.348	39.86	41.121	35.888	42.388
N<0	0	0	2	2	0	0

Table D-2. Summary statistics for NH₃ emissions (kg hd⁻¹ d⁻¹) from high rise layer sites.

Statistic	CA2BH5	CA2BH6	IN2HH6	IN2HH7	NC2BH3	NC2BH4
Mean	0.9626	0.9623	1.0384	1.2152	0.6134	0.6449
St. Dev	0.4940	0.8716	0.4185	1.4782	0.2168	0.2371
Median	0.9907	0.6175	1.0100	1.1361	0.5969	0.6046
Min	0.0657	0.0611	-0.3867	-0.5617	0.1196	0.1524
Max	2.3429	3.9534	3.2128	30.7551	1.4497	1.8541

Table D-3. Summary statistics for H₂S emissions (g hd⁻¹ d⁻¹) from high rise layer sites.

Statistic	CA2BH5	CA2BH6	IN2HH6	IN2HH7	NC2BH3	NC2BH4
Mean	45.41	39.96	301.54	269.27	57.36	62.31
St. Dev	24.04	29.76	277.81	281.05	35.23	43.69
N	601	620	401	410	701	701
Median	44.18	31.44	261.84	199.26	47.74	49.54
Min	-17.3	-23.31	-266.56	-458.66	-24.55	-49.83
Max	125.13	144.55	1,920.23	1,839.53	230.48	311.42
CV(%)	52.939	74.485	92.13	104.377	61.419	70.11
N<0	12	1	20	24	10	22

Table D-4. Summary statistics for H₂S emissions (g hd⁻¹ d⁻¹) from high rise layer sites.

Statistic	CA2BH5	СА2ВН6	IN2HH6	IN2HH7	NC2BH3	NC2BH4
Mean	0.0013	0.0012	0.0014	0.0012	0.0006	0.0007
St. Dev	0.0007	0.0009	0.0012	0.0013	0.0004	0.0004
Median	0.0013	0.0009	0.0012	0.0009	0.0005	0.0005
Min	-0.0005	-0.0007	-0.0012	-0.0020	-0.0001	-0.0001
Max	0.0038	0.0050	0.0088	0.0118	0.0024	0.0032

Table D-5. Summary statistics for PM₁₀ emissions (g d⁻¹) from high rise layer sites.

Statistic	CA2BH5	CA2BH6	IN2HH6	IN2HH7	NC2BH3	NC2BH4
Mean	1,272.88	959.86	3,677.6	4,897.59	1,523.08	2,200.01
St. Dev	1,021.92	795.37	3,230.21	4,004.38	635.89	1,129.83
N	451	527	417	421	383	565
Median	934.54	712.65	3,192.71	3,410.38	1,480.56	2,016.24
Min	-14.69	-570.18	-7,803.59	-1,285.68	-196.7	-296.13
Max	8,138.77	4,689.59	1,4674	24,934.32	5,243.27	9,382.02
CV(%)	80.284	82.864	87.835	81.762	41.751	51.356
N<0	1	8	30	8	3	4

Table D-6. Summary statistics for PM₁₀ emissions (g hd⁻¹ d⁻¹) from high rise layer sites.

Statistic	CA2BH5	СА2ВН6	IN2HH6	IN2HH7	NC2BH3	NC2BH4
Mean	0.0376	0.0296	0.0170	0.0268	0.0162	0.0240
St. Dev	0.0303	0.0267	0.0147	0.0936	0.0061	0.0116
Median	0.0275	0.0211	0.0146	0.0150	0.0156	0.0218
Min	-0.0004	-0.0173	-0.0337	-0.0205	0.0028	0.0016
Max	0.2308	0.2761	0.0662	1.8953	0.0537	0.0983

Table D-7. Summary statistics for PM_{2.5} emissions (g d⁻¹) from high rise layer sites.

Statistic	CA2BH5	CA2BH6	IN2HH6	IN2HH7	NC2BH3	NC2BH4
Mean	237.47	167.71	206.87	101.77	31.14	144.89
St. Dev	530.69	337.07	253.89	22.93	79.04	168.78
N	40	43	16	9	21	48
Median	-8.73	-4.31	128.78	104.42	48.88	81.14
Min	-38.47	-131.01	45.5	56.48	-248.73	-213.08
Max	1,890.27	1,146.13	1,112.38	128.71	97.98	470.88
CV(%)	223.474	200.986	122.732	22.53	253.824	116.483
N<0	29	25	0	0	3	7

Table D-8. Summary statistics for PM_{2.5} emissions (g hd⁻¹ d⁻¹) from high rise layer sites.

Statistic	CA2BH5	СА2ВН6	IN2HH6	IN2HH7	NC2BH3	NC2BH4
Mean	0.0067	0.0052	0.0010	0.0004	0.0003	0.0018
St. Dev	0.0149	0.0103	0.0013	0.0001	0.0008	0.0017
Median	-0.0003	-0.0001	0.0006	0.0005	0.0005	0.0009
Min	-0.0011	-0.0038	0.0002	0.0002	-0.0026	-0.0022
Max	0.0532	0.0352	0.0054	0.0006	0.0010	0.0051

Table D-9. Summary statistics for TSP emissions (g d⁻¹) from high rise layer sites.

Statistic	CA2BH5	СА2ВН6	IN2HH6	IN2HH7	NC2BH3	NC2BH4
Mean	2,436.9	2,765.33	7,803.04	4,652.2	3,433.66	4,147.85
St. Dev	1,382.51	1,455.05	2,804.68	3,739.97	1,514.99	2,428.9
N	36	32	19	21	45	87
Median	2,168.2	2,413.27	7,142.04	3,767.52	3,295.67	4,414.7
Min	177.51	1241.37	4613.03	657.55	983.17	-5.59
Max	5,954.19	7,411.92	15,092.75	13,310.67	7,758.45	16,620.07
CV(%)	56.732	52.618	35.943	80.391	44.122	58.558
N<0	0	0	0	0	0	2

Table D-10. Summary statistics for TSP emissions (g hd⁻¹ d⁻¹) from high rise layer sites.

Statistic	CA2BH5	СА2ВН6	IN2HH6	IN2HH7	NC2BH3	NC2BH4
Mean	0.0719	0.0840	0.0381	0.0219	0.0354	0.0511
St. Dev	0.0410	0.0444	0.0148	0.0177	0.0156	0.0216
Median	0.0625	0.0726	0.0336	0.0178	0.0345	0.0491
Min	0.0054	0.0374	0.0204	0.0029	0.0103	0.0200
Max	0.1772	0.2258	0.0680	0.0620	0.0812	0.1802

Table D-11. Summary statistics for NH₃ emissions (kg d⁻¹) from manure belt houses and manure shed.

Statistic	IN2BH8	IN2HH9	IN2BMS
Mean	71.54	67.05	4.74
St. Dev	37.52	42.99	6.52
N	621	629	518
Median	61.96	57.45	2.82
Min	10.36	1.47	-0.21
Max	294.14	414.61	88.37
CV(%)	52.442	64.126	137.353
N<0	0	0	1

Table D-12. Summary statistics for NH₃ emissions (kg hd⁻¹ d⁻¹) from manure belt houses and manure shed.

Statistic	IN2BH8	IN2BH9	IN2BMS
Mean	0.2863	0.2894	0.0097
St. Dev	0.1477	0.3153	0.0131
Count	621	627	518
Median	0.2473	0.2382	0.0057
Min	0.0428	0.0158	-0.0004
Max	1.1611	6.7292	0.1752
CV(%)	51.61	108.93	135.03
N<0	621	627	1

Table D-13. Summary statistics for H₂S emissions (g d-1) from manure belt houses and manure shed.

Statistic	IN2BH8	IN2HH9	IN2BMS	
Mean	492.07	471.07	35.88	
St. Dev	245.83	268.05	52.37	
N	631	645	534	
Median	440.34	431.86	19.17	
Min	64.85	-59.46	-9.88	
Max	2,163.8	1,915.52	528.18	
CV(%)	49.957	56.903	145.966	
N<0	0	8	5	

Table D-14. Summary statistics for H₂S emissions (g hd⁻¹ d⁻¹) from manure belt houses and manure shed.

Statistic	IN2BH8	IN2BH9	IN2BMS
Mean	0.0020	0.0019	0.0001
St. Dev	0.0009	0.0013	0.0001
Count	631	643	534
Median	0.0018	0.0018	0.00004
Min	0.0003	-0.0118	-0.00002
Max	0.0083	0.0077	0.0010
CV(%)	47.89	66.24	145.05
N<0	631	635	5

Table D-15. Summary statistics for PM₁₀ emissions (g d⁻¹) from manure belt houses and manure shed.

Statistic	IN2BH8	IN2HH9	IN2BMS	
Mean	3,038.68	6,076.17	133.63	
St. Dev	4,812.86	8,237.73	293.18	
N	251	361	307	
Median	1,528.26	3,898.46	58.44	
Min	-3,473.99	-6,845.25	-358.27	
Max	33,224.92	66,503.44	2,557.25	
CV(%)	158.386	135.574	219.404	
N<0	40	48	85	

Table D-16. Summary statistics for PM₁₀ emissions (g hd⁻¹ d⁻¹) from manure belt houses and manure shed.

Statistic	IN2BH8	IN2BH9	IN2BMS
Mean	0.0122	0.0148	0.0003
St. Dev	0.0193	0.1395	0.0006
Count	251	361	307
Median	0.0064	0.0158	0.0001
Min	-0.0144	-2.2450	-0.0009
Max	0.1359	0.2693	0.0050
CV(%)	158.22	943.04	217.91
N<0	211	313	85

Table D-17. Summary statistics for PM_{2.5} emissions (g d⁻¹) from manure belt houses and manure shed.

Statistic	IN2BH8	IN2HH9	IN2BMS
Mean	-85.13	113.21	48.26
St. Dev	690.04	250.55	74.25
N	25	31	30
Median	-235.63	114.89	15.41
Min	-1,329.54	-198.59	3.23
Max	1,387.77	688.04	320.54
CV(%)	-810.558	221.313	153.847
N<0	17	13	0

Table D-18. Summary statistics for PM_{2.5} emissions (g hd⁻¹ d⁻¹) from manure belt houses and manure shed.

Statistic	IN2BH8	IN2BH9	IN2BMS
Mean	-0.0004	0.0005	0.0001
St. Dev	0.0028	0.0010	0.0001
Count	25	31	30
Median	-0.0010	0.0005	0.00003
Min	-0.0055	-0.0008	0.00001
Max	0.0056	0.0028	0.0006
CV(%)	-746.00	220.03	151.76
N<0	8	18	0

Table D-19. Summary statistics for TSP emissions (g d⁻¹) from manure belt houses and manure shed.

Statistic	IN2BH8	IN2HH9	IN2BMS	
Mean	8,136.31	21,870.99	295.23	
St. Dev	5,866.62	22,337.67	413.03	
N	35	34	24	
Median	8,301.25	13,406.91	173.82	
Min	-694.57	11.95	-212.01	
Max	2,8130.03	92,287.15	1,763.37	
CV(%)	72.104	102.134	139.902	
N<0	1	0	3	

Table D-20. Summary statistics for TSP emissions (g hd⁻¹ d⁻¹) from manure belt houses and manure shed.

Statistic	tic IN2BH8 IN2BH9		IN2BMS
Mean	0.0333	0.0888	0.0006
St. Dev	0.0245	0.0913	0.0008
Count	35	34	24
Median	0.0329	0.0540	0.0004
Min	-0.0029	0.0000	-0.0004
Max	0.1180	0.3791	0.0036
CV(%)	73.46	102.81	139.25
N<0	34	34	3

Table D-21. Summary statistics of environmental and production parameters at high rise layer sites.

Davamatav	Chatiatia	CARRIE	CARRILL	INIZILILIC	18121117	NCODIIO	NCODIIA
Parameter	Statistic	CA2BH5	CA2BH6	IN2HH6	IN2HH7	NC2BH3	NC2BH4
	Mean	33,103.66	32,160.67	218,158.23	218,847.77	93,403.79	89,377.73
	St. Dev	6,032.20	5,963.76	21,556.81	20,610.54	14,057.81	19,925.12
Inventory	N	731	731	730	728	822	813
(head)	Median	34,223.00	33,183.00	221,708.00	221,097.00	95,448.75	93,065.51
	Min	0.00	0.00	0.00	0.00	0.00	0.00
	Max	35,675	34,697	232,157	232,382	97,995	97,346
	CV(%)	18.22	18.54	9.88	9.42	15.05	22.29
	Mean	1.66	1.59	1.44	1.46	1.52	1.47
	St. Dev	0.13	0.12	0.06	0.08	0.09	0.05
Hen	N	707	706	723	727	805	776
weight	Median	1.71	1.63	1.46	1.47	1.51	1.49
(kg)	Min	1.26	1.25	1.23	1.21	1.21	1.21
	Max	1.81	1.72	1.60	1.60	1.68	1.59
	CV(%)	7.67	7.47	4.27	5.82	6.21	3.60
	Mean	55,092.43	51,131.61	316,317.70	318,424.45	141,782.72	130,992.28
Live	St. Dev	10,831.62	10,205.98	24,541.93	31,468.14	22,884.01	29,180.99
Animal	N	729	729	724	728	822	813
Weight	Median	57,424.08	54,072.44	317,443.63	323,738.24	142,494.48	137,546.59
(kg)	Min	0.00	0.00	0.00	0.00	0.00	0.00
	Max	64,075.81	59,195.52	347,358.00	355,361.82	160,364.60	142,885.31
	CV(%)	19.66	19.96	7.76	9.88	16.14	22.28
	Mean	239.00	195.50	29.50	356.50	307.00	210.36
	St. Dev	137.84	112.73	16.89	205.68	177.10	143.63
Hen Age	N	477	390	58	712	613	537
(days)	Median	239.00	195.50	29.50	356.50	307.00	202.00
	Min	1.00	1.00	1.00	1.00	1.00	1.00
	Max	477.00	390.00	58.00	712.00	613.00	470.00
	CV(%)	57.68	57.66	57.24	57.70	57.69	68.28
	Mean	57.92	60.81	48.72	51.59	68.40	68.26
House	St. Dev	6.63	8.49	5.51	6.15	7.76	7.90
Relative	N	695	704	700	699	811	811
Humidity	Median	57.90	60.30	48.50	52.20	69.18	69.11
(%)	Min	37.20	36.90	31.80	34.50	43.88	45.73
	Max	75.30	82.70	65.50	66.90	85.47	86.17
	CV(%)	11.45	13.96	11.31	11.92	11.34	11.58
	Mean	22.33	22.30	22.33	22.53	24.63	24.43
	St. Dev	3.28	3.36	3.69	3.67	2.77	3.10
House	N Na - di	719	719	700	700	816	811
Temperature (°C)	Median	22.40	22.30	22.45	22.60	24.69	24.60
(C)	Min	15.00	13.40	9.80	8.50	13.50	11.09
	Max	28.60	29.10	30.00	30.70	31.72	31.37
	CV(%)	14.71	15.06	16.54	16.30	11.25	12.69
	Mean	47.76	46.22	176.37	183.89	109.15	111.90
	St. Dev	20.67	21.19	162.18	174.51	93.25	92.69
Airflow	N	642	672	657	575	812	803
(dsm3/s)	Median	45.99	46.09	92.17	93.48	65.21	73.44
	Min	15.63	12.24	13.42	32.22	1.77	15.28
	Max	86.80	102.83	657.95	675.84	278.30	281.63
	CV(%)	43.27	45.84	91.96	94.89	85.43	82.83

Table D-22. Summary statistics of environmental and production parameters at manure belt houses and manure shed.

Parameter	Statistic	IN2BH8	IN2BH9	IN2BMS
	Mean	250,110.53	241,817.18	491,927.85
	St. Dev	10,256.47	29,674.90	31,413.57
	N	731	731	731
Inventory	Median	253,110.00	246,430.00	498,340.00
(head)	Min	228,650.00	0.00	256,320.00
	Max	262,610.00	256,970.00	519,580.00
	CV(%)	4.10	12.27	6.39
	Mean	354,622.32	348,080.54	701,478.59
Average	St. Dev	16,692.23	21,459.59	36,029.59
Live	N	731	710	712
Animal	Median	359,758.00	347,859.95	704,602.80
Weight	Min	300,096.00	264,707.00	363,974.40
(kg)	Max	382,432.40	377,625.00	755,783.00
	CV(%)	4.71	6.17	5.14
	Mean	376.00	217.00	371.00
	St. Dev	211.17	125.14	125.14
Llon Age	N	731	433	433
Hen Age (days)	Median	376.00	217.00	371.00
(uays)	Min	11.00	1.00	155.00
	Max	741.00	433.00	587.00
	CV(%)	56.16	57.67	33.73
	Mean	1.42	1.41	1.41
	St. Dev	0.05	0.08	0.04
	N	731	710	712
Hen weight (kg)	Median	1.43	1.42	1.43
(Kg)	Min	1.20	1.15	1.25
	Max	1.52	1.53	1.47
	CV(%)	3.62	5.56	2.98
	Mean	26.84	26.47	
	St. Dev	2.04	2.46	
Barn	N	712	712	
Temperature	Median	27.05	26.60	
(°C)	Min	20.00	14.90	
	Max	31.00	31.20	
	CV(%)	7.58	9.29	
	Mean	58.23	64.41	
Dorn	St. Dev	6.02	7.84	
Barn Relative	N	651	712	
Humidity	Median	58.40	64.55	
(%)	Min	40.90	43.50	
V/	Max	78.30	83.60	
	CV(%)	10.34	12.17	
	Mean	129.53	127.11	13.28
	St. Dev	91.54	88.67	11.27
Airflow	N	628	632	566
(dsm3/s)	Median	91.45	91.33	9.67
(331113)3)	Min	9.62	19.70	2.68
	Max	542.30	561.77	127.83
	CV(%)	70.67	69.76	84.85

Table D-23. Summary statistics of ambient meteorological parameters at high rise layer sites.

	a		IN2H/IN2B	
Parameter	Statistic	CA2B	Manure Shed	NC2B
	Mean	58.82	68.21	70.61
A la : a t	St. Dev	12.73	13.32	14.17
Ambient Relative	N	661	668	812
Humidity	Median	57.60	67.80	70.75
(%)	Min	31.40	34.60	27.92
(70)	Max	91.40	97.80	100.00
	CV(%)	21.64	19.53	20.07
	Mean	17.49	12.05	16.51
	St. Dev	6.86	11.55	8.34
Ambient	N	731	731	811
Temperature	Median	17.10	13.50	17.23
(°C)	Min	3.70	-24.30	-5.63
	Max	32.10	29.80	31.81
	CV(%)	39.25	95.85	50.51
	Mean		0.73	
	St. Dev		0.77	
Mind Connad	N		693	
Wind Speed	Median		0.49	
(ms ⁻¹)	Min		0.04	
	Max		8.54	
	CV(%)		105.56	

Table D-24. Summary statistics of manure parameters at high rise layer sites.

Parameter	Statistic	CA2BH5	СА2ВН6	IN2HH6	IN2HH7	NC2BH3	NC2BH4
	Mean	71.45	81.31	172.73	342.50	139.28	140.33
	St. Dev	49.06	50.86	100.90	197.60	104.39	105.32
Manure	N	731	731	684	684	660	664
Age	Median	62.00	78.00	171.50	342.50	110.50	111.00
(days)	Min	1.00	1.00	1.00	1.00	1.00	1.00
	Max	192.00	192.00	371.00	684.00	377.00	380.00
	CV(%)	68.66	62.54	58.42	57.69	74.95	75.05
	Mean	7.82	7.60	8.33	8.50	8.19	8.03
	St. Dev	0.36	0.76	0.26	0.20	0.44	0.41
	N	5	8	5	5	6	5
рН	Median	7.82	7.46	8.20	8.50	8.30	8.17
	Min	7.40	6.40	8.08	8.24	7.51	7.57
	Max	8.30	8.72	8.70	8.79	8.77	8.55
	CV(%)	4.65	10.03	3.14	2.36	5.40	5.14
	Mean	66.62	67.71	52.66	52.15	43.08	53.66
	St. Dev	10.89	17.38	8.07	9.89	5.25	20.42
Solids	N	5	8	5	5	6	5
(%)	Median	62.80	70.05	53.43	57.47	41.75	43.10
(70)	Min	55.20	35.60	41.87	39.70	36.20	40.90
	Max	79.30	88.90	63.20	61.73	49.60	89.20
	CV(%)	16.35	25.66	15.32	18.97	12.19	38.05
	Mean	0.37	0.34	0.55	0.63	0.49	0.40
TAN	St. Dev	0.08	0.13	0.13	0.13	0.20	0.22
TAN (% wet	N	5	8	5	5	6	5
(% wet weight	Median	0.41	0.30	0.56	0.69	0.47	0.43
basis)	Min	0.27	0.15	0.38	0.48	0.29	0.10
503137	Max	0.45	0.52	0.73	0.75	0.85	0.68
	CV(%)	20.86	37.88	23.95	19.72	41.19	55.31

Table D-25. Summary statistics of manure parameters at manure belt houses and manure shed.

Parameter	Statistic	IN2BH8	IN2BH9	IN2BMS
Manure Age (days)	Mean		-	195.76
	St. Dev	-	1	132.58
	N	-	1	681
	Median	-	1	171.00
	Min			1.00
	Max			471.00
	CV(%)			67.73
	Mean	7.98	7.77	7.93
	St. Dev	0.54	0.54	0.53
	N	7	7	8
рН	Median	8.24	7.62	7.92
	Min	7.16	7.08	7.31
	Max	8.47	8.53	8.76
	CV(%)	6.78	6.90	6.73
	Mean	39.72	33.04	68.00
	St. Dev	17.04	11.20	14.37
Calida	N	7	7	9
Solids (%)	Median	33.70	28.10	68.30
	Min	23.90	25.40	49.90
	Max	65.10	56.90	86.05
	CV(%)	42.90	33.91	21.14
TAN (% wet weight basis)	Mean	0.59	0.68	0.43
	St. Dev	0.19	0.09	0.17
	N	7	7	8
	Median	0.53	0.71	0.42
	Min	0.38	0.49	0.26
	Max	0.86	0.75	0.80
	CV(%)	32.71	13.79	39.97
TKN (% wet weight basis)	Mean			3.48
	St. Dev	1	-	0.77
	N			7
	Median			3.53
	Min			2.38
	Max			4.69
	CV(%)		-	22.19

Appendix E - **Time Series Plots**

List of Figures

1
-4 -
-5
-6
-7
-8
-9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
E-
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
S.
52
53
54
111111111222222233333E 333444444555

Figure E-52. Trends in manure pH trends at a layer manure shed during NAEMS.	E-55
Figure E-53. Trends in manure solids content trends at a layer manure shed during NAEMS	
Figure E-54. Trends in TAN trends at a layer manure shed during NAEMS.	
Figure E-55. Trends in TKN trends at a layer manure shed during NAEMS	E-58




Figure E-1. NAEMS high rise layer confinement site NH₃ emissions, by site.




Figure E-2. NAEMS high rise layer confinement site H_2S emissions, by site.




Figure E-3. NAEMS high rise layer confinement site PM₁₀ emissions, by site.




Figure E-4. NAEMS high rise layer confinement site $PM_{2.5}$ emissions, by site.




Figure E-5. NAEMS high rise layer confinement site TSP emissions, by site.




Figure E-6. NAEMS manure belt layer confinement site $NH_{\rm 3}$ emissions, by site.




Figure E-7. NAEMS manure belt layer confinement site H₂S emissions, by site.




Figure E-8. NAEMS manure belt layer confinement site PM₁₀ emissions, by site.




Figure E-9. NAEMS manure belt layer confinement site PM_{2.5} emissions, by site.




Figure E-10. NAEMS manure belt layer confinement site TSP emissions, by site.




Figure E-11. NAEMS layer manure shed NH_3 emissions, by site.




Figure E-12. NAEMS layer manure shed H₂S emissions, by site.




Figure E-13. NAEMS layer manure shed PM_{10} emissions, by site.




Figure E-14. NAEMS layer manure shed PM_{2.5} emissions, by site.




Figure E-15. NAEMS layer manure shed TSP emissions, by site.




Figure E-16. Trends in inventory at high rise layer confinement sites, by site, during NAEMS.




Figure E-17. Trends in hen weight at high rise layer confinement sites, by site, during the NAEMS.




Figure E-18. Trends in live animal weight at high rise layer confinement sites, by site, during NAEMS.




Figure E-19. Trends in hen age at high rise layer confinement sites, by site, during NAEMS.




Figure E-20. Trends in house relative humidity at high rise layer confinement sites, by site, during NAEMS.




Figure E-21. Trends in house temperature at high rise layer confinement sites, by site, during NAEMS.




Figure E-22. Trends in airflow at high rise layer confinement sites, by site, during NAEMS.




Figure E-23. Trends in ambient relative humidity at high rise layer confinement sites, by site, during NAEMS.




Figure E-24. Trends in ambient temperature at high rise layer confinement sites, by site, during NAEMS.




Figure E-25. Trends in inventory at manure belt layer confinement sites, by site, during NAEMS.




Figure E-26. Trends in hen weight at manure belt layer confinement sites, by site, during NAEMS.




Figure E-27. Trends in live animal weight at manure belt layer confinement sites, by site, during NAEMS.




Figure E-28. Trends in hen age at manure belt layer confinement sites, by site, during NAEMS.




Figure E-29. Trends in house relative humidity at manure belt layer confinement sites, by site, during NAEMS.




Figure E-30. Trends in house temperature at manure belt layer confinement sites, by site, during NAEMS.




Figure E-31. Trends in house airflow at manure belt layer confinement sites, by site, during NAEMS.




Figure E-32. Trends in ambient relative humidity at manure belt layer confinement sites, by site, during NAEMS.




Figure E-33. Trends in ambient temperature at manure belt layer confinement sites, by site, during NAEMS.




Figure E-34. Trends in inventory at a layer manure shed during NAEMS.




Figure E-35. Trends in inventory with a 5 day lag at a layer manure shed during NAEMS.




Figure E-36. Trends in hen weight at a layer manure shed during NAEMS.




Figure E-37. Trends in live animal weight at a layer manure shed during NAEMS.




Figure E-38. Trends in live animal weight, with a 5 day lag, at a layer manure shed during NAEMS.




Figure E-39. Trends in hen age at a layer manure shed during NAEMS.




Figure E-40. Trends in house airflow at a layer manure shed during NAEMS.




Figure E-41. Trends in ambient relative humidity at a layer manure shed during NAEMS.




Figure E-42. Trends in ambient temperature at a layer manure shed during NAEMS.




Figure E-43. Trends in wind speed at a layer manure shed during NAEMS.




Figure E-44. Trends in manure age trends at high rise layer confinement sites, by site, during NAEMS.




Figure E-45. Trends in manure pH at high rise layer confinement sites, by site, during NAEMS.




Figure E-46. Trends in manure solids content at high rise layer confinement sites, by site, during NAEMS.




Figure E-47. Trends in TAN at high rise layer confinement sites, by site, during NAEMS.




Figure E-48. Trends in manure pH trends at manure belt layer confinement sites, by site, during NAEMS.




Figure E-49. Trends in manure solids content trends at manure belt layer confinement sites, by site, during NAEMS.




Figure E-50. Trends in TAN trends at manure belt layer confinement sites, by site, during NAEMS.




Figure E-51. Trends in manure age trends at a layer manure shed during NAEMS.




Figure E-52. Trends in manure pH trends at a layer manure shed during NAEMS.




Figure E-53. Trends in manure solids content trends at a layer manure shed during NAEMS.




Figure E-54. Trends in TAN trends at a layer manure shed during NAEMS.




Figure E-55. Trends in TKN trends at a layer manure shed during NAEMS.

Appendix F - Scatter Plots

List of Tables

Table F-1: Relationship classification based on R ² values	
Table F-2: Summary of high rise house R ² values	
Table F-3: Summary of belted battery house R ² values	
Table F-4: Summary of manure sned K* values	F-129
List of Figures	
· ·	
Figure F-1. Scatter plot of high rise NH ₃ emissions versus inventory and scatter plot with regression.	F-8
Figure F-2. Scatter plot of high rise H ₂ S emissions versus inventory and scatter plot with regression.	
Figure F-3. Scatter plot of high rise PM ₁₀ emissions versus inventory and scatter plot with regression.	
Figure F-4. Scatter plot of high rise PM _{2.5} emissions versus inventory and scatter plot with regression	
Figure F-5. Scatter plot of high rise TSP emissions versus inventory and scatter plot with regression	
Figure F-6. Scatter plot of high rise NH ₃ emissions versus hen weight and scatter plot with regression.	
Figure F-7. Scatter plot of high rise H_2S emissions versus hen weight and scatter plot with regression. Figure F-8. Scatter plot of high rise PM_{10} emissions versus hen weight and scatter plot with regression.	
Figure F-9. Scatter plot of high rise PM _{2.5} emissions versus hen weight and scatter plot with regression.	
Figure F-10. Scatter plot of high rise TSP emissions versus hen weight and scatter plot with regression.	
Figure F-11. Scatter plot of high rise NH ₃ emissions versus live animal weight and scatter plot with regression.	F-18
Figure F-12. Scatter plot of high rise H ₂ S emissions versus live animal weight and scatter plot with regression	
Figure F-13. Scatter plot of high rise PM ₁₀ emissions versus live animal weight and scatter plot with regression.	
Figure F-14. Scatter plot of high rise PM _{2.5} emissions versus live animal weight and scatter plot with regression.	
Figure F-15. Scatter plot of high rise TSP emissions versus live animal weight and scatter plot with regression.	
Figure F-16. Scatter plot of high rise NH ₃ emissions versus hen age and scatter plot with regression	
Figure F-18. Scatter plot of high rise PM ₁₀ emissions versus hen age and scatter plot with regression.	
Figure F-19. Scatter plot of high rise PM _{2.5} emissions versus hen age and scatter plot with regression.	
Figure F-20. Scatter plot of high rise TSP emissions versus hen age and scatter plot with regression	F-27
Figure F-21. Scatter plot of high rise NH ₃ emissions versus exhaust temperature and scatter plot with regression	F-28
Figure F-22. Scatter plot of high rise H ₂ S emissions versus exhaust temperature and scatter plot with regression.	
Figure F-23. Scatter plot of high rise PM ₁₀ emissions versus exhaust temperature and scatter plot with regression	
Figure F-24. Scatter plot of high rise PM _{2.5} emissions versus exhaust temperature and scatter plot with regression	F-31
Figure F-26. Scatter plot of high rise NH ₃ emissions versus house relative humidity and scatter plot with regression	г-32 F_33
Figure F-27. Scatter plot of high rise H ₂ S emissions versus house relative humidity and scatter plot with regression.	
Figure F-28. Scatter plot of high rise PM_{10} emissions versus house relative humidity and scatter plot with regression	
Figure F-29. Scatter plot of high rise PM _{2.5} emissions versus house relative humidity and scatter plot with regression	
Figure F-30. Scatter plot of high rise TSP emissions versus house relative humidity and scatter plot with regression	
Figure F-31. Scatter plot of high rise NH ₃ emissions versus airflow and scatter plot with regression.	
Figure F-32. Scatter plot of high rise H ₂ S emissions versus airflow and scatter plot with regression.	
Figure F-33. Scatter plot of high rise PM ₁₀ emissions versus airflow and scatter plot with regression. Figure F-34. Scatter plot of high rise PM _{2.5} emissions versus airflow and scatter plot with regression.	
Figure F-35. Scatter plot of high rise TSP emissions versus airflow and scatter plot with regression.	
Figure F-36. Scatter plot of high rise NH ₃ emissions versus ambient temperature and scatter plot with regression.	
Figure F-37. Scatter plot of high rise H ₂ S emissions versus ambient temperature and scatter plot with regression	
Figure F-38. Scatter plot of high rise PM ₁₀ emissions versus ambient temperature and scatter plot with regression	
Figure F-39. Scatter plot of high rise PM _{2.5} emissions versus ambient temperature and scatter plot with regression	
Figure F-40. Scatter plot of high rise TSP emissions versus ambient temperature and scatter plot with regression.	
Figure F-41. Scatter plot of high rise NH ₃ emissions versus ambient relative humidity and scatter plot with regression	
Figure F-42. Scatter plot of high rise H ₂ S emissions versus ambient relative humidity and scatter plot with regression	
Figure F-44. Scatter plot of high rise PM_{25} emissions versus ambient relative humidity and scatter plot with regression	
Figure F-45. Scatter plot of high rise TSP emissions versus ambient relative humidity and scatter plot with regression.	
Figure F-46. Scatter plot of high rise NH ₃ emissions versus manure age and scatter plot with regression.	
Figure F-47. Scatter plot of high rise H ₂ S emissions versus manure age and scatter plot with regression	
Figure F-48. Scatter plot of high rise PM ₁₀ emissions versus manure age and scatter plot with regression	F-55
Figure F-49. Scatter plot of high rise PM _{2.5} emissions versus manure age and scatter plot with regression.	
Figure F-50. Scatter plot of high rise TSP emissions versus manure age and scatter plot with regression.	
Figure F-51. Scatter plot of high rise NH ₃ emissions versus manure pH and scatter plot with regression	
Figure F-52. Scatter plot of high rise H ₂ S emissions versus manure pH and scatter plot with regression	
Figure F-54. Scatter plot of high rise TSP emissions versus manure pH and scatter plot with regression.	
Figure F-55. Scatter plot of high rise NH ₃ emissions versus manure solids content and scatter plot with regression.	
Figure F-56. Scatter plot of high rise H ₂ S emissions versus manure solids content and scatter plot with regression.	
Figure F-57. Scatter plot of high rise PM ₁₀ emissions versus manure solids content and scatter plot with regression	
Figure F-58. Scatter plot of high rise TSP emissions versus manure solids content and scatter plot with regression	F-65

Figure F-59. Scatter plot of high rise NH ₃ emissions versus TAN and scatter plot with regression.	.F-6	66
Figure F-60. Scatter plot of high rise H ₂ S emissions versus TAN and scatter plot with regression.	.F-6	67
Figure F-61. Scatter plot of high rise PM ₁₀ emissions versus TAN and scatter plot with regression.	.F-6	68
Figure F-62. Scatter plot of high rise TSP emissions versus TAN and scatter plot with regression		
Figure F-63. Scatter plot of belted battery NH ₃ emissions versus inventory and scatter plot with regression	.F-7	72
Figure F-64. Scatter plot of belted battery H ₂ S emissions versus inventory and scatter plot with regression.	F-7	73
Figure F-65. Scatter plot of belted battery PM ₁₀ emissions versus inventory and scatter plot with regression.	F-0	74
Figure F-66. Scatter plot of belted battery PM _{2.5} emissions versus inventory and scatter plot with regression.	F-1	, . 75
Figure F-67. Scatter plot of belted battery TSP emissions versus inventory and scatter plot with regression.		
Figure F-68. Scatter plot of belted battery NH ₃ emissions versus hen weight and scatter plot with regression.	E 2	70 77
Figure F-69. Scatter plot of belted battery H ₂ S emissions versus hen weight and scatter plot with regression.		
Figure F-70. Scatter plot of belted battery PM ₁₀ emissions versus hen weight and scatter plot with regression		
Figure F-71. Scatter plot of belted battery PM _{2.5} emissions versus hen weight and scatter plot with regression	.F-8	80
Figure F-72. Scatter plot of belted battery TSP emissions versus hen weight and scatter plot with regression	.F-8	81
Figure F-73. Scatter plot of belted battery NH ₃ emissions versus live animal weight and scatter plot with regression	.F-8	82
Figure F-74. Scatter plot of belted battery H ₂ S emissions versus live animal weight and scatter plot with regression	.F-8	83
Figure F-75. Scatter plot of belted battery PM ₁₀ emissions versus live animal weight and scatter plot with regression	.F-8	84
Figure F-76. Scatter plot of belted battery PM2.5 emissions versus live animal weight and scatter plot with regression		
Figure F-77. Scatter plot of belted battery TSP emissions versus live animal weight and scatter plot with regression	.F-8	86
Figure F-78. Scatter plot of belted battery NH ₃ emissions versus hen age and scatter plot with regression	.F-8	87
Figure F-79. Scatter plot of belted battery H ₂ S emissions versus hen age and scatter plot with regression.	.F-8	88
Figure F-80. Scatter plot of belted battery PM ₁₀ emissions versus hen age and scatter plot with regression.	.F-8	89
Figure F-81. Scatter plot of belted battery PM _{2.5} emissions versus hen age and scatter plot with regression	.F-	90
Figure F-82. Scatter plot of belted battery TSP emissions versus hen age and scatter plot with regression.	F-0	91
Figure F-83. Scatter plot of belted battery NH ₃ emissions versus exhaust temperature and scatter plot with regression.	F-0	92
Figure F-84. Scatter plot of belted battery H ₁ 3 emissions versus exhaust temperature and scatter plot with regression	F_0	92
Figure F-85. Scatter plot of belted battery PM ₁₀ emissions versus exhaust temperature and scatter plot with regression.	F (01
Figure F-86. Scatter plot of belted battery PM _{2.5} emissions versus exhaust temperature and scatter plot with regression.		
Figure F-87. Scatter plot of belted battery TSP emissions versus exhaust temperature and scatter plot with regression.	F (93 04
Figure F-88. Scatter plot of belted battery NH ₃ emissions versus house relative humidity and scatter plot with regression.	.Г-5	90 07
Figure F-89. Scatter plot of belted battery H ₂ S emissions versus house relative humidity and scatter plot with regression.		
Figure F-90. Scatter plot of belted battery PM ₁₀ emissions versus house relative humidity and scatter plot with regression		
Figure F-91. Scatter plot of belted battery PM _{2.5} emissions versus house relative humidity and scatter plot with regression		
Figure F-92. Scatter plot of belted battery TSP emissions versus house relative humidity and scatter plot with regression		
Figure F-93. Scatter plot of belted battery NH ₃ emissions versus airflow and scatter plot with regression		
Figure F-94. Scatter plot of belted battery H ₂ S emissions versus airflow and scatter plot with regressionF	F-1(03
Figure F-95. Scatter plot of belted battery PM ₁₀ emissions versus airflow and scatter plot with regressionF		
Figure F-96. Scatter plot of belted battery PM _{2.5} emissions versus airflow and scatter plot with regressionF		
Figure F-97. Scatter plot of belted battery TSP emissions versus airflow and scatter plot with regressionF	F-10	06
Figure F-98. Scatter plot of belted battery NH3 emissions versus ambient temperature and scatter plot with regressionF	F-10	07
Figure F-99. Scatter plot of belted battery H ₂ S emissions versus ambient temperature and scatter plot with regression		
Figure F-100. Scatter plot of belted battery PM ₁₀ emissions versus ambient temperature and scatter plot with regression	F-10	09
Figure F-101. Scatter plot of belted battery PM _{2.5} emissions versus ambient temperature and scatter plot with regression	F-11	10
Figure F-102. Scatter plot of belted battery TSP emissions versus ambient temperature and scatter plot with regression	F-11	11
Figure F-103. Scatter plot of belted battery NH ₃ emissions versus ambient relative humidity and scatter plot with regression		
Figure F-104. Scatter plot of belted battery H ₂ S emissions versus ambient relative humidity and scatter plot with regression	F-11	13
Figure F-105. Scatter plot of belted battery PM ₁₀ emissions versus ambient relative humidity and scatter plot with regression		
Figure F-106. Scatter plot of belted battery $PM_{2.5}$ emissions versus ambient relative humidity and scatter plot with regression	F_11	15
Figure F-107. Scatter plot of belted battery TSP emissions versus ambient relative humidity and scatter plot with regression	F_11	16
Figure F-108. Scatter plot of belted battery NH ₃ emissions versus manure pH and scatter plot with regression		
Figure F-109. Scatter plot of belted battery H ₂ S emissions versus manure pH and scatter plot with regression.		
Figure F-119. Scatter plot of belted battery PM ₁₀ emissions versus manure pH and scatter plot with regression.		
Figure F-110. Scatter plot of belted battery TSP emissions versus manure pH and scatter plot with regression. Figure F-111. Scatter plot of belted battery TSP emissions versus manure pH and scatter plot with regression. Figure F-110. Scatter plot of belted battery TSP emissions versus manure pH and scatter plot with regression.		
Figure F-112. Scatter plot of belted battery NH ₃ emissions versus manure solids content and scatter plot with regression		
Figure F-113. Scatter plot of belted battery H ₂ S emissions versus manure solids content and scatter plot with regression		
Figure F-114. Scatter plot of belted battery PM ₁₀ emissions versus manure solids content and scatter plot with regression		
Figure F-115. Scatter plot of belted battery TSP emissions versus manure solids content and scatter plot with regression		
Figure F-116. Scatter plot of belted battery NH ₃ emissions versus TAN and scatter plot with regressionF		
Figure F-117. Scatter plot of belted battery H ₂ S emissions versus TAN and scatter plot with regressionF		
Figure F-118. Scatter plot of belted battery PM ₁₀ emissions versus TAN and scatter plot with regression		
Figure F-119. Scatter plot of belted battery TSP emissions versus TAN and scatter plot with regressionF		
Figure F-120. Scatter plot, with regression of manure shed NH ₃ emissions versus inventory (left) and inventory lagged by 5 days (right)F		
Figure F-121. Scatter plot, with regression of manure shed H ₂ S emissions versus inventory (left) and inventory lagged by 5 days (right)F	F-13	32
Figure F-122. Scatter plot, with regression of manure shed PM ₁₀ emissions versus inventory (left) and inventory lagged by 5 days (right)F	F-13	33
Figure F-123. Scatter plot, with regression of manure shed PM _{2.5} emissions versus inventory (left) and inventory lagged by 5 days (right)F	F-13	33
Figure F-124. Scatter plot, with regression of manure shed TSP emissions versus inventory (left) and inventory lagged by 5 days (right)F		
Figure F-125. Scatter plot of manure shed NH ₃ emissions versus hen weight and scatter plot with regression.		
Figure F-126. Scatter plot of manure shed H ₂ S emissions versus hen weight and scatter plot with regression		
Figure F-127. Scatter plot of manure shed PM_{10} emissions versus hen weight and scatter plot with regression		
Figure F-128. Scatter plot of manure shed $PM_{2.5}$ emissions versus hen weight and scatter plot with regression.		
U :		_

Figure F-129. Scatter plot of manure shed TSP emissions versus hen weight and scatter plot with regression	
Figure F-130. Scatter plot, with regression of manure shed NH ₃ emissions versus live animal weight (left) and live animal weight lagged b (right).	
(right). Figure F-131. Scatter plot, with regression of manure shed H ₂ S emissions versus live animal weight (left) and live animal weight lagged by (right).	y 5 days
Figure F-132. Scatter plot, with regression of manure shed PM ₁₀ emissions versus live animal weight (left) and live animal weight lagged by (right).	y 5 days
Figure F-133. Scatter plot, with regression of manure shed PM _{2.5} emissions versus live animal weight (left) and live animal weight lagged b (right).	y 5 days
Figure F-134. Scatter plot, with regression of manure shed TSP emissions versus live animal weight (left) and live animal weight lagged b	y 5 days
(right)	F-139
Figure F-135. Scatter plot of manure shed NH ₃ emissions versus hen age and scatter plot with regression.	
Figure F-136. Scatter plot of manure shed H ₂ S emissions versus hen age and scatter plot with regression.	
Figure F-137. Scatter plot of manure shed PM ₁₀ emissions versus hen age and scatter plot with regression	F-140
Figure F-138. Scatter plot of manure shed PM _{2.5} emissions versus hen age and scatter plot with regression.	F-141
Figure F-139. Scatter plot of manure shed TSP emissions versus hen age and scatter plot with regression.	F-141
Figure F-140. Scatter plot of manure shed NH ₃ emissions versus airflow and scatter plot with regression.	F-142
Figure F-141. Scatter plot of manure shed H ₂ S emissions versus airflow and scatter plot with regression	F-142
Figure F-142. Scatter plot of manure shed PM ₁₀ emissions versus airflow and scatter plot with regression	F-143
Figure F-143. Scatter plot of manure shed PM _{2.5} emissions versus airflow and scatter plot with regression	
Figure F-144. Scatter plot of manure shed TSP emissions versus airflow and scatter plot with regression.	
Figure F-145. Scatter plot of manure shed NH ₃ emissions versus ambient temperature and scatter plot with regression	
Figure F-146. Scatter plot of manure shed H ₂ S emissions versus ambient temperature and scatter plot with regression	F-145
Figure F-147. Scatter plot of manure shed PM ₁₀ emissions versus ambient temperature and scatter plot with regression	F-145
Figure F-148. Scatter plot of manure shed PM _{2.5} emissions versus ambient temperature and scatter plot with regression	
Figure F-149. Scatter plot of manure shed TSP emissions versus ambient temperature and scatter plot with regression.	
Figure F-150. Scatter plot of manure shed NH ₃ emissions versus ambient relative humidity and scatter plot with regression	
Figure F-151. Scatter plot of manure shed H ₂ S emissions versus ambient relative humidity and scatter plot with regression	
Figure F-152. Scatter plot of manure shed PM ₁₀ emissions versus ambient relative humidity and scatter plot with regression	F-148
Figure F-153. Scatter plot of manure shed PM _{2.5} emissions versus ambient relative humidity and scatter plot with regression	F-148
Figure F-154. Scatter plot of manure shed TSP emissions versus ambient relative humidity and scatter plot with regression	
Figure F-155. Scatter plot of manure shed NH ₃ emissions versus wind speed and scatter plot with regression	
Figure F-156. Scatter plot of manure shed H ₂ S emissions versus wind speed and scatter plot with regression	
Figure F-157. Scatter plot of manure shed PM ₁₀ emissions versus wind speed and scatter plot with regression.	F-150
Figure F-158. Scatter plot of manure shed PM _{2.5} emissions versus wind speed and scatter plot with regression	F-151
Figure F-159. Scatter plot of manure shed TSP emissions versus wind speed and scatter plot with regression	F-151
Figure F-160. Scatter plot of manure shed NH ₃ emissions versus manure age and scatter plot with regression.	F-152
Figure F-161. Scatter plot of manure shed H ₂ S emissions versus manure age and scatter plot with regression	F-152
Figure F-162. Scatter plot of manure shed PM ₁₀ emissions versus manure age and scatter plot with regression	
Figure F-163. Scatter plot of manure shed PM _{2.5} emissions versus manure age and scatter plot with regression.	F-153
Figure F-164. Scatter plot of manure shed TSP emissions versus manure age and scatter plot with regression.	F-154
Figure F-165. Scatter plot of manure shed NH ₃ emissions versus manure pH and scatter plot with regression	F-154
Figure F-166. Scatter plot of manure shed H ₂ S emissions versus manure pH and scatter plot with regression.	F-155
Figure F-167. Scatter plot of manure shed PM ₁₀ emissions versus manure pH and scatter plot with regression	F-155
Figure F-168. Scatter plot of manure shed NH ₃ emissions versus manure solids content and scatter plot with regression.	
Figure F-169. Scatter plot of manure shed H ₂ S emissions versus manure solids content and scatter plot with regression	
Figure F-170. Scatter plot of manure shed PM_{10} emissions versus manure solids content and scatter plot with regression	F-157
Figure F-171. Scatter plot of manure shed NH ₃ emissions versus TAN and scatter plot with regression.	
Figure F-172. Scatter plot of manure shed H ₂ S emissions versus TAN and scatter plot with regression.	F-158
Figure F-173. Scatter plot of manure shed PM_{10} emissions versus TAN and scatter plot with regression.	F-158
Figure F-174. Scatter plot of manure shed NH ₃ emissions versus TKN and scatter plot with regression.	
Figure F-175. Scatter plot of manure shed H ₂ S emissions versus TKN and scatter plot with regression.	
Figure F-176. Scatter plot of manure shed PM ₁₀ emissions versus TKN and scatter plot with regression.	
Provide Manager Street 11.10 emissions versus 11.11 and seatter province respectively	100

High Rise – Summary

To further explore the trends between the predictor variables and emissions and determine whether the parameter should be included in developing an EEM, EPA prepared scatter plots of emissions versus the process, environmental, and manure parameters and conducted least squares regression analysis to assess the influence of each variable on emissions. For the regressions, EPA classified the linear relationships based on the ranges in Table 8-1

Table F-1: Relationship classification based on R² values

Range of R ²	Relationship strength
$R^2 = 0$	none
$0 < R^2 \le 0.2$	slight or weak
$0.2 < R^2 \le 0.4$	modest
$0.4 < R^2 \le 0.6$	moderate
$0.6 < R^2 \le 0.8$	moderately strong
$R^2 > 0.8$	strong

Table F-2: Summary of high rise house R² values

Pollutant	Parameter	R ²	Strength	Figure
NH3	Inventory	0.6781	moderately strong	E-1
H2S	Inventory	0.3377	modest	E-2
PM10	Inventory	0.2583	modest	E-3
PM2.5	Inventory	0.0008	Slight or weak	E-4
TSP	Inventory	0.2475	modest	E-5
NH3	Hen weight	0.1012	Slight or weak	E-6
H2S	Hen weight	0.0489	Slight or weak	E-7
PM10	Hen weight	0.1969	Slight or weak	E-8
PM2.5	Hen weight	0.1942	Slight or weak	E-9
TSP	Hen weight	0.2080	modest	E-10
NH3	Live animal weight	0.6869	moderately strong	E-11
H2S	Live animal weight	0.3328	modest	E-12
PM10	Live animal weight	0.2269	modest	E-13
PM2.5	Live animal weight	0.0020	Slight or weak	E-14
TSP	Live animal weight	0.2247	modest	E-15
NH3	Hen age	0.0253	Slight or weak	E-16
H2S	Hen age	0.0008	Slight or weak	E-17
PM10	Hen age	0.0244	Slight or weak	E-18
PM2.5	Hen age	0.0279	Slight or weak	E-19
TSP	Hen age	0.0017	Slight or weak	E-20
NH3	Exhaust temperature	0.0343	Slight or weak	E-21
H2S	Exhaust temperature	0.0367	Slight or weak	E-22

Pollutant	Parameter	R ²	Strength	Figure
PM10	Exhaust temperature	0.0278	Slight or weak	E-23
PM2.5	Exhaust temperature	0.4068	moderate	E-24
TSP	Exhaust temperature	0.0143	Slight or weak	E-25
NH3	House relative humidity	0.1701	Slight or weak	E-26
H2S	House relative humidity	0.0514	Slight or weak	E-27
PM10	House relative humidity	0.0852	Slight or weak	E-28
PM2.5	House relative humidity	0.0410	Slight or weak	E-29
TSP	House relative humidity	0.0650	Slight or weak	E-30
NH3	Airflow	0.0463	Slight or weak	E-31
H2S	Airflow	0.3029	modest	E-32
PM10	Airflow	0.1737	Slight or weak	E-33
PM2.5	Airflow	0.1331	Slight or weak	E-34
TSP	Airflow	0.0057	Slight or weak	E-35
NH3	Ambient temperature	0.1192	Slight or weak	E-36
H2S	Ambient temperature	0.0341	Slight or weak	E-37
PM10	Ambient temperature	0.0165	Slight or weak	E-38
PM2.5	Ambient temperature	0.3637	modest	E-39
TSP	Ambient temperature	0.0959	Slight or weak	E-40
NH3	Ambient relative humidity	0.0684	Slight or weak	E-41
H2S	Ambient relative humidity	0.0088	Slight or weak	E-42
PM10	Ambient relative humidity	0.0018	Slight or weak	E-43
PM2.5	Ambient relative humidity	0.2103	modest	E-44
TSP	Ambient relative humidity	0.0051	Slight or weak	E-45
NH3	Manure age	0.1874	Slight or weak	E-46
H2S	Manure age	0.0037	Slight or weak	E-47
PM10	Manure age	0.1625	Slight or weak	E-48
PM2.5	Manure age	0.1126	Slight or weak	E-49
TSP	Manure age	0.0616	Slight or weak	E-50
NH3	рН	0.2607	modest	E-51
H2S	рН	0.0963	Slight or weak	E-52
PM10	рН	0.1496	Slight or weak	E-53
PM2.5	рН		a	
TSP	рН		a	E-54
NH3	Solids	0.1597	Slight or weak	E-55
H2S	Solids	0.0114	Slight or weak	E-56
PM10	Solids	0.1154	Slight or weak	E-57
PM2.5	Solids		а	
TSP	Solids		a	E-58

Pollutant	Parameter	R ²	Strength	Figure
NH3	TAN	0.4703	moderate	E-59
H2S	TAN	0.0796	Slight or weak	E-60
PM10	TAN	0.1273	Slight or weak	E-61
PM2.5	TAN	a		
TSP	TAN	а		E-62

^a EPA did not have sufficient measurement data from NAEMS to conduct a linear regression analysis.




Figure F-1. Scatter plot of high rise NH₃ emissions versus inventory and scatter plot with regression.




Figure F-2. Scatter plot of high rise H₂S emissions versus inventory and scatter plot with regression.




Figure F-3. Scatter plot of high rise PM₁₀ emissions versus inventory and scatter plot with regression.




Figure F-4. Scatter plot of high rise PM_{2.5} emissions versus inventory and scatter plot with regression.




Figure F-5. Scatter plot of high rise TSP emissions versus inventory and scatter plot with regression.




Figure F-6. Scatter plot of high rise NH₃ emissions versus hen weight and scatter plot with regression.




Figure F-7. Scatter plot of high rise H₂S emissions versus hen weight and scatter plot with regression.




Figure F-8. Scatter plot of high rise PM₁₀ emissions versus hen weight and scatter plot with regression.




Figure F-9. Scatter plot of high rise PM_{2.5} emissions versus hen weight and scatter plot with regression.




Figure F-10. Scatter plot of high rise TSP emissions versus hen weight and scatter plot with regression.




Figure F-11. Scatter plot of high rise NH₃ emissions versus live animal weight and scatter plot with regression.




Figure F-12. Scatter plot of high rise H₂S emissions versus live animal weight and scatter plot with regression.




Figure F-13. Scatter plot of high rise PM₁₀ emissions versus live animal weight and scatter plot with regression.




Figure F-14. Scatter plot of high rise PM_{2.5} emissions versus live animal weight and scatter plot with regression.




Figure F-15. Scatter plot of high rise TSP emissions versus live animal weight and scatter plot with regression.




Figure F-16. Scatter plot of high rise NH₃ emissions versus hen age and scatter plot with regression.




Figure F-17. Scatter plot of high rise H₂S emissions versus hen age and scatter plot with regression.




Figure F-18. Scatter plot of high rise PM₁₀ emissions versus hen age and scatter plot with regression.




Figure F-19. Scatter plot of high rise PM_{2.5} emissions versus hen age and scatter plot with regression.




Figure F-20. Scatter plot of high rise TSP emissions versus hen age and scatter plot with regression.




Figure F-21. Scatter plot of high rise NH₃ emissions versus exhaust temperature and scatter plot with regression.





Figure F-22. Scatter plot of high rise H₂S emissions versus exhaust temperature and scatter plot with regression.



Fit Plot for PM10Emission_g_d




Figure F-24. Scatter plot of high rise PM_{2.5} emissions versus exhaust temperature and scatter plot with regression.




Figure F-25. Scatter plot of high rise TSP emissions versus exhaust temperature and scatter plot with regression.




Figure F-26. Scatter plot of high rise NH₃ emissions versus house relative humidity and scatter plot with regression.




Figure F-27. Scatter plot of high rise H₂S emissions versus house relative humidity and scatter plot with regression.




Figure F-28. Scatter plot of high rise PM₁₀ emissions versus house relative humidity and scatter plot with regression.




Figure F-29. Scatter plot of high rise PM_{2.5} emissions versus house relative humidity and scatter plot with regression.




Figure F-30. Scatter plot of high rise TSP emissions versus house relative humidity and scatter plot with regression.




Figure F-31. Scatter plot of high rise NH₃ emissions versus airflow and scatter plot with regression.




Figure F-32. Scatter plot of high rise H₂S emissions versus airflow and scatter plot with regression.




Figure F-33. Scatter plot of high rise PM₁₀ emissions versus airflow and scatter plot with regression.




Figure F-34. Scatter plot of high rise PM_{2.5} emissions versus airflow and scatter plot with regression.




Figure F-35. Scatter plot of high rise TSP emissions versus airflow and scatter plot with regression.





Figure F-36. Scatter plot of high rise NH₃ emissions versus ambient temperature and scatter plot with regression.



Highrise layers




Figure F-38. Scatter plot of high rise PM₁₀ emissions versus ambient temperature and scatter plot with regression.




Figure F-39. Scatter plot of high rise PM_{2.5} emissions versus ambient temperature and scatter plot with regression.




Figure F-40. Scatter plot of high rise TSP emissions versus ambient temperature and scatter plot with regression.




Figure F-41. Scatter plot of high rise NH₃ emissions versus ambient relative humidity and scatter plot with regression.




Figure F-42. Scatter plot of high rise H₂S emissions versus ambient relative humidity and scatter plot with regression.




Figure F-43. Scatter plot of high rise PM₁₀ emissions versus ambient relative humidity and scatter plot with regression.




Figure F-44. Scatter plot of high rise PM_{2.5} emissions versus ambient relative humidity and scatter plot with regression.




Figure F-45. Scatter plot of high rise TSP emissions versus ambient relative humidity and scatter plot with regression.




Figure F-46. Scatter plot of high rise NH₃ emissions versus manure age and scatter plot with regression.




Figure F-47. Scatter plot of high rise H₂S emissions versus manure age and scatter plot with regression.




Figure F-48. Scatter plot of high rise PM₁₀ emissions versus manure age and scatter plot with regression.




Figure F-49. Scatter plot of high rise PM_{2.5} emissions versus manure age and scatter plot with regression.




Figure F-50. Scatter plot of high rise TSP emissions versus manure age and scatter plot with regression.




Figure F-51. Scatter plot of high rise NH₃ emissions versus manure pH and scatter plot with regression.




Figure F-52. Scatter plot of high rise H₂S emissions versus manure pH and scatter plot with regression.




Figure F-53. Scatter plot of high rise PM₁₀ emissions versus manure pH and scatter plot with regression.




Figure F-54. Scatter plot of high rise TSP emissions versus manure pH and scatter plot with regression.




Figure F-55. Scatter plot of high rise NH₃ emissions versus manure solids content and scatter plot with regression.




Figure F-56. Scatter plot of high rise H₂S emissions versus manure solids content and scatter plot with regression.




Figure F-57. Scatter plot of high rise PM₁₀ emissions versus manure solids content and scatter plot with regression.




Figure F-58. Scatter plot of high rise TSP emissions versus manure solids content and scatter plot with regression.




Figure F-59. Scatter plot of high rise NH₃ emissions versus TAN and scatter plot with regression.




Figure F-60. Scatter plot of high rise H₂S emissions versus TAN and scatter plot with regression.




Figure F-61. Scatter plot of high rise PM₁₀ emissions versus TAN and scatter plot with regression.




Figure F-62. Scatter plot of high rise TSP emissions versus TAN and scatter plot with regression.

Table F-3: Summary of belted battery house R² values

Table F-3: Summary of belted battery nouse R ² values						
Pollutant	Parameter	R ²	Figure	Strength		
H ₂ S	Airflow	0.0450	E-94	Slight or weak		
NH ₃	Airflow	0.0078	E-93	Slight or weak		
PM ₁₀	Airflow	0.0016	E-95	Slight or weak		
PM _{2.5}	Airflow	0.2047	E-96	Modest		
TSP	Airflow	0.1518	E-97	Slight or weak		
H ₂ S	Ambient relative humidity	0.0010	E-104	Slight or weak		
NH ₃	Ambient relative humidity	0.0883	E-103	Slight or weak		
PM ₁₀	Ambient relative humidity	0.0008	E-105	Slight or weak		
PM _{2.5}	Ambient relative humidity	0.0863	E-106	Slight or weak		
TSP	Ambient relative humidity	0.0242	E-107	Slight or weak		
H ₂ S	Ambient temperature	0.0162	E-99	Slight or weak		
NH ₃	Ambient temperature	0.0964	E-98	Slight or weak		
PM ₁₀	Ambient temperature	0.0010	E-100	Slight or weak		
PM _{2.5}	Ambient temperature	0.1353	E-101	Slight or weak		
TSP	Ambient temperature	0.0733	E-102	Slight or weak		
H ₂ S	Exhaust temperature	0.0716	E-84	Slight or weak		
NH ₃	Exhaust temperature	0.00004	E-83	Slight or weak		
PM ₁₀	Exhaust temperature	0.0033	E-85	Slight or weak		
PM _{2.5}	Exhaust temperature	0.1947	E-86	Slight or weak		
TSP	Exhaust temperature	0.0201	E-87	Slight or weak		
H ₂ S	Hen age	0.1609	E-79	Slight or weak		
NH ₃	Hen age	0.0405	E-78	Slight or weak		
PM ₁₀	Hen age	0.1006	E-80	Slight or weak		
PM _{2.5}	Hen age	0.0593	E-81	Slight or weak		
TSP	Hen age	0.0626	E-82	Slight or weak		
H ₂ S	Hen weight	0.0161	E-69	Slight or weak		
NH ₃	Hen weight	0.0461	E-68	Slight or weak		
PM ₁₀	Hen weight	0.0009	E-70	Slight or weak		
PM _{2.5}	Hen weight	0.1552	E-71	Slight or weak		
TSP	Hen weight	0.0087	E-72	Slight or weak		
H ₂ S	House relative humidity	0.0398	E-89	Slight or weak		
NH ₃	House relative humidity	0.1474	E-88	Slight or weak		

Pollutant	Parameter	R ²	Figure	Strength
PM ₁₀	House relative humidity	0.0031	E-90	Slight or weak
PM _{2.5}	House relative humidity	0.0006	E-91	Slight or weak
TSP	House relative humidity	0.0302	E-92	Slight or weak
H ₂ S	Inventory	0.0870	E-64	Slight or weak
NH ₃	Inventory	0.0524	E-63	Slight or weak
PM ₁₀	Inventory	0.0241	E-65	Slight or weak
PM _{2.5}	Inventory	0.3679	E-66	modest
TSP	Inventory	0.0012	E-67	Slight or weak
H ₂ S	Live animal weight	0.0819	E-74	Slight or weak
NH ₃	Live animal weight	0.0833	E-73	Slight or weak
PM ₁₀	Live animal weight	0.0207	E-75	Slight or weak
PM _{2.5}	Live animal weight	0.0357	E-76	Slight or weak
TSP	Live animal weight	0.0096	E-77	Slight or weak
H ₂ S	рН	0.0783	E-109	Slight or weak
NH ₃	рН	0.0231	E-108	Slight or weak
PM ₁₀	рН	0.1230	E-110	Slight or weak
PM _{2.5}	рН	a		
TSP	рН	0.02	E-111	Slight or weak
H ₂ S	Solids	0.0122	E-113	Slight or weak
NH ₃	Solids	0.1110	E-112	Slight or weak
PM ₁₀	Solids	0.2291	E-114	modest
PM _{2.5}	Solids	a		
TSP	Solids	0.057	E-115	Slight or weak
H ₂ S	TAN	0.1739	E-117	Slight or weak
NH ₃	TAN	0.0115	E-116	Slight or weak
PM ₁₀	TAN	0.0119	E-118	Slight or weak
PM _{2.5}	TAN	a		
TSP	TAN	0.219	E-119	modest

^a EPA did not have sufficient measurement data from NAEMS to conduct a linear regression analysis.




Figure F-63. Scatter plot of belted battery NH₃ emissions versus inventory and scatter plot with regression.




Figure F-64. Scatter plot of belted battery H₂S emissions versus inventory and scatter plot with regression.




Figure F-65. Scatter plot of belted battery PM₁₀ emissions versus inventory and scatter plot with regression.




Figure F-66. Scatter plot of belted battery PM_{2.5} emissions versus inventory and scatter plot with regression.




Figure F-67. Scatter plot of belted battery TSP emissions versus inventory and scatter plot with regression.




Figure F-68. Scatter plot of belted battery NH₃ emissions versus hen weight and scatter plot with regression.




Figure F-69. Scatter plot of belted battery H₂S emissions versus hen weight and scatter plot with regression.




Figure F-70. Scatter plot of belted battery PM₁₀ emissions versus hen weight and scatter plot with regression.




Figure F-71. Scatter plot of belted battery PM_{2.5} emissions versus hen weight and scatter plot with regression.




Figure F-72. Scatter plot of belted battery TSP emissions versus hen weight and scatter plot with regression.




Figure F-73. Scatter plot of belted battery NH₃ emissions versus live animal weight and scatter plot with regression.




Figure F-74. Scatter plot of belted battery H₂S emissions versus live animal weight and scatter plot with regression.




Figure F-75. Scatter plot of belted battery PM₁₀ emissions versus live animal weight and scatter plot with regression.




Figure F-76. Scatter plot of belted battery PM_{2.5} emissions versus live animal weight and scatter plot with regression.




Figure F-77. Scatter plot of belted battery TSP emissions versus live animal weight and scatter plot with regression.




Figure F-78. Scatter plot of belted battery NH₃ emissions versus hen age and scatter plot with regression.




Figure F-79. Scatter plot of belted battery H₂S emissions versus hen age and scatter plot with regression.




Figure F-80. Scatter plot of belted battery PM₁₀ emissions versus hen age and scatter plot with regression.




Figure F-81. Scatter plot of belted battery PM_{2.5} emissions versus hen age and scatter plot with regression.




Figure F-82. Scatter plot of belted battery TSP emissions versus hen age and scatter plot with regression.




Figure F-83. Scatter plot of belted battery NH₃ emissions versus exhaust temperature and scatter plot with regression.




Figure F-84. Scatter plot of belted battery H₂S emissions versus exhaust temperature and scatter plot with regression.




Figure F-85. Scatter plot of belted battery PM₁₀ emissions versus exhaust temperature and scatter plot with regression.




Figure F-86. Scatter plot of belted battery PM_{2.5} emissions versus exhaust temperature and scatter plot with regression.




Figure F-87. Scatter plot of belted battery TSP emissions versus exhaust temperature and scatter plot with regression.




Figure F-88. Scatter plot of belted battery NH₃ emissions versus house relative humidity and scatter plot with regression.




Figure F-89. Scatter plot of belted battery H₂S emissions versus house relative humidity and scatter plot with regression.




Figure F-90. Scatter plot of belted battery PM₁₀ emissions versus house relative humidity and scatter plot with regression.




Figure F-91. Scatter plot of belted battery PM_{2.5} emissions versus house relative humidity and scatter plot with regression.




Figure F-92. Scatter plot of belted battery TSP emissions versus house relative humidity and scatter plot with regression.




Figure F-93. Scatter plot of belted battery NH₃ emissions versus airflow and scatter plot with regression.




Figure F-94. Scatter plot of belted battery H₂S emissions versus airflow and scatter plot with regression.




Figure F-95. Scatter plot of belted battery PM₁₀ emissions versus airflow and scatter plot with regression.




Figure F-96. Scatter plot of belted battery PM_{2.5} emissions versus airflow and scatter plot with regression.




Figure F-97. Scatter plot of belted battery TSP emissions versus airflow and scatter plot with regression.




Figure F-98. Scatter plot of belted battery NH₃ emissions versus ambient temperature and scatter plot with regression.




Figure F-99. Scatter plot of belted battery H₂S emissions versus ambient temperature and scatter plot with regression.




Figure F-100. Scatter plot of belted battery PM₁₀ emissions versus ambient temperature and scatter plot with regression.




Figure F-101. Scatter plot of belted battery PM_{2.5} emissions versus ambient temperature and scatter plot with regression.




Figure F-102. Scatter plot of belted battery TSP emissions versus ambient temperature and scatter plot with regression.




Figure F-103. Scatter plot of belted battery NH₃ emissions versus ambient relative humidity and scatter plot with regression.




Figure F-104. Scatter plot of belted battery H₂S emissions versus ambient relative humidity and scatter plot with regression.




Figure F-105. Scatter plot of belted battery PM₁₀ emissions versus ambient relative humidity and scatter plot with regression.




Figure F-106. Scatter plot of belted battery PM_{2.5} emissions versus ambient relative humidity and scatter plot with regression.




Figure F-107. Scatter plot of belted battery TSP emissions versus ambient relative humidity and scatter plot with regression.




Figure F-108. Scatter plot of belted battery NH₃ emissions versus manure pH and scatter plot with regression.




Figure F-109. Scatter plot of belted battery H₂S emissions versus manure pH and scatter plot with regression.




Figure F-110. Scatter plot of belted battery PM₁₀ emissions versus manure pH and scatter plot with regression.




Figure F-111. Scatter plot of belted battery TSP emissions versus manure pH and scatter plot with regression.




Figure F-112. Scatter plot of belted battery NH₃ emissions versus manure solids content and scatter plot with regression.




Figure F-113. Scatter plot of belted battery H₂S emissions versus manure solids content and scatter plot with regression.




Figure F-114. Scatter plot of belted battery PM₁₀ emissions versus manure solids content and scatter plot with regression.




Figure F-115. Scatter plot of belted battery TSP emissions versus manure solids content and scatter plot with regression.




Figure F-116. Scatter plot of belted battery NH₃ emissions versus TAN and scatter plot with regression.




Figure F-117. Scatter plot of belted battery H₂S emissions versus TAN and scatter plot with regression.




Figure F-118. Scatter plot of belted battery PM₁₀ emissions versus TAN and scatter plot with regression.




Figure F-119. Scatter plot of belted battery TSP emissions versus TAN and scatter plot with regression.

Table F-4: Summary of manure shed R² values

Pollutant	Parameter	R ²	Strength	Figure
NH ₃	Inventory	0.0152	Slight or weak	E-120
H ₂ S	Inventory	0.0158	Slight or weak	E-121
PM ₁₀	Inventory	0.3074	modest	E-122
PM _{2.5}	Inventory	0.3074	modest	E-123
TSP	Inventory	0.0216	Slight or weak	E-124
NH ₃	Inventory, 5 day lag	0.0147	Slight or weak	E-120
H ₂ S	Inventory, 5 day lag	0.0120	Slight or weak	E-121
PM ₁₀	Inventory, 5 day lag	0.3074	modest	E-122
PM _{2.5}	Inventory, 5 day lag	0.3074	modest	E-123
TSP	Inventory, 5 day lag	0.0206	Slight or weak	E-124
NH ₃	Hen weight	0.0003	Slight or weak	E-125
H ₂ S	Hen weight	0.0278	Slight or weak	E-126
PM ₁₀	Hen weight	0.0214	Slight or weak	E-127
PM _{2.5}	Hen weight	0.1815	Slight or weak	E-128
TSP	Hen weight	0.0404	Slight or weak	E-129
NH ₃	Live animal weight	0.0026	Slight or weak	E-130
H ₂ S	Live animal weight	0.0014	Slight or weak	E-131
PM ₁₀	Live animal weight	0.1730	Slight or weak	E-132
PM _{2.5}	Live animal weight	0.2852	modest	E-133
TSP	Live animal weight	0.0838	Slight or weak	E-134
NH ₃	Live animal weight, 5 day lag	0.0000	Slight or weak	E-130
H ₂ S	Live animal weight, 5 day lag	0.0102	Slight or weak	E-131
PM ₁₀	Live animal weight, 5 day lag	0.0164	Slight or weak	E-132
PM _{2.5}	Live animal weight, 5 day lag	0.2464	modest	E-133
TSP	Live animal weight, 5 day lag	0.0954	Slight or weak	E-134
NH ₃	Hen age	0.0206	Slight or weak	E-135
H ₂ S	Hen age	0.0530	Slight or weak	E-136
PM ₁₀	Hen age	0.0006	Slight or weak	E-137
PM _{2.5}	Hen age	0.4424	moderate	E-138

Pollutant	Parameter	R ²	Strength	Figure
TSP	Hen age	0.0717	Slight or weak	E-139
NH ₃	Airflow	0.4745	moderate	E-140
H ₂ S	Airflow	0.3917	modest	E-141
PM ₁₀	Airflow	0.1877	Slight or weak	E-142
PM _{2.5}	Airflow	0.6428	moderately strong	E-143
TSP	Airflow	0.0689	Slight or weak	E-144
NH ₃	Ambient temperature	0.0328	Slight or weak	E-145
H ₂ S	Ambient temperature	0.0813	Slight or weak	E-146
PM ₁₀	Ambient temperature	0.0030	Slight or weak	E-147
PM _{2.5}	Ambient temperature	0.0040	Slight or weak	E-148
TSP	Ambient temperature	0.0069	Slight or weak	E-149
NH ₃	Ambient relative humidity	0.0057	Slight or weak	E-150
H ₂ S	Ambient relative humidity	0.0200	Slight or weak	E-151
PM ₁₀	Ambient relative humidity	0.0065	Slight or weak	E-152
PM _{2.5}	Ambient relative humidity	0.1697	Slight or weak	E-153
TSP	Ambient relative humidity	0.0066	Slight or weak	E-154
NH ₃	Wind Speed	0.0017	Slight or weak	E-155
H ₂ S	Wind Speed	0.0022	Slight or weak	E-156
PM ₁₀	Wind Speed	0.0058	Slight or weak	E-157
PM _{2.5}	Wind Speed	0.0086	Slight or weak	E-158
TSP	Wind Speed	0.0001	Slight or weak	E-159
NH ₃	Manure age	0.0277	Slight or weak	E-160
H₂S	Manure age	0.0251	Slight or weak	E-161
PM ₁₀	Manure age	0.0010	Slight or weak	E-162
PM _{2.5}	Manure age	0.0080	Slight or weak	E-163
TSP	Manure age	0.0904	Slight or weak	E-164
NH ₃	рН	0.4836	moderate	E-165
H ₂ S	рН	0.7927	moderately strong	E-166
PM ₁₀	рН	0.0869	Slight or weak	E-167
PM _{2.5}	рН		a	
TSP	рН		a	

Pollutant	Parameter	R ²	Strength	Figure
NH ₃	Solids	0.0066	Slight or weak	E-168
H ₂ S	Solids	0.0406	Slight or weak	E-169
PM ₁₀	Solids	0.1666	Slight or weak	E-170
PM _{2.5}	Solids	a		
TSP	Solids		a	
NH ₃	TAN	0.1008	Slight or weak	E-171
H ₂ S	TAN	0.0067	Slight or weak	E-172
PM ₁₀	TAN	0.9321	strong	E-173
PM _{2.5}	TAN		a	
TSP	TAN		a	
NH ₃	TKN	0.2271	modest	E-174
H₂S	TKN	0.1150	Slight or weak	E-175
PM ₁₀	TKN	0.7744	moderately strong	E-176
PM _{2.5}	TKN		a	
TSP	TKN		a	

^a EPA did not have sufficient measurement data from NAEMS to conduct a linear regression analysis.






Figure F-120. Scatter plot, with regression of manure shed NH₃ emissions versus inventory (left) and inventory lagged by 5 days (right).




Figure F-121. Scatter plot, with regression of manure shed H₂S emissions versus inventory (left) and inventory lagged by 5 days (right).




Figure F-122. Scatter plot, with regression of manure shed PM₁₀ emissions versus inventory (left) and inventory lagged by 5 days (right).




Figure F-123. Scatter plot, with regression of manure shed $PM_{2.5}$ emissions versus inventory (left) and inventory lagged by 5 days (right).




Figure F-124. Scatter plot, with regression of manure shed TSP emissions versus inventory (left) and inventory lagged by 5 days (right).




Figure F-125. Scatter plot of manure shed NH₃ emissions versus hen weight and scatter plot with regression.




Figure F-126. Scatter plot of manure shed H₂S emissions versus hen weight and scatter plot with regression.




Figure F-127. Scatter plot of manure shed PM₁₀ emissions versus hen weight and scatter plot with regression.




Figure F-128. Scatter plot of manure shed PM_{2.5} emissions versus hen weight and scatter plot with regression.




Figure F-129. Scatter plot of manure shed TSP emissions versus hen weight and scatter plot with regression.

Manure Shed - LAW




Figure F-130. Scatter plot, with regression of manure shed NH₃ emissions versus live animal weight (left) and live animal weight lagged by 5 days (right).




Figure F-131. Scatter plot, with regression of manure shed H₂S emissions versus live animal weight (left) and live animal weight lagged by 5 days (right).




Figure F-132. Scatter plot, with regression of manure shed PM₁₀ emissions versus live animal weight (left) and live animal weight lagged by 5 days (right).




Figure F-133. Scatter plot, with regression of manure shed PM_{2.5} emissions versus live animal weight (left) and live animal weight lagged by 5 days (right).




Figure F-134. Scatter plot, with regression of manure shed TSP emissions versus live animal weight (left) and live animal weight lagged by 5 days (right).




Figure F-135. Scatter plot of manure shed NH₃ emissions versus hen age and scatter plot with regression.




Figure F-136. Scatter plot of manure shed H₂S emissions versus hen age and scatter plot with regression.




Figure F-137. Scatter plot of manure shed PM₁₀ emissions versus hen age and scatter plot with regression.




Figure F-138. Scatter plot of manure shed $PM_{2.5}$ emissions versus hen age and scatter plot with regression.




Figure F-139. Scatter plot of manure shed TSP emissions versus hen age and scatter plot with regression.

Manure Shed - Airflow




Figure F-140. Scatter plot of manure shed NH₃ emissions versus airflow and scatter plot with regression.




Figure F-141. Scatter plot of manure shed H₂S emissions versus airflow and scatter plot with regression.




Figure F-142. Scatter plot of manure shed PM₁₀ emissions versus airflow and scatter plot with regression.




Figure F-143. Scatter plot of manure shed PM_{2.5} emissions versus airflow and scatter plot with regression.




Figure F-144. Scatter plot of manure shed TSP emissions versus airflow and scatter plot with regression.




Figure F-145. Scatter plot of manure shed NH_3 emissions versus ambient temperature and scatter plot with regression.




Figure F-146. Scatter plot of manure shed H₂S emissions versus ambient temperature and scatter plot with regression.




Figure F-147. Scatter plot of manure shed PM_{10} emissions versus ambient temperature and scatter plot with regression.




Figure F-148. Scatter plot of manure shed $PM_{2.5}$ emissions versus ambient temperature and scatter plot with regression.




Figure F-149. Scatter plot of manure shed TSP emissions versus ambient temperature and scatter plot with regression.

Manure Shed – Ambient Relative Humidity




Figure F-150. Scatter plot of manure shed NH₃ emissions versus ambient relative humidity and scatter plot with regression.




Figure F-151. Scatter plot of manure shed H_2S emissions versus ambient relative humidity and scatter plot with regression.




Figure F-152. Scatter plot of manure shed PM_{10} emissions versus ambient relative humidity and scatter plot with regression.




Figure F-153. Scatter plot of manure shed $PM_{2.5}$ emissions versus ambient relative humidity and scatter plot with regression.




Figure F-154. Scatter plot of manure shed TSP emissions versus ambient relative humidity and scatter plot with regression.




Figure F-155. Scatter plot of manure shed NH₃ emissions versus wind speed and scatter plot with regression.




Figure F-156. Scatter plot of manure shed H₂S emissions versus wind speed and scatter plot with regression.




Figure F-157. Scatter plot of manure shed PM_{10} emissions versus wind speed and scatter plot with regression.




Figure F-158. Scatter plot of manure shed $PM_{2.5}$ emissions versus wind speed and scatter plot with regression.




Figure F-159. Scatter plot of manure shed TSP emissions versus wind speed and scatter plot with regression.

Manure Shed – Manure Age




Figure F-160. Scatter plot of manure shed NH₃ emissions versus manure age and scatter plot with regression.




Figure F-161. Scatter plot of manure shed H₂S emissions versus manure age and scatter plot with regression.




Figure F-162. Scatter plot of manure shed PM₁₀ emissions versus manure age and scatter plot with regression.




Figure F-163. Scatter plot of manure shed PM_{2.5} emissions versus manure age and scatter plot with regression.




Figure F-164. Scatter plot of manure shed TSP emissions versus manure age and scatter plot with regression.




Figure F-165. Scatter plot of manure shed NH₃ emissions versus manure pH and scatter plot with regression.




Figure F-166. Scatter plot of manure shed H₂S emissions versus manure pH and scatter plot with regression.




Figure F-167. Scatter plot of manure shed PM₁₀ emissions versus manure pH and scatter plot with regression.

Manure Shed - Solids




Figure F-168. Scatter plot of manure shed NH₃ emissions versus manure solids content and scatter plot with regression.




Figure F-169. Scatter plot of manure shed H₂S emissions versus manure solids content and scatter plot with regression.




Figure F-170. Scatter plot of manure shed PM₁₀ emissions versus manure solids content and scatter plot with regression.




Figure F-171. Scatter plot of manure shed NH_3 emissions versus TAN and scatter plot with regression.




Figure F-172. Scatter plot of manure shed H_2S emissions versus TAN and scatter plot with regression.




Figure F-173. Scatter plot of manure shed PM₁₀ emissions versus TAN and scatter plot with regression.

Manure Shed - TKN




Figure F-174. Scatter plot of manure shed NH₃ emissions versus TKN and scatter plot with regression.




Figure F-175. Scatter plot of manure shed H_2S emissions versus TKN and scatter plot with regression.




Figure F-176. Scatter plot of manure shed PM₁₀ emissions versus TKN and scatter plot with regression.

Appendix G - Daily Models

List of Tables

Table G-1. Parameter combinations tested as models for NH ₃ and H ₂ S emissions	G-4
Table G-2. Parameter combinations tested as models for PM ₁₀ , PM _{2.5} , and TSP emissions	
Table G-3. Parameter and estimates for high rise layer NH ₃ emission models tested	
Table G-4. Fit and evaluation statistics for the high rise house NH ₃ models tested	
Table G-5. Parameter and estimates for high rise layer H ₂ S emission models tested	
Table G-6. Fit and evaluation statistics for the high rise house H ₂ S models tested	
Table G-7. Parameter and estimates for high rise layer PM ₁₀ emission models tested	
Table G-8. Fit and evaluation statistics for the high rise house PM ₁₀ models tested	
Table G-9. Parameter and estimates for high rise layer PM _{2.5} emission models tested	
Table G-10. Fit and evaluation statistics for the high rise house PM _{2.5} models tested	
Table G-11. Parameter and estimates for high rise layer TSP emission models tested	
Table G-12. Fit and evaluation statistics for the high rise house TSP models tested	
Table G-13. Parameter combinations tested as models for manure belt house NH ₃ and H ₂ S emis	
25	
Table G-14. Parameter combinations tested as models for manure belt house PM ₁₀ , PM _{2.5}	, and TSP
emissions	
Table G-15. Parameter and estimates for manure belt house NH ₃ emission models tested	G-26
Table G-16. Fit and evaluation statistics for the manure belt house NH ₃ models tested	G-27
Table G-17. Parameter and estimates for manure belt house H ₂ S emission models tested	G-30
Table G-18. Fit and evaluation statistics for the manure belt house H ₂ S models tested	G-31
Table G-19. Parameter and estimates for manure belt house PM ₁₀ emission models tested	G-34
Table G-20. Fit and evaluation statistics for the manure belt house PM ₁₀ models tested	G-35
Table G-21. Parameter and estimates for manure belt house PM _{2.5} emission models tested	G-38
Table G-22. Fit and evaluation statistics for the manure belt house PM _{2.5} models tested	G-39
Table G-23. Parameter and estimates for manure belt house TSP emission models tested	G-42
Table G-24. Fit and evaluation statistics for the manure belt house TSP models tested	G-43
Table G-25. Parameter combinations tested as models for layer manure shed NH ₃ and H ₂ S emis	sionsG-
46	
Table G-26. Parameter combinations tested as models for layer manure shed PM ₁₀ , PM _{2.5}	, and TSP
emissions	G-46
Table G-27. Parameter and estimates for layer manure shed NH ₃ emission models tested	G-47
Table G-28. Fit and evaluation statistics for the layer manure shed NH ₃ models tested	G-49
Table G-29. Parameter and estimates for layer manure shed H ₂ S emission models tested	G-53
Table G-30. Fit and evaluation statistics for the layer manure shed H ₂ S models tested	G-55
Table G-31. Parameter and estimates for layer manure shed PM ₁₀ emission models tested	G-59
Table G-32. Fit and evaluation statistics for the layer manure shed PM ₁₀ models tested	G-60
Table G-33. Parameter and estimates for layer manure shed PM _{2.5} emission models tested	G-63
Table G-34. Fit and evaluation statistics for the layer manure shed PM _{2.5} models tested	
Table G-35. Parameter and estimates for layer manure shed TSP emission models tested	G-67
Table G-36. Fit and evaluation statistics for the layer manure shed TSP models tested	G-68

List of Figures

Figure G-1 High rise layer house NH ₃ one-to-one plots models 1 through 6	G-7
Figure G-2 High rise layer house NH ₃ one-to-one plots models 7 through 10	G-8
Figure G-3 High rise layer house H ₂ S one-to-one plots models 1 through 6	G-11
Figure G-4 High rise layer house H ₂ S one-to-one plots models 7 through 10	G-12
Figure G-5 High rise layer house PM ₁₀ one-to-one plots for models 1 through 8	G-15
Figure G-6 High rise layer house PM ₁₀ one-to-one plots for models 9 through 15	G-16
Figure G-7 High rise layer house PM _{2.5} one-to-one plots models 1 through 8	G-19
Figure F-G-8 High rise layer house PM _{2.5} one-to-one plots models 9 through 15	G-20
Figure G-9 High rise layer house TSP one-to-one plots models 1 through 8	
Figure G-10 High rise layer house TSP one-to-one plots models 9 through 15	G-24
Figure G-11. Manure belt house NH ₃ one-to-one plots models 1 through 6.	
Figure G-12. Manure belt house NH ₃ one-to-one plots models 7 through 12.	
Figure G-13. Manure belt house H ₂ S one-to-one plots models 1 through 6	
Figure G-14. Manure belt house H ₂ S one-to-one plots models 7 through 12	
Figure G-15. Manure belt house PM ₁₀ one-to-one plots models 1 through 8	
Figure G-16. Manure belt house PM ₁₀ one-to-one plots models 9 through 16	
Figure G-17. Manure belt house PM _{2.5} one-to-one plots models 1 through 8.	
Figure G-18. Manure belt house PM _{2.5} one-to-one plots models 9 through 16	
Figure G-19. Manure belt house TSP one-to-one plots models 1 through 8.	
Figure G-20. Manure belt house TSP one-to-one plots models 9 through 16.	
Figure G-21. Layer manure shed NH_3 one-to-one plots models 1 through 9.	
Figure G-22. Layer manure shed NH ₃ one-to-one plots models 10 through 18	
Figure G-23. Layer manure shed NH ₃ one-to-one plots models 19 through 27	
Figure G-24. Layer manure shed H ₂ S one-to-one plots models 1 through 10	
Figure G-25. Layer manure shed H ₂ S one-to-one plots models 11 through 19	
Figure G-26. Layer manure shed H ₂ S one-to-one plots models 20 through 27	
Figure G-27. Layer manure shed PM ₁₀ one-to-one plots models 1 through 7	
Figure G-28. Layer manure shed PM ₁₀ one-to-one plots models 8 through 14	
Figure G-29. Layer manure shed PM _{2.5} one-to-one plots models 1 through 7	
Figure G-30. Layer manure shed PM _{2.5} one-to-one plots models 7 through 14	
Figure G-31. Layer manure shed TSP one-to-one plots models 1 through 7	
Figure G-32. Layer manure shed TSP one-to-one plots models 8 through 14	G-70

Table G-1. Parameter combinations tested as models for NH₃ and H₂S emissions.

Model	Parameter
G-1	Intercept, Live animal weight, Ambient temperature
G-2	Intercept, Live animal weight, Exhaust temperature
G-3	Intercept, Inventory, Ambient temperature
G-4	Intercept, Inventory, Exhaust temperature
G-5	Intercept, Inventory, Hen age, Ambient temperature
G-6	Intercept, Inventory, Ambient temperature, Management phase (manure cleanouts (C),
G-6	flock emptying and replacement (E), full flock (F), molting (M), and transition (T))
G-7	Intercept, Inventory, Manure age, Ambient temperature
G-8	Intercept, Inventory, Ambient temperature, Ambient relative humidity
G-9	Intercept, Inventory, Ambient temperature, Exhaust relative humidity
G-10	Intercept, Inventory, Ambient temperature, Ambient relative humidity, Management phase
G-10	(C,E,F,M,T)

Table G-2. Parameter combinations tested as models for PM₁₀, PM_{2.5}, and TSP emissions.

Model	Parameter
P-1	Intercept, Inventory
P-2	Intercept, Inventory, Ambient relative humidity
P-3	Intercept, Inventory, Exhaust relative humidity
P-4	Intercept, Inventory, Ambient relative humidity, Ambient temperature
P-5	Intercept, Inventory, Ambient temperature, Exhaust relative humidity
P-6	Intercept, Inventory, Ambient relative humidity, Exhaust temperature
P-7	Intercept, Inventory, Exhaust temperature, Exhaust relative humidity
P-8	Intercept, Live animal weight, Ambient temperature, Ambient relative humidity
P-9	Intercept, Live animal weight, Ambient temperature, Exhaust relative humidity
P-10	Intercept, Live animal weight, Ambient relative humidity, Exhaust temperature
P-11	Intercept, Live animal weight, Exhaust temperature, Exhaust relative humidity
P-12	Intercept, Hen age, Inventory, Ambient relative humidity
P-13	Intercept, Inventory, Ambient relative humidity, Management phase (C,E,F,M,T)
P-14	Intercept, Inventory, Ambient temperature, Ambient relative humidity, Management phase
P-14	(C,E,F,M,T)
P-15	Intercept, Inventory, Manure age, Ambient relative humidity

Table G-3. Parameter and estimates for high rise layer NH₃ emission models tested.

Model	Parameter	Estimate	Standard Error	p-value
	Intercept	2.7356	0.1963	<.0001
G-1	Live animal weight	0.0043	0.0008	<.0001
	Ambient temperature	0.0385	0.0010	<.0001
G-2	Intercept	1.0666	0.2248	<.0001
	Live animal weight	0.0025	0.0010	0.0131
	Exhaust temperature	0.1034	0.0023	<.0001
	Intercept	2.7852	0.2047	<.0001
G-3	Inventory	0.0059	0.0012	<.0001
	Ambient temperature	0.0385	0.0010	<.0001
	Intercept	1.1075	0.2308	<.0001
G-4	Inventory	0.0034	0.0015	0.0259
	Exhaust temperature	0.1032	0.0023	<.0001
	Intercept	2.8780	0.1420	<.0001
6.5	Inventory	0.0038	0.0008	<.0001
G-5	Hen age	0.0002	0.0003	0.6459
	Ambient temperature	0.0411	0.0012	<.0001
	Intercept	2.8863	0.2312	<.0001
	Inventory	0.0049	0.0016	0.0017
	Ambient temperature	0.0384	0.0010	<.0001
	С	0.0118	0.0804	0.8836
G-6	E	-0.2943	0.1268	0.0204
	F	-0.0035	0.0759	0.9632
	M	-0.0610	0.0949	0.5204
	Т	0.0000		
	Intercept	2.3118	0.1459	<.0001
	Inventory	0.0097	0.0009	<.0001
G-7	Manure age	0.0006	0.0002	0.0134
	Ambient temperature	0.0408	0.0011	<.0001
	Intercept	2.6598	0.2257	<.0001
	Inventory	0.0059	0.0013	<.0001
G-8	Ambient temperature	0.0387	0.0010	<.0001
	Ambient relative humidity	0.0018	0.0003	<.0001
	Intercept	2.2104	0.2775	<.0001
	Inventory	0.0060	0.0014	<.0001
G-9	Ambient temperature	0.0388	0.0010	<.0001
	Exhaust relative humidity	0.0082	0.0007	<.0001
	Intercept	2.7728	0.2504	<.0001
	Inventory	0.0049	0.0016	0.0029
	Ambient temperature	0.0386	0.0010	<.0001
	Ambient relative humidity	0.0018	0.0003	<.0001
G-10	C	0.0037	0.0795	0.9626
3 _0	E	-0.3120	0.1282	0.0151
	F	-0.0149	0.0755	0.8435
	M	-0.0665	0.0985	0.4996
	T	0.0000	0.0303	0.4550

Table G-4. Fit and evaluation statistics for the high rise house NH₃ models tested.

						LNMEa	NMEb	ME ^b	MB ^b	NMB ^b
Model	2LogL	AIC	AICc	BIC	Corr.	(%)	(%)	(kg day ⁻¹)	(kg day ⁻¹)	(%)
G-1	-1638	-1608	-1608	-1611	0.543	16.834	59.626	62.555	0.351	0.334
G-2	-2058	-2028	-2028	-2031	0.425	18.743	68.8	72.179	2.42	2.307
G-3	-1635	-1605	-1605	-1608	0.52	17.118	61.181	64.182	1.093	1.042
G-4	-2052	-2022	-2022	-2026	0.402	18.946	69.969	73.401	3.189	3.04
G-5	-1219	-1199	-1199	-1201	0.263	18.838	74.74	71.793	6.855	7.136
G-6	-1644	-1606	-1606	-1610	0.453	17.909	64.732	67.907	2.299	2.192
G-7	-1199	-1187	-1187	-1188	0.686	15.104	59.245	62.519	8.739	8.281
G-8	-1627	-1595	-1595	-1598	0.541	16.851	59.12	60.818	0.165	0.161
G-9	-1778	-1746	-1746	-1749	0.51	17.628	63.48	66.595	1.368	1.304
G-10	-1637	-1597	-1596	-1601	0.474	17.68	62.777	64.581	1.382	1.344




Figure G-1 High rise layer house NH₃ one-to-one plots models 1 through 6.




Figure G-2 High rise layer house NH₃ one-to-one plots models 7 through 10.

Table G-5. Parameter and estimates for high rise layer H₂S emission models tested.

Model	Parameter	Estimate	Standard Error	p-value
	Intercept	3.0046	0.0681	<.0001
G-1	Live animal weight	0.0066	0.0004	<.0001
	Ambient temperature	0.0211	0.0016	<.0001
G-2	Intercept	1.8773	0.1120	<.0001
	Live animal weight	0.0067	0.0004	<.0001
	Exhaust temperature	0.0590	0.0036	<.0001
	Intercept	3.0042	0.0663	<.0001
G-3	Inventory	0.0096	0.0005	<.0001
	Ambient temperature	0.0210	0.0016	<.0001
	Intercept	1.8299	0.1083	<.0001
G-4	Inventory	0.0100	0.0005	<.0001
	Exhaust temperature	0.0597	0.0036	<.0001
	Intercept	3.0811	0.0789	<.0001
6.5	Inventory	0.0083	0.0006	<.0001
G-5	Hen age	0.0002	0.0002	0.2544
	Ambient temperature	0.0228	0.0018	<.0001
	Intercept	3.4171	0.1452	<.0001
	Inventory	0.0082	0.0005	<.0001
	Ambient temperature	0.0217	0.0016	<.0001
C C	С	-0.0773	0.1401	0.5815
G-6	Е	-1.0659	0.1816	<.0001
	F	-0.2473	0.1317	0.0606
	M	-0.5080	0.1512	0.0008
	Т	0.0000		•
	Intercept	3.1374	0.0671	<.0001
G-7	Inventory	0.0096	0.0005	<.0001
G-7	Manure age	-0.0010	0.0002	<.0001
	Ambient temperature	0.0272	0.0021	<.0001
	Intercept	2.7231	0.0726	<.0001
G-8	Inventory	0.0098	0.0005	<.0001
U-8	Ambient temperature	0.0210	0.0016	<.0001
	Ambient relative humidity	0.0038	0.0005	<.0001
	Intercept	1.9138	0.0976	<.0001
G-9	Inventory	0.0117	0.0005	<.0001
U-3	Ambient temperature	0.0229	0.0016	<.0001
	Exhaust relative humidity	0.0124	0.0011	<.0001
	Intercept	3.1737	0.1471	<.0001
	Inventory	0.0082	0.0005	<.0001
	Ambient temperature	0.0217	0.0016	<.0001
	Ambient relative humidity	0.0038	0.0005	<.0001
G-10	С	-0.0987	0.1393	0.4785
	E	-1.0994	0.1781	<.0001
	F	-0.2662	0.1308	0.0421
	M	-0.5322	0.1513	0.0004
	Т	0.0000		

Table G-6. Fit and evaluation statistics for the high rise house H₂S models tested.

						LNME ^a	NME ^b	ME ^b	MB ^b	NMB ^b
Model	2LogL	AIC	AICc	BIC	Corr.	(%)	(%)	(g day ⁻¹)	(g day ⁻¹)	(%)
G-1	1600	1630	1630	1626	0.722	9.972	53.393	58.611	-3.776	-3.44
G-2	1519	1549	1550	1546	0.703	10.867	56.378	61.888	1.676	1.527
G-3	1606	1636	1637	1633	0.721	10.017	53.521	58.714	-2.056	-1.874
G-4	1520	1550	1551	1547	0.704	11.014	56.725	62.228	5.399	4.922
G-5	482	514	514	511	0.725	8.471	50.304	50.219	-8.569	-8.584
G-6	1547	1585	1585	1581	0.743	9.499	53.029	58.174	-9.457	-8.621
G-7	2369	2381	2381	2380	0.746	9.436	48.869	54.842	-3.805	-3.391
G-8	1597	1629	1629	1625	0.729	9.962	52.695	58.631	-1.931	-1.735
G-9	1484	1516	1517	1513	0.727	10.666	55.142	60.492	8.507	7.755
G-10	1534	1574	1574	1570	0.752	9.384	52.358	58.257	-10.55	-9.479




Figure G-3 High rise layer house H₂S one-to-one plots models 1 through 6.




Figure G-4 High rise layer house H₂S one-to-one plots models 7 through 10.

Table G-7. Parameter and estimates for high rise layer PM₁₀ emission models tested.

			Standard	
Model	Parameter	Estimate	Error	p-value
P-1	Intercept	6.9224	0.0627	<.0001
P-1	Inventory	0.0073	0.0006	<.0001
	Intercept	7.0833	0.0688	<.0001
P-2	Inventory	0.0072	0.0006	<.0001
	Ambient relative humidity	-0.0023	0.0005	<.0001
	Intercept	7.4170	0.0988	<.0001
P-3	Inventory	0.0073	0.0007	<.0001
	Exhaust relative humidity	-0.0076	0.0011	<.0001
	Intercept	6.8702	0.0675	<.0001
D 4	Inventory	0.0077	0.0005	<.0001
P-4	Ambient relative humidity	-0.0030	0.0005	<.0001
	Ambient temperature	0.0145	0.0015	<.0001
	Intercept	7.1518	0.0990	<.0001
5.5	Inventory	0.0076	0.0006	<.0001
P-5	Ambient temperature	0.0123	0.0015	<.0001
	Exhaust relative humidity	-0.0068	0.0011	<.0001
	Intercept	6.4350	0.1059	<.0001
	Inventory	0.0070	0.0005	<.0001
P-6	Ambient relative humidity	-0.0028	0.0005	<.0001
	Exhaust temperature	0.0300	0.0038	<.0001
	Intercept	6.7862	0.1286	<.0001
5.7	Inventory	0.0070	0.0006	<.0001
P-7	Exhaust temperature	0.0262	0.0037	<.0001
	Exhaust relative humidity	-0.0069	0.0011	<.0001
	Intercept	6.8633	0.0715	<.0001
D 0	Live animal weight	0.0051	0.0004	<.0001
P-8	Ambient temperature	0.0148	0.0016	<.0001
	Ambient relative humidity	-0.0030	0.0005	<.0001
	Intercept	7.1483	0.1021	<.0001
р 0	Live animal weight	0.0049	0.0004	<.0001
P-9	Ambient temperature	0.0126	0.0015	<.0001
	Exhaust relative humidity	-0.0067	0.0011	<.0001
	Intercept	6.4397	0.1090	<.0001
P-10	Live animal weight	0.0046	0.0004	<.0001
P-10	Ambient relative humidity	-0.0028	0.0005	<.0001
	Exhaust temperature	0.0302	0.0038	<.0001
	Intercept	6.7908	0.1316	<.0001
P-11	Live animal weight	0.0045	0.0004	<.0001
L-11	Exhaust temperature	0.0264	0.0037	<.0001
	Exhaust relative humidity	-0.0068	0.0011	<.0001
	Intercept	7.3244	0.1003	<.0001
D 12	Hen age	-0.0004	0.0003	0.1586
P-12	Inventory	0.0063	0.0009	<.0001
	Ambient relative humidity	-0.0024	0.0006	0.0001

			Standard	
Model	Parameter	Estimate	Error	p-value
	Intercept	6.9875	0.1355	<.0001
	Inventory	0.0061	0.0006	<.0001
	Ambient relative humidity	-0.0022	0.0005	<.0001
P-13	С	0.4360	0.1437	0.0024
P-13	E	-0.4227	0.1547	0.0064
	F	0.1960	0.1251	0.1172
	M	0.3307	0.1596	0.0385
	Т	0.0000	•	
	Intercept	6.7482	0.1334	<.0001
	Inventory	0.0067	0.0006	<.0001
	Ambient temperature	0.0143	0.0015	<.0001
	Ambient relative humidity	-0.0029	0.0005	<.0001
P-14	С	0.4552	0.1401	0.0012
	E	-0.3960	0.1491	0.008
	F	0.2117	0.1219	0.0827
	M	0.3381	0.1543	0.0286
	Т	0.0000	•	
	Intercept	7.0446	0.1363	<.0001
P-15	Inventory	0.0128	0.0013	<.0001
	Manure age	-0.0017	0.0002	<.0001
	Ambient relative humidity	-0.0031	0.0006	<.0001

Table G-8. Fit and evaluation statistics for the high rise house PM_{10} models tested.

						LNME ^a	NME ^b	ME ^b	MB ^b	NMB ^b
Model	2LogL	AIC	AICc	BIC	Corr.	(%)	(%)	(g day ⁻¹)	(g day ⁻¹)	(%)
P-1	730	758	758	755	0.591	5.596	57.434	1379.5	178.68	7.44
P-2	718	748	748	745	0.595	5.465	55.407	1343.4	137.45	5.669
P-3	686	716	717	713	0.593	5.666	59.428	1427.3	266.97	11.115
P-4	633	665	666	662	0.644	4.977	50.241	1218.1	111.63	4.604
P-5	623	655	655	652	0.636	5.206	53.831	1292.9	208.25	8.671
P-6	662	694	694	690	0.638	5.018	50.714	1229.6	57.862	2.386
P-7	641	673	673	670	0.633	5.179	53.465	1284.1	145	6.037
P-8	642	674	674	671	0.626	5.087	51.586	1250.7	63.451	2.617
P-9	632	664	664	660	0.62	5.249	53.941	1295.5	125.04	5.206
P-10	671	703	704	700	0.62	5.118	52.064	1262.3	22.303	0.92
P-11	651	683	683	680	0.617	5.238	53.949	1295.7	82.732	3.445
P-12	116	148	148	144	0.611	5.868	55.809	1507.1	-67.5	-2.5
P-13	686	724	724	720	0.608	5.313	52.088	1262.9	12.381	0.511
P-14	598	638	639	634	0.657	4.85	47.508	1151.9	5.412	0.223
P-15	747	779	779	775	0.549	7.281	105.52	2600.6	1600.4	64.938




Figure G-5 High rise layer house PM₁₀ one-to-one plots for models 1 through 8.




Figure G-6 High rise layer house PM₁₀ one-to-one plots for models 9 through 15.

Table G-9. Parameter and estimates for high rise layer PM_{2.5} emission models tested.

	_		Standard	
Model	Parameter	Estimate	Error	p-value
P-1	Intercept	4.1539	0.2067	<.0001
	Inventory	0.0034	0.0009	0.0037
P-2	Intercept	4.7488	0.2275	<.0001
	Inventory	0.0046	0.0004	<.0001
	Ambient relative humidity	-0.0110	0.0040	0.0068
P-3	Intercept	5.1075	0.9129	<.0001
	Inventory	0.0018	0.0017	0.2927
	Exhaust relative humidity	-0.0120	0.0116	0.3062
	Intercept	4.6219	0.4614	<.0001
P-4	Inventory	0.0080	0.0034	0.0295
	Ambient relative humidity	-0.0181	0.0037	<.0001
	Ambient temperature	0.0510	0.0115	<.0001
	Intercept	4.7297	0.9469	<.0001
P-5	Inventory	0.0020	0.0018	0.2521
	Ambient temperature	0.0234	0.0128	0.074
	Exhaust relative humidity	-0.0088	0.0121	0.4655
	Intercept	3.5383	0.6251	<.0001
P-6	Inventory	0.0055	0.0034	0.1253
	Ambient relative humidity	-0.0171	0.0037	<.0001
	Exhaust temperature	0.0861	0.0223	0.0002
	Intercept	4.0751	1.0128	0.0001
P-7	Inventory	0.0017	0.0016	0.2918
P-/	Exhaust temperature	0.0566	0.0243	0.0229
	Exhaust relative humidity	-0.0145	0.0115	0.2102
	Intercept	4.5909	0.4649	<.0001
P-8	Live animal weight	0.0056	0.0023	0.0266
1-0	Ambient temperature	0.0514	0.0115	<.0001
	Ambient relative humidity	-0.0182	0.0037	<.0001
	Intercept	4.6253	0.8931	<.0001
P-9	Live animal weight	0.0016	0.0012	0.1698
1-9	Ambient temperature	0.0238	0.0130	0.0733
	Exhaust relative humidity	-0.0082	0.0114	0.4712
	Intercept	3.5260	0.6291	<.0001
P-10	Live animal weight	0.0038	0.0023	0.1242
P-10	Ambient relative humidity	-0.0171	0.0037	<.0001
	Exhaust temperature	0.0861	0.0223	0.0002
	Intercept	4.0456	0.9670	<.0001
D 11	Live animal weight	0.0013	0.0011	0.2376
P-11	Exhaust temperature	0.0563	0.0244	0.024
	Exhaust relative humidity	-0.0145	0.0109	0.1841
	Intercept	4.6600	1.6333	0.0208
D 10	Hen age	0.0044	0.0069	0.5523
P-12	Inventory	0.0052	0.0123	0.6927
	Ambient relative humidity	-0.0183	0.0056	0.002
P-13	Intercept	4.7737	0.2244	<.0001

			Standard	
Model	Parameter	Estimate	Error	p-value
	Inventory	0.0045	0.0003	<.0001
	Ambient relative humidity	-0.0110	0.0039	0.0065
	С	0.3491	0.4115	0.4015
	E	-0.7150	1.6965	0.6876
	F	0.0000		
	M	•	•	
	Т	•	•	
	Intercept	4.8303	0.4447	<.0001
	Inventory	0.0065	0.0031	0.0526
	Ambient temperature	0.0516	0.0112	<.0001
	Ambient relative humidity	-0.0183	0.0036	<.0001
P-14	С	0.3710	0.3437	0.2837
	E	-1.0497	0.8527	0.244
	F	0.0000		
	M			
	Т	•		
P-15	Intercept	7.3299	1.0741	<.0001
	Inventory	0.0069	0.0052	0.211
N-12	Manure age	-0.0103	0.0064	0.1291
	Ambient relative humidity	-0.0262	0.0056	<.0001

Table G-10. Fit and evaluation statistics for the high rise house PM_{2.5} models tested.

						LNME ^a	NMEb	ME ^b	MBb	NMB ^b
Model	2LogL	AIC	AICc	BIC	Corr.	(%)	(%)	(g day ⁻¹)	kg day ⁻¹)	(%)
P-1	211	239	242	236	0.273	22.312	125.79	233.25	15.997	8.627
P-2	173	203	207	200	0.364	20.116	116.3	244.83	9.944	4.724
P-3	210	240	243	236	0.272	22.548	121.8	225.86	4.823	2.601
P-4	177	193	194	191	0.733	16.51	78.095	164.4	-44.37	-21.08
P-5	205	237	241	234	0.641	21.312	98.335	182.34	-32.15	-17.34
P-6	181	197	198	195	0.753	16.605	84.537	177.96	-43.62	-20.72
P-7	203	235	239	231	0.688	20.204	98.937	183.45	-32.67	-17.62
P-8	177	193	194	191	0.735	16.479	78.393	165.02	-44.7	-21.23
P-9	205	237	241	234	0.65	21.206	98.682	182.98	-31.46	-16.97
P-10	181	197	198	195	0.747	16.686	85.319	179.6	-42.6	-20.23
P-11	203	235	239	231	0.681	20.164	99.649	184.77	-31.55	-17.02
P-12	97	113	115	111	0.284	24.645	161.21	351.42	122.17	56.045
P-13	171	205	210	202	0.381	19.925	114.6	241.25	7.779	3.695
P-14	174	194	196	192	0.747	15.934	72.31	152.22	-49.53	-23.53
P-15	148	164	165	162	0.688	18.715	88.026	223.43	-15.7	-6.186




Figure G-7 High rise layer house PM_{2.5} one-to-one plots models 1 through 8.




Figure F-G-8 High rise layer house PM_{2.5} one-to-one plots models 9 through 15.

Table G-11. Parameter and estimates for high rise layer TSP emission models tested.

			Standard	
Model	Parameter	Estimate	Error	p-value
P-1	Intercept	7.5811	0.1685	<.0001
	Inventory	0.0059	0.0014	0.0021
	Intercept	7.9009	0.2166	<.0001
P-2	Inventory	0.0068	0.0010	<.0001
	Ambient relative humidity	-0.0059	0.0028	0.0378
	Intercept	8.3610	0.3542	<.0001
P-3	Inventory	0.0054	0.0011	0.0004
	Exhaust relative humidity	-0.0123	0.0053	0.0207
	Intercept	7.5995	0.3020	<.0001
P-4	Inventory	0.0079	0.0012	<.0001
P-4	Ambient relative humidity	-0.0058	0.0028	0.0423
	Ambient temperature	0.0137	0.0092	0.1417
	Intercept	8.1788	0.4033	<.0001
. -	Inventory	0.0061	0.0013	0.0001
P-5	Ambient temperature	0.0088	0.0084	0.2948
	Exhaust relative humidity	-0.0125	0.0053	0.0198
	Intercept	7.1395	0.5337	<.0001
	Inventory	0.0073	0.0008	<.0001
P-6	Ambient relative humidity	-0.0056	0.0028	0.0485
	Exhaust temperature	0.0312	0.0205	0.1296
	Intercept	7.8037	0.5826	<.0001
	Inventory	0.0058	0.0013	0.0009
P-7	Exhaust temperature	0.0235	0.0199	0.2409
	Exhaust relative humidity	-0.0125	0.0053	0.0201
	Intercept	7.5804	0.3157	<.0001
	Live animal weight	0.0049	0.0009	
P-8				<.0001
	Ambient temperature	0.0129	0.0094	0.1751
	Ambient relative humidity	-0.0049	0.0029	0.0899
	Intercept	8.1632	0.4205	<.0001
P-9	Live animal weight	0.0041	0.0012	0.0036
	Ambient temperature	0.0080	0.0085	0.3531
	Exhaust relative humidity	-0.0120	0.0054	0.0267
	Intercept	7.0480	0.5505	<.0001
P-10	Live animal weight	0.0044	0.0008	<.0001
	Ambient relative humidity	-0.0050	0.0028	0.0786
	Exhaust temperature	0.0350	0.0208	0.0949
	Intercept	7.7461	0.6150	<.0001
P-11	Live animal weight	0.0039	0.0013	0.0134
	Exhaust temperature	0.0249	0.0208	0.2321
	Exhaust relative humidity	-0.0120	0.0054	0.0276
	Intercept	7.3039	0.2786	<.0001
D 13	Hen age	0.0005	0.0006	0.3878
P-12	Inventory	0.0102	0.0012	<.0001
	Ambient relative humidity	-0.0007	0.0023	0.7637
P-13	Intercept	8.2312	0.2697	<.0001

			Standard	
Model	Parameter	Estimate	Error	p-value
	Inventory	0.0063	0.0009	<.0001
	Ambient relative humidity	-0.0054	0.0028	0.0601
	С	0.4012	0.1700	0.0256
	E	-3.4233	0.6369	<.0001
	F	-0.2566	0.2260	0.2902
	M		•	
	Т	0.0000	•	
	Intercept	7.9620	0.3193	<.0001
	Inventory	0.0073	0.0012	<.0001
	Ambient temperature	0.0141	0.0089	0.117
	Ambient relative humidity	-0.0052	0.0029	0.0698
P-14	С	0.3540	0.1712	0.0477
	E	-3.3807	0.6245	<.0001
	F	-0.2943	0.2154	0.2055
	M		•	
	Т	0.0000	•	
	Intercept	7.9217	0.2282	<.0001
P-15	Inventory	0.0071	0.0013	0.0001
N-12	Manure age	-0.0003	0.0010	0.7503
	Ambient relative humidity	-0.0059	0.0028	0.0389

Table G-12. Fit and evaluation statistics for the high rise house TSP models tested.

						LNME	NME ^b	ME ^b	MB ^b	NMB ^b
Model	2LogL	AIC	AICc	BIC	Corr.	(%)	(%)	(g day ⁻¹)	(g day ⁻¹)	(%)
P-1	325	353	355	350	0.408	7.541	43.046	1695.7	92.032	2.336
P-2	315	345	348	342	0.504	7.35	39.647	1592.4	38.911	0.969
P-3	320	350	352	346	0.43	7.533	42.667	1680.8	124.87	3.17
P-4	313	345	347	341	0.532	7.467	40.869	1641.5	64.602	1.608
P-5	318	350	353	347	0.454	7.591	43.365	1708.3	130.37	3.31
P-6	313	345	348	342	0.587	7.289	39.629	1591.7	20.431	0.509
P-7	318	350	353	347	0.499	7.465	42.682	1681.4	104.42	2.651
P-8	316	348	351	345	0.523	7.633	42.094	1690.7	26.341	0.656
P-9	321	353	355	350	0.45	7.666	44.261	1743.6	112.75	2.862
P-10	316	348	350	344	0.592	7.47	40.796	1638.6	-11.48	-0.286
P-11	321	353	355	349	0.503	7.546	43.495	1713.4	79.389	2.015
P-12	76	108	112	104	0.424	6.189	54.975	2264.4	830.37	20.16
P-13	298	334	337	330	0.746	6.506	48.702	1956.1	1365.6	34.001
P-14	295	333	337	329	0.747	6.636	48.65	1954	1322.8	32.933
P-15	315	347	350	344	0.522	7.252	38.515	1546.9	-1.332	-0.033




Figure G-9 High rise layer house TSP one-to-one plots models 1 through 8.




Figure G-10 High rise layer house TSP one-to-one plots models 9 through 15.

Table G-13. Parameter combinations tested as models for manure belt house NH₃ and H₂S emissions.

Model	Parameter
G-1	Intercept, Live animal weight, Ambient temperature
G-2	Intercept, Live animal weight, Exhaust temperature
G-3	Intercept, Inventory, Ambient temperature
G-4	Intercept, Inventory, Exhaust temperature
G-5	Intercept, Inventory, Hen age, Ambient temperature
G-6	Intercept, Inventory, Ambient temperature, Management phase (flock emptying and
G-0	replacement (E), full flock (F), molting (M), and transition (T))
G-7	Intercept, Inventory, Ambient temperature, Ambient relative humidity
G-8	Intercept, Inventory, Ambient temperature, Exhaust relative humidity
G-9	Intercept, Inventory, Ambient temperature, Ambient relative humidity, Management phase
G-9	(E,F,M,T)
G-10	Intercept, Inventory
G-11	Intercept, Inventory, Exhaust temperature, Ambient relative humidity
G-12	Intercept, Inventory, Exhaust temperature, Exhaust relative humidity

Table G-14. Parameter combinations tested as models for manure belt house PM_{10} , $PM_{2.5}$, and TSP emissions.

Model	Parameter
P-1	Intercept, Inventory
P-2	Intercept, Inventory, Ambient relative humidity
P-3	Intercept, Inventory, Exhaust relative humidity
P-4	Intercept, Inventory, Ambient relative humidity, Ambient temperature
P-5	Intercept, Inventory, Ambient temperature, Exhaust relative humidity
P-6	Intercept, Inventory, Ambient relative humidity, Exhaust temperature
P-7	Intercept, Inventory, Exhaust temperature, Exhaust relative humidity
P-8	Intercept, Live animal weight, Ambient temperature, Ambient relative humidity
P-9	Intercept, Live animal weight, Ambient temperature, Exhaust relative humidity
P-10	Intercept, Live animal weight, Ambient relative humidity, Exhaust temperature
P-11	Intercept, Live animal weight, Exhaust temperature, Exhaust relative humidity
P-12	Intercept, Hen age, Inventory, Ambient relative humidity
P-13	Intercept, Inventory, Ambient relative humidity, Management phase (C,E,F,M,T)
P-14	Intercept, Inventory, Ambient temperature, Ambient relative humidity, Management phase (C,E,F,M,T)
P-15	Intercept, Inventory, Ambient temperature
P-16	Intercept, Inventory, Exhaust temperature

Table G-15. Parameter and estimates for manure belt house NH₃ emission models tested.

			Standard	
Model	Parameter	Estimate	Error	p-value
	Intercept	2.1289	0.4683	<.0001
G-1	Ambient temperature	0.0291	0.0020	<.0001
	Live animal weight	0.0046	0.0013	0.0005
	Intercept	-0.9761	0.4848	0.0448
G-2	Exhaust temperature	0.1698	0.0090	<.0001
	Live animal weight	0.0015	0.0013	0.2374
	Intercept	2.5569	0.3831	<.0001
G-3	Ambient temperature	0.0293	0.0021	<.0001
	Inventory	0.0048	0.0015	0.0015
	Intercept	-0.9143	0.4132	0.0272
G-4	Exhaust temperature	0.1666	0.0088	<.0001
	Inventory	0.0022	0.0014	0.1142
	Intercept	2.1020	3.4798	0.5496
6.5	Ambient temperature	0.0264	0.0022	<.0001
G-5	Inventory	0.0074	0.0135	0.588
	Hen age	-0.0002	0.0006	0.7011
	Intercept	2.5910	0.3934	<.0001
	E	0.3330	0.1926	0.0841
	F	-0.1036	0.2313	0.6543
G-6	M	-0.2996	0.2681	0.2641
	T	0.0000		
	Ambient temperature	0.0294	0.0021	<.0001
	Inventory	0.0051	0.0016	0.0014
	Intercept	2.4392	0.3808	<.0001
6.7	Ambient temperature	0.0294	0.0021	<.0001
G-7	Inventory	0.0047	0.0015	0.0015
	Ambient relative humidity	0.0019	0.0008	0.0211
	Intercept	2.0188	0.3852	<.0001
6.0	Ambient temperature	0.0294	0.0021	<.0001
G-8	Inventory	0.0046	0.0015	0.0014
	Exhaust relative humidity	0.0090	0.0018	<.0001
	Intercept	2.4867	0.3905	<.0001
	E	0.2968	0.1905	0.1194
	F	-0.1157	0.2279	0.612
	M	-0.3139	0.2642	0.2351
G-9	Т	0.0000		
	Ambient temperature	0.0295	0.0021	<.0001
	Inventory	0.0051	0.0016	0.0014
	Ambient relative humidity	0.0018	0.0008	0.0289
G-10	Intercept	2.5726	0.3433	<.0001

			Standard	
Model	Parameter	Estimate	Error	p-value
	Inventory	0.0061	0.0014	<.0001
	Intercept	-0.7763	0.4047	0.0555
C 11	Inventory	0.0021	0.0014	0.1302
G-11	Exhaust temperature	0.1595	0.0085	<.0001
	Ambient relative humidity	0.0011	0.0008	0.16
	Intercept	-1.2123	0.4148	0.0036
C 40	Inventory	0.0022	0.0014	0.113
G-12	Exhaust temperature	0.1610	0.0090	<.0001
	Exhaust relative humidity	0.0071	0.0017	<.0001

Table G-16. Fit and evaluation statistics for the manure belt house NH₃ models tested.

						LNME	NMEb	MEb	MB ^b	NMB ^b
Model	2LogL	AIC	AICc	BIC	Corr.	(%)	(%)	(kg day ⁻¹)	(kg day ⁻¹)	(%)
G-1	199	209	209	202	-0.09	12.16	59.25	41.16	10.72	15.43
G-2	178	188	188	182	0.108	11.43	55.73	39.09	9.516	13.57
G-3	316	326	326	319	-0.05	12.77	60.08	41.22	10.65	15.53
G-4	269	279	279	273	0.277	11.47	54.81	37.97	8.49	12.26
G-5	85	97	97	89	-0.19	12.1	57.55	41.26	9.484	13.23
G-6	310	326	327	316	-0.04	12.89	60.91	41.79	11.24	16.39
G-7	288	300	300	292	-0.03	12.55	58.8	40.2	9.866	14.43
G-8	227	239	239	231	0.034	12.11	56.04	38.17	8.501	12.48
G-9	283	301	301	289	-0.02	12.68	59.68	40.8	10.48	15.33
G-10	564	572	572	567	0.487	9.444	38.93	26.97	-0.622	-0.898
G-11	157	169	169	161	0.311	11.12	52.15	35.65	7.084	10.36
G-12	205	217	217	209	0.36	10.69	49.86	34.31	5.944	8.639




Figure G-11. Manure belt house NH₃ one-to-one plots models 1 through 6.




Figure G-12. Manure belt house NH₃ one-to-one plots models 7 through 12.

Table G-17. Parameter and estimates for manure belt house H₂S emission models tested.

Model	Parameter	Estimate	Standard Error	p-value
	Intercept	3.9631	0.4108	<.0001
G-1	Ambient temperature	0.0212	0.0018	<.0001
	Live animal weight	0.0055	0.0012	<.0001
	Intercept	1.6607	0.4924	0.0009
G-2	Exhaust temperature	0.1239	0.0080	<.0001
	Live animal weight	0.0034	0.0013	0.0117
	Intercept	4.0179	0.3453	<.0001
G-3	Ambient temperature	0.0212	0.0018	<.0001
	Inventory	0.0076	0.0014	<.0001
	Intercept	1.4304	0.4292	0.001
G-4	Exhaust temperature	0.1225	0.0078	<.0001
	Inventory	0.0058	0.0015	0.0002
	Intercept	5.1432	2.5993	0.0607
G-5	Ambient temperature	0.0189	0.0019	<.0001
u-3	Inventory	0.0042	0.0100	0.6779
	Hen age	-0.0006	0.0005	0.2025
	Intercept	4.0262	0.3491	<.0001
	E	0.7578	0.1809	<.0001
	F	-0.1727	0.2152	0.4225
G-6	M	-0.4080	0.2413	0.0913
	Т	0.0000	•	
	Ambient temperature	0.0209	0.0018	<.0001
	Inventory	0.0084	0.0015	<.0001
	Intercept	3.7391	0.3430	<.0001
G-7	Ambient temperature	0.0222	0.0018	<.0001
"	Inventory	0.0073	0.0014	<.0001
	Ambient relative humidity	0.0048	0.0007	<.0001
	Intercept	3.2727	0.3467	<.0001
G-8	Ambient temperature	0.0230	0.0018	<.0001
	Inventory	0.0075	0.0013	<.0001
	Exhaust relative humidity	0.0121	0.0015	<.0001
	Intercept	3.7730	0.3467	<.0001
	E	0.6634	0.1771	0.0002
	F	-0.2113	0.2105	0.316
G-9	M	-0.4586	0.2363	0.0526
	Т	0.0000		
	Ambient temperature	0.0219	0.0018	<.0001
	Inventory	0.0082	0.0014	<.0001
	Ambient relative humidity	0.0047	0.0007	<.0001
G-10	Intercept	4.3868	0.3996	<.0001
0.10	Inventory	0.0071	0.0016	<.0001

Model	Parameter	Estimate	Standard Error	p-value
	Intercept	1.3286	0.3758	0.0004
G-11	Inventory	0.0054	0.0013	<.0001
G-11	Exhaust temperature	0.1196	0.0072	<.0001
	Ambient relative humidity	0.0041	0.0006	<.0001
	Intercept	0.9834	0.3850	0.0109
C 10	Inventory	0.0049	0.0013	0.0001
G-12	Exhaust temperature	0.1240	0.0086	<.0001
	Exhaust relative humidity	0.0096	0.0016	<.0001

Table G-18. Fit and evaluation statistics for the manure belt house H₂S models tested.

						LNME	NMEb	MEb	MB ^b	NMB ^b
Model	2LogL	AIC	AICc	BIC	Corr.	(%)	(%)	(g day ⁻¹)	(g day ⁻¹)	(%)
G-1	-136	-122	-122	-131	0.385	5.998	39.52	194.2	16.27	3.312
G-2	43	57	57	48	0.37	6.085	39.88	194.9	19.28	3.944
G-3	-74	-60	-59	-69	0.482	6.227	39.09	189.5	11.6	2.394
G-4	76	90	90	81	0.522	6.051	38.77	187	13.41	2.781
G-5	-215	-199	-199	-210	0.411	5.861	36.81	186.3	6.171	1.219
G-6	-97	-77	-77	-90	0.516	6.087	38.3	185.6	10.62	2.191
G-7	-122	-106	-106	-116	0.505	6.180	38.39	186	7.55	1.559
G-8	-165	-149	-149	-159	0.566	5.938	36.77	176.4	3.694	0.77
G-9	-143	-121	-120	-135	0.544	6.013	37.44	181.4	6.212	1.282
G-10	304	316	316	308	0.49	6.137	37.19	179.4	-5.611	-1.163
G-11	-226	-210	-210	-220	0.56	5.968	38.11	184.6	8.143	1.681
G-12	147	159	159	151	0.576	5.744	36.21	173	4.882	1.022




Figure G-13. Manure belt house H₂S one-to-one plots models 1 through 6.




Figure G-14. Manure belt house H_2S one-to-one plots models 7 through 12.

Table G-19. Parameter and estimates for manure belt house PM₁₀ emission models tested.

Model	Parameter	Estimate	Standard Error	p-value
P-1	Intercept	6.631005	0.74268	<.0001
P-1	Inventory	0.007205	0.00304	0.0186
	Intercept	6.701076	0.76935	<.0001
P-2	Inventory	0.007479	0.00306	0.0153
	Ambient relative humidity	-0.001965	0.00392	0.6168
	Intercept	7.118863	0.83707	<.0001
P-3	Inventory	0.007319	0.00304	0.0168
	Exhaust relative humidity	-0.008078	0.00731	0.2704
	Intercept	6.916558	0.80465	<.0001
P-4	Inventory	0.007424	0.00306	0.016
P-4	Ambient relative humidity	-0.003651	0.00426	0.3923
	Ambient temperature	-0.006758	0.00683	0.3233
	Intercept	7.331233	0.86758	<.0001
P-5	Inventory	0.007218	0.00303	0.0181
P-5	Exhaust relative humidity	-0.009859	0.00751	0.1902
	Ambient temperature	-0.005743	0.00643	0.3728
	Intercept	8.177526	0.96556	<.0001
P-6	Inventory	0.010336	0.00316	0.0012
P-6	Ambient relative humidity	-0.005075	0.00404	0.2102
	Exhaust temperature	-0.073958	0.02775	0.0083
	Intercept	8.165323	0.97841	<.0001
P-7	Inventory	0.009413	0.00313	0.003
P-/	Exhaust relative humidity	-0.008534	0.00722	0.2386
	Exhaust temperature	-0.057587	0.02677	0.033
	Intercept	7.630943	1.11232	<.0001
P-8	Live animal weight	0.00319	0.00309	0.3041
P-0	Ambient temperature	-0.008317	0.00699	0.2353
	Ambient relative humidity	-0.003303	0.00431	0.4437
	Intercept	7.95699	1.07648	<.0001
P-9	Live animal weight	0.003787	0.0031	0.2245
F-9	Ambient temperature	-0.008271	0.00666	0.216
	Exhaust relative humidity	-0.0119	0.00775	0.1256
	Intercept	8.87744	1.1733	<.0001
P-10	Live animal weight	0.006543	0.00322	0.0444
P-10	Exhaust temperature	-0.090975	0.03098	0.0036
	Ambient relative humidity	-0.004612	0.00408	0.2588
	Intercept	8.890437	1.10805	<.0001
P-11	Live animal weight	0.006797	0.00329	0.0411
 L-TT	Exhaust temperature	-0.07933	0.03034	0.0096
	Exhaust relative humidity	-0.011355	0.00736	0.1241
P-12	Intercept	7.065611	2.94541	0.0182
L-12	Hen age	-0.001711	0.00041	<.0001

Model	Parameter	Estimate	Standard Error	p-value
	Inventory	0.008101	0.01154	0.4844
	Ambient relative humidity	-0.00372	0.0041	0.3646
	Intercept	6.435778	0.77396	<.0001
	F	-2.23531	1.07099	0.0376
P-13	M	-2.175865	1.11679	0.0522
P-13	Т	0	·	
	Inventory	0.017791	0.00573	0.0021
	Ambient relative humidity	-0.00269	0.00396	0.4973
	Intercept	6.648885	0.8072	<.0001
	F	-2.265321	1.06807	0.0346
	M	-2.265121	1.11403	0.0428
P-14	Т	0	•	
	Inventory	0.017891	0.00571	0.0019
	Ambient temperature	-0.007059	0.00688	0.3062
	Ambient relative humidity	-0.004377	0.00426	0.3052
	Intercept	6.705426	0.75615	<.0001
P-15	Inventory	0.007092	0.00305	0.0209
	Ambient temperature	-0.003622	0.00626	0.5633
	Intercept	7.687405	0.90053	<.0001
P-16	Inventory	0.009193	0.00312	0.0035
	Exhaust temperature	-0.058102	0.02672	0.031

Table G-20. Fit and evaluation statistics for the manure belt house PM₁₀ models tested.

						LNME ^a	NMEb	ME ^b	MB ^b	NMB ^b
Model	2LogL	AIC	AICc	BIC	Corr.	(%)	(%)	(g day ⁻¹)	(g day ⁻¹)	(%)
P-1	1,590	1,602	1,602	1,594	0.166	9.608	85.2	4,620	-61.72	-1.138
P-2	1,580	1,594	1,594	1,585	0.174	9.603	84.9	4,624	-76.38	-1.403
P-3	1,569	1,583	1,583	1,574	0.174	9.625	84.61	4,574	-77.96	-1.442
P-4	1,579	1,595	1,595	1,585	0.2	9.597	84.89	4,623	-48.7	-0.894
P-5	1,568	1,584	1,584	1,574	0.196	9.616	84.76	4,582	-51.06	-0.945
P-6	1,574	1,590	1,590	1,579	0.283	9.291	83.43	4,543	-22.48	-0.413
P-7	1,565	1,581	1,581	1,570	0.264	9.384	83.9	4,535	-32.44	-0.6
P-8	1,574	1,590	1,590	1,579	0.148	9.806	86.29	4,722	-24.61	-0.45
P-9	1,562	1,578	1,578	1,568	0.153	9.827	85.7	4,655	-39.03	-0.718
P-10	1,568	1,584	1,584	1,573	0.271	9.482	83.79	4,585	-48.54	-0.887
P-11	1,558	1,574	1,574	1,563	0.259	9.544	83.52	4,537	-63.89	-1.176
P-12	1,209	1,225	1,225	1,214	0.386	8.113	73.18	3,295	58.71	1.304
P-13	1,576	1,594	1,594	1,582	0.247	9.361	83.65	4,555	-101.5	-1.864
P-14	1,575	1,595	1,595	1,582	0.268	9.361	83.51	4,548	-74.76	-1.373
P-15	1,590	1,604	1,604	1,595	0.178	9.61	85.29	4,624	-40.12	-0.74
P-16	1,586	1,600	1,600	1,591	0.253	9.374	84.4	4,576	-0.421	-0.008




Figure G-15. Manure belt house PM₁₀ one-to-one plots models 1 through 8.




Figure G-16. Manure belt house PM₁₀ one-to-one plots models 9 through 16.

Table G-21. Parameter and estimates for manure belt house PM_{2.5} emission models tested.

Model	Parameter	Estimate	Standard Error	p-value
P-1	Intercept	-127.4489	61.0184	0.0681
P-T	Inventory	0.534577	0.24656	0.0604
	Intercept	-126.759	60.9479	0.0904
P-2	Inventory	0.531631	0.24755	0.0819
	Ambient relative humidity	0.000565	0.03137	0.9858
	Intercept	37.035887	29.4546	0.2794
P-3	Inventory	-0.094695	0.11911	0.474
	Exhaust relative humidity	-0.128115	0.03386	0.0049
	Intercept	-117.9991	89.9341	0.2219
P-4	Inventory	0.495946	0.36299	0.2041
P-4	Ambient relative humidity	0.002538	0.03018	0.9336
	Ambient temperature	-0.01164	0.0707	0.8712
	Intercept	-133.8634	83.2194	0.1316
P-5	Inventory	0.591108	0.33948	0.1046
P-5	Exhaust relative humidity	-0.115588	0.07929	0.1596
	Ambient temperature	-0.026486	0.06113	0.67
	Intercept	-121.6364	85.7067	0.1828
P-6	Inventory	0.513706	0.32419	0.143
P-6	Ambient relative humidity	0.001609	0.03052	0.9583
	Exhaust temperature	-0.029674	0.33459	0.9305
	Intercept	-131.4002	80.2027	0.1237
P-7	Inventory	0.585801	0.30869	0.0795
P-/	Exhaust relative humidity	-0.096143	0.06461	0.1506
	Exhaust temperature	-0.099247	0.29081	0.7373
	Intercept	15.313806	17.5534	0.4379
P-8	Live animal weight	-0.023061	0.05245	0.6874
P-0	Ambient temperature	-0.027824	0.0518	0.6043
	Ambient relative humidity	-0.028845	0.03358	0.3982
	Intercept	35.229203	11.5459	0.0059
P-9	Live animal weight	-0.050831	0.02658	0.0715
P-3	Ambient temperature	-0.045538	0.03403	0.1936
	Exhaust relative humidity	-0.174683	0.04196	0.0004
	Intercept	16.694338	10.5833	0.2096
P-10	Live animal weight	-0.019203	0.03677	0.6328
F-10	Exhaust temperature	-0.107616	0.21044	0.624
	Ambient relative humidity	-0.030398	0.03209	0.3531
	Intercept	33.520989	12.4481	0.0166
P-11	Live animal weight	-0.037796	0.02139	0.104
₋₁₁	Exhaust temperature	-0.19216	0.16949	0.2727
	Exhaust relative humidity	-0.148561	0.0359	0.0009
P-12	Intercept	-127.1099	54.6499	0.0449
F-12	Hen age	0.000199	0.0049	0.9701

Model	Parameter	Estimate	Standard Error	p-value
	Inventory	0.516362	0.21867	0.0419
	Ambient relative humidity	0.056541	0.02641	0.0536
	Intercept	-94.28449	51.693	0.0942
	F	-1.46145	1.06395	0.1939
P-13	M	0	•	
	Inventory	0.411612	0.20817	0.0726
	Ambient relative humidity	-0.024659	0.03084	0.4305
	Intercept	-96.14378	76.8941	0.2282
	F	-1.475749	1.37378	0.3033
P-14	M	0	•	
P-14	Inventory	0.419524	0.31091	0.195
	Ambient temperature	0.004319	0.07322	0.9543
	Ambient relative humidity	-0.026112	0.03459	0.4569
	Intercept	-118.1904	87.2564	0.199
P-15	Inventory	0.497444	0.3514	0.1809
	Ambient temperature	-0.011043	0.06202	0.8608
	Intercept	-119.6086	85.0107	0.1815
P-16	Inventory	0.507481	0.3233	0.1408
	Exhaust temperature	-0.044218	0.3062	0.8869

Table G-22. Fit and evaluation statistics for the manure belt house PM_{2.5} models tested.

						LNME	NME ^b	ME ^b	MBb	NMB ^b
Model	2LogL	AIC	AICc	BIC	Corr.	(%)	(%)	(g day ⁻¹)	(g day ⁻¹)	(%)
P-1	108	120	123	112	0.216	24.72	158.3	485.9	322.4	105.1
P-2	104	118	123	109	0.229	24.48	153.1	483	314.4	99.61
P-3	103	117	122	108	0.326	19.45	152	466.3	202.7	66.04
P-4	104	120	127	110	0.201	25.86	172.6	544.7	393.5	124.7
P-5	101	117	123	106	0.366	27.26	486.8	1536	1415	448.3
P-6	104	120	127	110	0.221	25.05	159.3	502.6	344.4	109.1
P-7	104	120	126	110	0.356	26.5	362	1111	990.8	322.9
P-8	106	122	128	112	0.326	22.59	91.43	288.5	44.7	14.16
P-9	97	113	119	102	0.514	18.95	175.4	553.6	415.2	131.6
P-10	106	122	128	112	0.351	22.05	89.05	281	35.57	11.27
P-11	102	118	124	108	0.514	16.93	116	356	197.7	64.42
P-12	83	99	107	88	0.105	31.62	385.4	1141	1021	345
P-13	101	117	124	107	0.424	22.27	150.6	475.3	337.3	106.9
P-14	101	119	128	108	0.441	21.66	141.2	445.5	305.2	96.72
P-15	104	118	123	109	0.209	25.63	168.4	531.5	375	118.8
P-16	108	122	126	112	0.206	25.58	166.8	511.8	361.7	117.9




Figure G-17. Manure belt house PM_{2.5} one-to-one plots models 1 through 8.




Figure G-18. Manure belt house PM_{2.5} one-to-one plots models 9 through 16.

Table G-23. Parameter and estimates for manure belt house TSP emission models tested.

Model	Parameter	Estimate	Standard Error	p-value
P-1	Intercept	6.936206	8.87165	0.4404
P-I	Inventory	0.00987	0.03594	0.7855
P-2	Intercept	5.99155	9.72176	0.5423
	Inventory	0.01539	0.0404	0.706
	Ambient relative humidity	-0.005782	0.01267	0.6515
	Intercept	5.400088	9.93986	0.5915
P-3	Inventory	0.014807	0.03983	0.7129
	Exhaust relative humidity	0.004945	0.02397	0.8381
	Intercept	0.931643	10.4965	0.9299
P-4	Inventory	0.032681	0.04235	0.4466
P-4	Ambient relative humidity	0.001295	0.01358	0.9247
	Ambient temperature	0.02696	0.02172	0.2243
	Intercept	-0.557657	10.6841	0.9588
٦.	Inventory	0.039052	0.04225	0.3639
P-5	Exhaust relative humidity	0.000474	0.0226	0.9834
	Ambient temperature	0.026653	0.01919	0.176
	Intercept	4.570047	11.8127	0.7017
D. C	Inventory	0.018506	0.0432	0.6716
P-6	Ambient relative humidity	-0.00509	0.01302	0.6987
	Exhaust temperature	0.022693	0.10278	0.827
	Intercept	3.270848	11.9588	0.7867
P-7	Inventory	0.019968	0.04294	0.6458
P-/	Exhaust relative humidity	0.004089	0.0239	0.8654
	Exhaust temperature	0.034352	0.10044	0.7354
	Intercept	10.02535	5.38972	0.0744
P-8	Live animal weight	-0.003082	0.01507	0.8398
P-8	Ambient temperature	0.01859	0.02478	0.4581
	Ambient relative humidity	0.003044	0.01382	0.8271
	Intercept	9.484187	5.35837	0.0889
P-9	Live animal weight	-0.000969	0.01539	0.9503
P-9	Ambient temperature	0.017318	0.02344	0.465
	Exhaust relative humidity	0.000939	0.02389	0.9689
	Intercept	12.886557	5.97738	0.0414
D 10	Live animal weight	-0.009111	0.01332	0.5007
P-10	Exhaust temperature	-0.010104	0.10065	0.9207
	Ambient relative humidity	-0.00111	0.01328	0.9339
	Intercept	11.921498	5.94847	0.0558
D 11	Live animal weight	-0.008098	0.01283	0.5329
P-11	Exhaust temperature	-0.004938	0.10048	0.9611
	Exhaust relative humidity	0.006196	0.02451	0.8022
D 13	Intercept	26.866895	13.2367	0.0528
P-12	Hen age	-0.003326	0.0014	0.0262

Model	Parameter	Estimate	Standard Error	p-value
	Inventory	-0.066133	0.05325	0.2254
	Ambient relative humidity	-0.004948	0.01273	0.7002
	Intercept	5.178835	9.95086	0.6065
	F	0.351267	0.93894	0.7148
P-13	M	0		
	Inventory	0.017207	0.04058	0.6746
	Ambient relative humidity	-0.005463	0.01268	0.6696
	Intercept	0.033107	10.7166	0.9976
	F	0.397099	0.97067	0.6898
P-14	M	0		
P-14	Inventory	0.034641	0.04251	0.4217
	Ambient temperature	0.027125	0.02169	0.221
	Ambient relative humidity	0.001645	0.01358	0.9044
	Intercept	0.948768	9.75704	0.9232
P-15	Inventory	0.033059	0.03914	0.4051
	Ambient temperature	0.025805	0.01899	0.1847
	Intercept	4.719899	10.9951	0.6709
P-16	Inventory	0.015134	0.03922	0.7024
	Exhaust temperature	0.034521	0.09845	0.7286

Table G-24. Fit and evaluation statistics for the manure belt house TSP models tested.

						LNME	NME ^b	ME ^b	MB ^b	NMB ^b
Model	2LogL	AIC	AICc	BIC	Corr.	(%)	(%)	(g day ⁻¹)	(g day ⁻¹)	(%)
P-1	238	250	252	243	0.074	9.95	78.29	11,670	82.75	0.555
P-2	233	247	249	238	0.085	10.020	78.37	11,760	-41.4	-0.276
P-3	232	246	248	237	0.12	10.07	78.03	11,800	-5.875	-0.039
P-4	232	248	250	237	0.041	10.09	76.75	11,520	-241.80	-1.612
P-5	230	246	249	236	0.056	10.1	75.66	11,450	-239	-1.580
P-6	233	249	252	239	0.068	10.06	78.53	11,780	-93.59	-0.624
P-7	232	248	251	238	0.088	10.08	78.22	11,830	-98.2	-0.649
P-8	232	248	251	238	-0.02	10.02	76.68	11,500	-180	-1.199
P-9	231	247	250	237	-0.017	10.01	75.92	11,490	-270.6	-1.789
P-10	233	249	251	238	0.008	10.090	80.19	12,030	120.4	0.803
P-11	232	248	250	237	0.024	10.1	78.76	11,910	-31.93	-0.211
P-12	226	242	245	232	0.32	10.04	68.49	10,370	-191.3	-1.264
P-13	233	249	252	239	0.07	9.979	77.61	11,640	-113.2	-0.76
P-14	231	249	253	238	0.043	9.936	75.66	11,350	-323.2	-2.154
P-15	237	251	252	241	0.044	9.941	75.94	11,320	-205.4	-1.378
P-16	238	252	254	243	0.043	9.958	78.36	11,680	-17.93	-0.12




Figure G-19. Manure belt house TSP one-to-one plots models 1 through 8.




Figure G-20. Manure belt house TSP one-to-one plots models 9 through 16.

Table G-25. Parameter combinations tested as models for layer manure shed NH₃ and H₂S emissions.

Model	Parameter
G-1	Intercept, Inventory, Ambient temperature
G-2	Intercept, Inventory (5 day lag), Ambient temperature
G-3	Intercept, Live animal weight, Ambient temperature
G-4	Intercept, Live animal weight (5 day lag), Ambient temperature
G-5	Intercept, Inventory (5 day lag), Ambient temperature, Ambient relative humidity
G-6	Intercept, Inventory (5 day lag), Ambient temperature, Wind speed
G-7	Intercept, Inventory (5 day lag), Ambient temperature, Ambient relative humidity, Wind speed
G-8	Intercept, Inventory (5 day lag), Ambient temperature, Average hen age
G-9	Intercept, Inventory (5 day lag), Ambient temperature, Management phase ((CFF), (FE), (FF),
G-3	(FM), (FT), and (MF))
G-10	Intercept, Inventory (5 day lag), Ambient temperature, Manure age
G-11	Intercept, Inventory (5 day lag), Wind speed
G-12	Intercept, Inventory (5 day lag), Manure age
G-13	Intercept, Inventory (5 day lag), Wind speed, Manure age
G-14	Intercept, Inventory (5 day lag), Ambient relative humidity, Manure age
G-15	Intercept, Inventory (5 day lag), Airflow
G-16	Intercept, Ambient temperature, Airflow
G-17	Intercept, Airflow, manure age
G-18	Intercept, Inventory (5 day lag), Manure age, Management phase (CFF, FE, FF, FM, FT, and MF)
G-19	Intercept, Ambient Temperature, Manure age, Management phase (CFF, FE, FF, FM, FT, and MF)
G-20	Intercept, Inventory (5 day lag)

Table G-26. Parameter combinations tested as models for layer manure shed PM_{10} , $PM_{2.5}$, and TSP emissions.

Model	Parameter
P-1	Intercept, Inventory, Airflow
P-2	Intercept, Inventory (5 day lag), Airflow
P-3	Intercept, Live animal weight, Airflow
P-4	Intercept, Live animal weight (5 day lag), Airflow
P-5	Intercept, Inventory (5 day lag), Wind speed
P-6	Intercept, Inventory (5 day lag), Ambient temperature
P-7	Intercept, Inventory (5 day lag), Ambient relative humidity
P-8	Intercept, Inventory (5 day lag), Average hen age
P-9	Intercept, Inventory (5 day lag), Management phase (CFF, FE, FF, FM, FT, and MF)
P-10	Intercept, Inventory (5 day lag), Manure age
P-11	Intercept, Inventory (5 day lag),
P-12	Intercept, Live animal weight (5 day lag)
P-13	Intercept, Airflow

Table G-27. Parameter and estimates for layer manure shed NH_3 emission models tested.

Model	Parameter	Estimate	Standard Error	p-value
	Intercept	-0.093535	0.53554	0.8615
G-1	Inventory	0.003706	0.00109	0.0008
	Ambient Temperature	-0.013247	0.00385	0.0007
	Intercept	-0.194945	0.5268	0.7116
G-2	Inventory (5 day lag)	0.003927	0.00108	0.0003
	Ambient Temperature	-0.013752	0.00385	0.0004
	Intercept	0.77162	0.72672	0.2893
G-3	Live animal weight	0.001413	0.00106	0.1819
	Ambient Temperature	-0.014411	0.00393	0.0003
	Intercept	1.607649	0.72587	0.0276
G-4	Live animal weight (5 day lag)	0.000211	0.00105	0.8415
	Ambient Temperature	-0.014591	0.00396	0.0003
	Intercept	-0.200801	0.5544	0.7175
C -	Inventory (5 day lag)	0.003926	0.00108	0.0003
G-5	Ambient Temperature	-0.013697	0.00417	0.0011
	Ambient relative humidity	0.000087	0.00257	0.973
	Intercept	-0.204953	0.52685	0.6976
6.6	Inventory (5 day lag)	0.003998	0.00108	0.0003
G-6	Ambient Temperature	-0.014176	0.00389	0.0003
	Wind speed	-0.028035	0.03929	0.4759
	Intercept	-0.234691	0.55617	0.6733
	Inventory (5 day lag)	0.003993	0.00108	0.0003
G-7	Ambient Temperature	-0.013919	0.00418	0.001
	Wind speed	-0.029245	0.03996	0.4646
	Ambient relative humidity	0.000435	0.00261	0.8677
	Intercept	-2.184307	2.10806	0.3013
6.0	Inventory (5 day lag)	0.008211	0.0041	0.0465
G-8	Ambient Temperature	-0.01487	0.00431	0.0007
	Average hen age	0.000045	0.00046	0.9215
	Intercept	1.365736	0.92967	0.1426
	CFF	-1.127859	0.45398	0.0134
	FE	-1.085898	0.61032	0.0759
	FF	-0.761009	0.19641	0.0001
G-9	FM	-0.21515	0.25873	0.4063
	FT	-1.19297	0.364	0.0011
	MF	0	•	
	Inventory (5 day lag)	0.001975	0.00183	0.2807
	Ambient Temperature	-0.005749	0.00421	0.173
	Intercept	-0.502174	0.52469	0.3394
C 40	Inventory (5 day lag)	0.003861	0.00106	0.0003
G-10	Ambient Temperature	-0.007305	0.00414	0.0786
	Manure age	0.001224	0.00032	0.0001
	Intercept	-0.353768	0.54826	0.5193
G-11	Inventory (5 day lag)	0.003864	0.00113	0.0007
	Wind speed	-0.009488	0.0388	0.8069

Model	Parameter	Estimate	Standard Error	p-value
	Intercept	-0.619946	0.53464	0.2472
G-12	Inventory (5 day lag)	0.003821	0.00109	0.0005
	Manure age	0.001395	0.00029	<.0001
	Intercept	-0.645153	0.52975	0.2243
C 13	Inventory (5 day lag)	0.003875	0.00108	0.0004
G-13	Manure age	0.001469	0.0003	<.0001
	Wind speed	-0.026953	0.03834	0.4825
	Intercept	-0.789483	0.54424	0.1479
6.44	Inventory (5 day lag)	0.003771	0.00107	0.0005
G-14	Manure age	0.001434	0.00029	<.0001
	Ambient relative humidity	0.00275	0.00234	0.2414
	Intercept	-0.383165	0.47248	0.418
G-15	Airflow	0.025665	0.00166	<.0001
	Inventory (5 day lag)	0.003203	0.00097	0.0011
	Intercept	1.35475	0.06234	<.0001
G-16	Airflow	0.026011	0.00172	<.0001
	Ambient Temperature	-0.013239	0.00335	<.0001
	Intercept	0.930519	0.06529	<.0001
G-17	Airflow	0.025631	0.00166	<.0001
	Manure age	0.001198	0.00026	<.0001
	Intercept	1.10654	0.91811	0.2288
	CFF	-0.677203	0.45526	0.1377
	FE	-1.109011	0.60165	0.0659
	FF	-0.650149	0.18947	0.0007
G-18	FM	-0.004042	0.2595	0.9876
	FT	-1.164419	0.35849	0.0013
	MF	0		
	Inventory (5 day lag)	0.001621	0.00181	0.3712
	Manure age	0.001206	0.00029	<.0001
	Intercept	1.889412	0.21118	<.0001
	CFF	-0.759612	0.45799	0.098
	FE	-1.43337	0.48585	0.0033
	FF	-0.659925	0.1943	0.0008
G-19	FM	-0.05961	0.25373	0.8144
	FT	-1.390809	0.25771	<.0001
	MF	0		
	Manure age	0.001276	0.00032	<.0001
	Ambient Temperature	0.000626	0.00446	0.8883
G-20	Intercept	-0.365539	0.55253	0.5087
G-20	Inventory (5 day lag)	0.003885	0.00113	0.0007

Table G-28. Fit and evaluation statistics for the layer manure shed NH₃ models tested.

						LNME	NME ^b	ME ^b	MB ^b	NMB ^b
Model	2LogL	AIC	AICc	BIC	Corr.	(%)	(%)	(kg day ⁻¹)	(kg day ⁻¹)	(%)
G-1	765	775	775	796	0.361	32.65	72.64	3.45	-0.084	-1.763
G-2	763	773	773	794	0.367	32.47	72.45	3.442	-0.086	-1.814
G-3	745	755	755	776	0.335	32.2	72.32	3.542	-0.063	-1.28
G-4	746	756	756	777	0.334	32.38	72.81	3.567	-0.051	-1.035
G-5	763	775	775	800	0.367	32.47	72.47	3.442	-0.086	-1.813
G-6	763	775	775	800	0.366	32.49	72.44	3.441	-0.086	-1.803
G-7	763	777	777	806	0.365	32.490	72.48	3.443	-0.086	-1.801
G-8	502	514	514	537	0.394	28.48	65	3.429	-0.098	-1.855
G-9	739	759	759	801	0.411	31.81	71.52	3.397	-0.027	-0.569
G-10	748	760	760	785	0.389	31.69	71.14	3.379	-0.051	-1.07
G-11	774	784	784	805	0.199	34.29	76.34	3.626	-0.033	-0.705
G-12	784	794	794	815	0.326	32.45	72.98	3.462	-0.019	-0.405
G-13	751	763	763	788	0.339	32.37	72.38	3.438	-0.023	-0.482
G-14	750	762	762	787	0.356	32.12	72.15	3.427	-0.036	-0.76
G-15	609	619	620	641	0.551	29.68	60.36	2.863	-0.193	-4.068
G-16	587	597	598	618	0.599	28.47	57.31	2.722	-0.272	-5.725
G-17	599	609	609	630	0.57	29.09	58.01	2.752	-0.235	-4.949
G-18	756	776	776	818	0.439	30.93	69.82	3.312	-0.007	-0.153
G-19	724	744	745	786	0.442	30.93	69.94	3.322	0.006	0.129
G-20	806	814	814	831	0.196	34.11	76.38	3.624	-0.032	-0.665




Figure G-21. Layer manure shed NH₃ one-to-one plots models 1 through 9.




Figure G-22. Layer manure shed NH₃ one-to-one plots models 10 through 18.




Figure G-23. Layer manure shed NH₃ one-to-one plots models 19 through 27.

Table G-29. Parameter and estimates for layer manure shed H_2S emission models tested.

Model	Parameter	Estimate	Standard Error	p-value
	Intercept	0.755232	0.60942	0.2163
G-1	Inventory	0.006078	0.00125	<.0001
	Ambient Temperature	-0.023815	0.00448	<.0001
	Intercept	1.295775	0.60993	0.0345
G-2	Inventory (5 day lag)	0.004976	0.00125	<.0001
	Ambient Temperature	-0.024164	0.00454	<.0001
	Intercept	2.764473	0.84436	0.0012
G-3	Live animal weight	0.001413	0.00123	0.2532
	Ambient Temperature	-0.023989	0.00471	<.0001
	Intercept	4.186178	0.84953	<.0001
G-4	Live animal weight (5 day lag)	-0.000664	0.00124	0.5924
	Ambient Temperature	-0.022925	0.00477	<.0001
	Intercept	1.263787	0.64345	0.0504
G-5	Inventory (5 day lag)	0.004968	0.00125	<.0001
u-5	Ambient Temperature	-0.02389	0.00488	<.0001
	Ambient relative humidity	0.000476	0.00305	0.876
	Intercept	1.274465	0.61052	0.0377
G-6	Inventory (5 day lag)	0.005129	0.00126	<.0001
U-0	Ambient Temperature	-0.025048	0.0046	<.0001
	Wind speed	-0.062217	0.0471	0.1872
	Intercept	1.191004	0.6457	0.0661
	Inventory (5 day lag)	0.005117	0.00126	<.0001
G-7	Ambient Temperature	-0.024391	0.0049	<.0001
	Wind speed	-0.065576	0.04787	0.1714
	Ambient relative humidity	0.001225	0.00309	0.6922
	Intercept	2.142455	2.53874	0.3998
G-8	Inventory (5 day lag)	0.004477	0.00495	0.367
U-0	Ambient Temperature	-0.026488	0.0053	<.0001
	Average hen age	-0.001243	0.00055	0.0255
	Intercept	1.95062	1.09308	0.0751
	CFF	-1.101759	0.55353	0.0472
	FE	-0.65747	0.72742	0.3665
	FF	-0.834148	0.23239	0.0004
G-9	FM	-0.101993	0.30247	0.7362
	FT	-0.86526	0.43079	0.0453
	MF	0		
	Inventory (5 day lag)	0.004885	0.00215	0.0236
	Ambient Temperature	-0.01348	0.00509	0.0085
	Intercept	1.137439	0.61664	0.0661
G-10	Inventory (5 day lag)	0.004942	0.00125	<.0001
0-10	Ambient Temperature	-0.020719	0.00496	<.0001
	Manure age	0.000624	0.00037	0.0924
	Intercept	1.133939	0.6545	0.0842
G-11	Inventory (5 day lag)	0.004602	0.00134	0.0007
	Wind speed	-0.032384	0.04695	0.4907

Model	Parameter	Estimate	Standard Error	p-value
	Intercept	0.92442	0.64839	0.1549
G-12	Inventory (5 day lag)	0.004517	0.00132	0.0007
	Manure age	0.001147	0.00035	0.0012
	Intercept	0.846057	0.64637	0.1915
C 13	Inventory (5 day lag)	0.004695	0.00132	0.0004
G-13	Manure age	0.001281	0.00036	0.0004
	Wind speed	-0.046784	0.04675	0.3175
	Intercept	0.553938	0.66249	0.4037
C 11	Inventory (5 day lag)	0.004553	0.0013	0.0005
G-14	Manure age	0.001229	0.00035	0.0005
	Ambient relative humidity	0.005119	0.00287	0.0753
	Intercept	1.108613	0.59271	0.0623
G-15	Airflow	0.026849	0.00212	<.0001
	Inventory (5 day lag)	0.003873	0.00122	0.0016
	Intercept	3.321478	0.08027	<.0001
G-16	Airflow	0.02715	0.00218	<.0001
	Ambient Temperature	-0.021926	0.00417	<.0001
	Intercept	2.818123	0.08072	<.0001
G-17	Airflow	0.026809	0.00213	<.0001
	Manure age	0.000895	0.00033	0.0063
	Intercept	1.8084	1.10569	0.1027
	CFF	-0.830047	0.55418	0.1349
	FE	-0.863555	0.72612	0.2349
	FF	-0.925039	0.22826	<.0001
G-18	FM	0.066221	0.31199	0.832
	FT	-0.972908	0.43179	0.0248
	MF	0		
	Inventory (5 day lag)	0.004587	0.00218	0.0356
	Manure age	0.000805	0.00035	0.0211
	Intercept	4.145601	0.25269	<.0001
	CFF	-0.940113	0.56368	0.0961
	FE	-1.621421	0.58794	0.0061
	FF	-0.810388	0.23545	0.0007
G-19	FM	-0.13455	0.30412	0.6585
	FT	-1.514196	0.30967	<.0001
	MF	0		•
	Manure age	0.00067	0.00038	0.0758
	Ambient Temperature	-0.009705	0.00551	0.0789
6.30	Intercept	1.170893	0.65633	0.0754
G-20	Inventory (5 day lag)	0.004474	0.00135	0.001

Table G-30. Fit and evaluation statistics for the layer manure shed H₂S models tested.

						LNME ^a	NME ^b	ME ^b	MBb	NMB ^b
Model	2LogL	AIC	AICc	BIC	Corr.	(%)	(%)	(g day ⁻¹)	(g day ⁻¹)	(%)
G-1	956	966	966	987	0.444	16.93	73.49	26.28	-1.4	-3.915
G-2	963	973	973	994	0.423	17.04	73.72	26.36	-1.37	-3.829
G-3	928	938	939	959	0.386	17.02	74.49	27.39	-1.041	-2.831
G-4	933	943	944	964	0.377	17.28	74.84	27.46	-0.999	-2.723
G-5	963	975	975	1001	0.423	17.04	73.75	26.38	-1.377	-3.849
G-6	962	974	974	999	0.42	17.08	73.73	26.37	-1.348	-3.768
G-7	961	975	976	1005	0.42	17.090	73.79	26.39	-1.366	-3.82
G-8	614	626	627	649	0.495	15.92	67.6	27.84	-1.576	-3.827
G-9	943	963	964	1006	0.451	16.82	73.52	26.29	-0.537	-1.502
G-10	960	972	973	998	0.418	16.99	73.50	26.29	-1.297	-3.627
G-11	988	998	998	1019	0.184	18.7	82.25	29.42	-0.264	-0.737
G-12	1015	1025	1025	1046	0.258	18.07	79.97	28.76	-0.515	-1.432
G-13	975	987	987	1012	0.264	18.07	78.66	28.13	-0.536	-1.499
G-14	973	985	985	1010	0.31	17.8	77.34	27.66	-0.884	-2.471
G-15	886	896	896	917	0.479	16.35	67.59	24.31	-1.73	-4.811
G-16	843	853	853	874	0.566	15.45	59.89	21.42	-2.664	-7.449
G-17	888	898	899	920	0.47	16.46	66.01	23.74	-2.125	-5.908
G-18	981	1001	1001	1043	0.417	17.05	75.06	26.99	-0.442	-1.229
G-19	945	965	966	1008	0.434	16.93	74.08	26.49	-0.186	-0.52
G-20	1025	1033	1033	1051	0.187	18.67	83.12	29.89	-0.238	-0.661




Figure G-24. Layer manure shed H₂S one-to-one plots models 1 through 10.




Figure G-25. Layer manure shed H₂S one-to-one plots models 11 through 19.




Figure G-26. Layer manure shed H₂S one-to-one plots models 20 through 27.

Table G-31. Parameter and estimates for layer manure shed PM₁₀ emission models tested.

Model	Parameter	Estimate	Standard Error	p-value
	Intercept	4.069675	1.25809	0.0015
P-1	Inventory	0.000441	0.00258	0.8643
	Airflow	0.050735	0.0077	<.0001
	Intercept	4.530912	1.13681	0.0001
P-2	Inventory (5 day lag)	-0.000509	0.00233	0.8273
	Airflow	0.050994	0.00771	<.0001
	Intercept	0.71194	1.69773	0.6757
P-3	Live animal weight	0.005127	0.00244	0.0376
	Airflow	0.048737	0.00777	<.0001
	Intercept	0.230996	1.75607	0.8956
P-4	Live animal weight (5 day lag)	0.005803	0.00252	0.0228
	Airflow	0.049074	0.00773	<.0001
	Intercept	4.617304	1.3095	0.0006
P-5	Inventory (5 day lag)	0.000414	0.00269	0.8779
	Wind speed	0.113139	0.11575	0.3292
	Intercept	4.557673	1.31245	0.0007
P-6	Inventory (5 day lag)	0.000511	0.00271	0.8504
	Ambient Temperature	0.005383	0.00907	0.5535
	Intercept	5.242112	1.41299	0.0003
P-7	Inventory (5 day lag)	0.00035	0.0027	0.8968
	Ambient relative humidity	-0.007909	0.00589	0.1804
	Intercept	-66.36761	74.762	0.377
P-8	Inventory (5 day lag)	0.134653	0.14143	0.3435
	Average hen age	0.01381	0.01456	0.3454
	Intercept	4.161863	1.6264	0.0116
	FF	-0.807121	1.09202	0.4608
P-9	FM	-1.766058	1.09532	0.1087
	FT	0	•	•
	Inventory (5 day lag)	0.003175	0.00496	0.523
	Intercept	4.343148	1.30873	0.0011
P-10	Inventory (5 day lag)	0.000766	0.00266	0.774
	Manure age	0.000939	0.00057	0.0993
P-11	Intercept	4.5366	1.31267	0.0007
1 -77	Inventory (5 day lag)	0.000732	0.00268	0.7853
P-12	Intercept	-0.684347	2.01647	0.7349
L-TZ	Live animal weight (5 day lag)	0.007952	0.00287	0.0065
P-13	Intercept	4.283943	0.11276	<.0001
L-13	Airflow	0.050837	0.00768	<.0001

Table G-32. Fit and evaluation statistics for the layer manure shed PM_{10} models tested.

						LNME	NMEb	ME ^b	MBb	NMB ^b
Model	2LogL	AIC	AICc	BIC	Corr.	(%)	(%)	(g day ⁻¹)	(g day ⁻¹)	(%)
P-1	737	747	747	765	0.409	15.22	85.51	147	-9.075	-5.279
P-2	737	747	747	765	0.41	15.2	85.43	146.9	-9.1	-5.294
P-3	725	735	735	752	0.429	15.15	86.01	148.5	-6.199	-3.59
P-4	724	734	734	752	0.433	15.1	86.13	148.7	-5.675	-3.287
P-5	774	784	784	802	0.077	16.43	104.7	180	0.623	0.362
P-6	774	784	785	802	0.036	16.41	103	177.1	-0.928	-0.54
P-7	773	783	783	801	0.071	16.48	102.8	176.8	-1.222	-0.711
P-8	507	517	518	533	0.067	16.68	104.7	154.7	-1.623	-1.098
P-9	770	782	782	803	0.156	16.29	102.9	176.9	-0.996	-0.579
P-10	772	782	782	800	0.105	16.47	105.2	180.8	2.566	1.493
P-11	775	783	783	797	0.021	16.48	104	178.8	-0.291	-0.169
P-12	759	767	767	782	0.193	16.45	103.6	178.9	0.766	0.443
P-13	737	745	745	759	0.409	15.22	85.49	147	-9.1	-5.294




Figure G-27. Layer manure shed PM₁₀ one-to-one plots models 1 through 7.




Figure G-28. Layer manure shed PM₁₀ one-to-one plots models 8 through 14.

Table G-33. Parameter and estimates for layer manure shed $PM_{2.5}$ emission models tested.

Model	Parameter	Estimate	Standard Error	p-value	
	Intercept	-14.84682	8.72133	0.1062	
P-1	Inventory	0.033562	0.01766	0.0736	
	Airflow	0.067728	0.00999	<.0001	
	Intercept	-14.83081	8.79651	0.1094	
P-2	Inventory (5 day lag)	0.033497	0.01779	0.0762	
	Airflow	0.067819	0.00999	<.0001	
	Intercept	-4.34117	4.60554	0.359	
P-3	Live animal weight	0.008795	0.00666	0.2036	
	Airflow	0.070025	0.01008	<.0001	
	Intercept	-2.158197	4.1939	0.6134	
P-4	Live animal weight (5 day lag)	0.005665	0.00608	0.3646	
	Airflow	0.070835	0.01017	<.0001	
	Intercept	-31.51962	15.7683	0.0601	
P-5	Inventory (5 day lag)	0.069985	0.03188	0.0407	
	Wind speed	-0.312909	0.43464	0.4776	
	Intercept	-23.98923	17.8262	0.195	
P-6	Inventory (5 day lag)	0.054672	0.03552	0.141	
	Ambient Temperature	-0.008898	0.02378	0.7126	
	Intercept	-27.46865	13.9511	0.0645	
P-7	Inventory (5 day lag)	0.064388	0.02812	0.0344	
	Ambient relative humidity	-0.022083	0.01865	0.2484	
	Intercept	490.18321	895.044	0.589	
P-8	Inventory (5 day lag)	-0.922294	1.70959	0.5945	
	Average hen age	-0.094642	0.15137	0.5377	
	Intercept	-26.39053	13.6003	0.0686	
P-9	FF	-0.906908	0.48512	0.0788	
P-9	MF	0	•		
	Inventory (5 day lag)	0.060338	0.02726	0.0404	
	Intercept	-24.92318	13.6853	0.0857	
P-10	Inventory (5 day lag)	0.055385	0.02762	0.0606	
	Manure age	0.002715	0.00142	0.0719	
D 11	Intercept	-30.57734	14.5816	0.0505	
P-11	Inventory (5 day lag)	0.067599	0.02933	0.0334	
D 13	Intercept	-5.178811	7.38577	0.4919	
P-12	Live animal weight (5 day lag)	0.011848	0.01065	0.2802	
P-13	Intercept	1.747983	0.24149	<.0001	
P-13	Airflow	0.071834	0.01021	<.0001	

Table G-34. Fit and evaluation statistics for the layer manure shed $PM_{2.5}$ models tested.

						LNME	NMEb	ME ^b	MBb	NMB ^b
Model	2LogL	AIC	AICc	BIC	Corr.	(%)	(%)	(g day ⁻¹)	(g day ⁻¹)	(%)
P-1	57	67	69	74	0.851	16.37	26.54	12.81	1.882	3.9
P-2	57	67	70	74	0.85	16.41	26.63	12.85	1.881	3.897
P-3	59	69	71	76	0.84	17.31	28.47	13.74	1.34	2.776
P-4	59	69	72	77	0.834	17.46	27.06	13.06	0.43	0.892
P-5	81	91	94	98	0.411	29.46	76.99	39.49	-6.222	-12.13
P-6	81	91	94	98	0.423	29.35	80.79	41.43	-6.569	-12.81
P-7	80	90	93	97	0.482	27.94	65.18	33.43	-4.768	-9.297
P-8	51	59	61	64	0.719	21.17	54.34	16.17	-0.486	-1.634
P-9	84	94	96	101	0.563	27.89	78.72	38	-4.163	-8.626
P-10	84	94	96	101	0.567	27.8	78.66	37.97	-4.023	-8.336
P-11	87	95	96	101	0.452	30.33	79.34	38.29	-6.885	-14.27
P-12	90	98	100	104	0.24	33.6	91.71	44.26	-5.069	-10.5
P-14	60	68	70	74	0.826	16.62	31.96	15.42	-0.695	-1.44




Figure G-29. Layer manure shed PM_{2.5} one-to-one plots models 1 through 7.




Figure G-30. Layer manure shed PM_{2.5} one-to-one plots models 7 through 14.

Table G-35. Parameter and estimates for layer manure shed TSP emission models tested.

Model	Parameter	Estimate	Standard Error	p-value
	Intercept	0.573094	12.2988	0.9636
P-1	Inventory	0.006529	0.02523	0.8001
	Airflow	0.096594	0.03436	0.0115
	Intercept	0.45394	12.3709	0.9713
P-2	Inventory (5 day lag)	0.006766	0.02535	0.7939
	Airflow	0.096613	0.03436	0.0115
	Intercept	5.424643	8.49308	0.5346
P-3	Live animal weight	-0.002454	0.01244	0.8468
	Airflow	0.096379	0.03437	0.0116
	Intercept	6.812837	8.29716	0.427
P-4	Live animal weight (5 day lag)	-0.004473	0.01209	0.7176
	Airflow	0.095721	0.03427	0.0119
	Intercept	4.283506	15.8341	0.7917
P-5	Inventory (5 day lag)	0.001656	0.03259	0.9604
	Wind speed	0.081591	0.35102	0.8195
	Intercept	3.895732	15.7494	0.8091
P-6	Inventory (5 day lag)	0.002475	0.03233	0.9403
	Ambient Temperature	0.003544	0.0278	0.9008
	Intercept	0.71152	16.7371	0.9668
P-7	Inventory (5 day lag)	0.007148	0.03345	0.8347
	Ambient relative humidity	0.01393	0.0231	0.5554
	Intercept	-1188.806	960.653	0.2399
P-8	Inventory (5 day lag)	2.253253	1.81283	0.2379
	Average hen age	0.236659	0.19058	0.2384
	Intercept	4.041666	15.7641	0.8022
P-9	FF	0	·	
	Inventory (5 day lag)	0.002286	0.03241	0.945
	Intercept	-4.789205	17.7638	0.7922
P-10	Inventory (5 day lag)	0.020979	0.03682	0.5797
	Manure age	-0.001618	0.0017	0.3612
D 11	Intercept	4.041666	15.7641	0.8022
P-11	Inventory (5 day lag)	0.002286	0.03241	0.945
D 43	Intercept	9.118039	10.5292	0.4043
P-12	Live animal weight (5 day lag)	-0.005805	0.01541	0.7133
D 43	Intercept	3.752372	0.53756	<.0001
P-13	Airflow	0.096097	0.0344	0.0119

Table G-36. Fit and evaluation statistics for the layer manure shed TSP models tested.

						LNME	NMEb	ME ^b	MB ^b	NMB ^b
Model	2LogL	AIC	AICc	BIC	Corr.	(%)	(%)	(kg day ⁻¹)	(kg day ⁻¹)	(%)
P-1	75	85	88	90	0.415	20.19	95.71	303.7	61.59	19.41
P-2	75	85	88	90	0.415	20.2	95.77	303.9	61.69	19.44
P-3	75	85	88	91	0.414	20	92.14	292.4	54.35	17.13
P-4	75	85	88	90	0.416	19.86	90.65	287.6	51.84	16.34
P-5	81	91	95	97	0.1	20.46	88.18	279.8	1.329	0.419
P-6	81	91	95	97		20.62	88.91	282.1	1.228	0.387
P-7	81	91	95	97	0.165	20.73	87.74	278.4	4.4	1.387
P-8	80	90	94	96	0.201	20.33	85.05	269.8	-1.817	-0.573
P-9	81	89	92	94	0.005	20.63	88.5	280.8	1.141	0.36
P-10	81	91	94	96	0.147	20.28	88.04	279.3	0.192	0.06
P-11	81	89	92	94	0.005	20.63	88.5	280.8	1.141	0.36
P-12	81	89	92	94	0.059	19.95	86.11	273.2	-5.478	-1.727
P-14	75	83	85	87	0.414	20.03	93.53	296.8	57.7	18.19




Figure G-31. Layer manure shed TSP one-to-one plots models 1 through 7.




Figure G-32. Layer manure shed TSP one-to-one plots models 8 through 14.

Appendix H - Model Performance Evaluation

List of Tables

Table H-1. Model performance statistics, overall
Table H-2. Model performance statistics by house
Table H-3. Model performance statistics by season
Table H-4. Model performance statistics by house, by season
List of Figures
Figure H-1. Time series comparison of model (points) and observed (line) NH ₃ emissionsH-3
Figure H-2. Time series comparison of model (points) and observed (line) H ₂ S emissionsH-3
Figure H-3. Time series comparison of model (points) and observed (line) PM ₁₀ emissionsH-4
Figure H-4. Scatter plots of model versus observed emissions
Figure H-5. Scatter plots of model versus observed emissions, color coded by house
Figure H-6. Scatter plots of model versus observed NH ₃ emissions by season
Figure H-7. Scatter plots of model versus observed H ₂ S emissions by season
Figure H-8. Scatter plots of model versus observed PM ₁₀ emissions by season




Figure H-1. Time series comparison of model (points) and observed (line) NH₃ emissions.




Figure H-2. Time series comparison of model (points) and observed (line) H₂S emissions.




Figure H-3. Time series comparison of model (points) and observed (line) PM₁₀ emissions.

Table H-1. Model performance statistics, overall

Pollutant	n	MB (kg)	ME (kg)	NMB (%)	NME (%)	r
NH ₃	544	-164.32	204.96	-51%	64%	-0.38
H ₂ S	578	-0.05	0.24	-13%	64%	0.63
PM ₁₀	560	-7,542.73	7938.27	-53%	56%	0.37




Figure H-4. Scatter plots of model versus observed emissions.

Table H-2. Model performance statistics by house

Pollutant	House	n	MB (kg)	ME (kg)	NMB (%)	NME (%)	r
NH ₃	B13	273	-219.44	249.51	-59%	67%	-0.39
NH ₃	B14	271	-108.80	160.09	-41%	60%	-0.41
H ₂ S	B13	290	-0.08	0.27	-21%	67%	0.60
H ₂ S	B14	288	-0.01	0.22	-4%	61%	0.71
PM ₁₀	B13	282	-9923.98	10008.56	-60%	61%	0.39
PM ₁₀	B14	278	-5127.22	5838.18	-43%	49%	0.48

H-5




Figure H-5. Scatter plots of model versus observed emissions, color coded by house.

Table H-3. Model performance statistics by season

Pollutant	House	n	MB (kg)	ME (kg)	NMB (%)	NME (%	r
NH ₃	spring (MAM)	166	-137.85	220.95	-46%	74%	-0.49
NH ₃	summer (JJA)	68	6.36	99.02	3%	40%	0.55
NH ₃	autumn (SON)	156	-73.79	81.15	-30%	34%	0.40
NH ₃	winter (DJF)	154	-359.93	359.93	-79%	79%	-0.25
H₂S	spring (MAM)	166	-0.11	0.29	-25%	65%	0.59
H ₂ S	summer (JJA)	94	-0.35	0.48	-45%	61%	0.31
H ₂ S	autumn (SON)	154	0.04	0.12	12%	39%	0.65
H ₂ S	winter (DJF)	164	0.10	0.19	65%	116%	0.26
PM ₁₀	spring (MAM)	162	-6503.98	6734.67	-49%	50%	0.12
PM ₁₀	summer (JJA)	97	-12010.28	12010.28	-59%	59%	-0.11
PM ₁₀	autumn (SON)	135	-8742.36	9012.29	-57%	59%	0.38
PM ₁₀	winter (DJF)	166	-4970.31	5859.97	-47%	55%	-0.19




Figure H-6. Scatter plots of model versus observed NH₃ emissions by season.




Figure H-7. Scatter plots of model versus observed H₂S emissions by season.




Figure H-8. Scatter plots of model versus observed PM₁₀ emissions by season.

Table H-4. Model performance statistics by house, by season

Pollutant	House	season	n	MB	MGE	NMB	NMGE	r
NH ₃	B13	spring (MAM)	81	-189.02	263.65	-55%	77%	-0.47
NH ₃	B13	summer (JJA)	34	-47.78	106.98	-16%	35%	0.59
NH ₃	B13	autumn (SON)	80	-97.76	99.65	-36%	37%	0.26
NH ₃	B13	winter (DJF)	78	-450.65	450.65	-82%	82%	-0.39
NH ₃	B14	spring (MAM)	85	-89.08	180.25	-34%	70%	-0.50
NH ₃	B14	summer (JJA)	34	60.51	91.07	32%	48%	0.54
NH ₃	B14	autumn (SON)	76	-48.57	61.67	-23%	29%	0.56
NH ₃	B14	winter (DJF)	76	-266.84	266.84	-73%	73%	0.14
H ₂ S	B13	spring (MAM)	81	-0.16	0.32	-34%	68%	0.75
H ₂ S	B13	summer (JJA)	47	-0.44	0.59	-50%	67%	0.29
H ₂ S	B13	autumn (SON)	79	0.03	0.11	11%	35%	0.57
H ₂ S	B13	winter (DJF)	83	0.08	0.21	43%	115%	0.32
H ₂ S	B14	spring (MAM)	85	-0.06	0.26	-15%	61%	0.79
H ₂ S	B14	summer (JJA)	47	-0.26	0.37	-38%	54%	0.31
H ₂ S	B14	autumn (SON)	75	0.04	0.12	14%	42%	0.74
H ₂ S	B14	winter (DJF)	81	0.13	0.17	92%	117%	0.30
PM ₁₀	B13	spring (MAM)	82	-8690.21	8980.75	-57%	59%	0.55
PM ₁₀	B13	summer (JJA)	47	-12943.57	12943.57	-61%	61%	-0.11
PM ₁₀	B13	autumn (SON)	66	-10949.46	10949.89	-62%	62%	0.45
PM ₁₀	B13	winter (DJF)	87	-8677.62	8677.62	-61%	61%	0.15
PM ₁₀	B14	spring (MAM)	80	-4263.08	4432.45	-37%	38%	0.23
PM ₁₀	B14	summer (JJA)	50	-11132.98	11132.98	-58%	58%	-0.21
PM ₁₀	B14	autumn (SON)	69	-6631.23	7158.93	-50%	54%	0.26
PM ₁₀	B14	winter (DJF)	79	-887.58	2757.00	-13%	41%	-0.35