


Draft Risk Evaluation for 1,1-Dichloroethane

Systematic Review Supplemental File:

Data Quality Evaluation and Data Extraction Information for Dermal Absorption

CASRN: 75-34-3



July 2024

This supplemental file contains information regarding the data extraction and evaluation results for data sources that met the PECO screening criteria for the *Draft Risk Evaluation for 1,1-Dichloroethane* and were used to characterize dermal absorption. All studies that met the PECO for 1,1-Dichloroethane dermal absorption were conducted using 1,2-Dichloroethane. EPA conducted data quality evaluation based on author-reported descriptions and results; additional analyses (*e.g.*, statistical analyses performed during data integration into the risk evaluation) potentially conducted by EPA are not contained in this supplemental file. Key parameters and corresponding data for each condition were extracted from the reference. EPA used the TSCA systematic review process described in the *Draft Systematic Review Protocol Supporting TSCA Risk Evaluations for Chemical Substances* (also referred to as the '2021 Draft Systematic Review Protocol'). Any updated steps in the systematic review process since the publication of the 2021 Draft Systematic Review Protocol are described in the *Draft Risk Evaluation for 1,1-Dichloroethane - Systematic Review Protocol*.

To evaluate dermal absorption references, EPA consulted several OECD documents when considering quality rankings for individual metrics. Each condition (*e.g.*, individual concentrations tested or different experimental designs) is evaluated independently within a given reference. Therefore each reference may have more than one overall quality determination (OQD) to more appropriately reflect the quality of each condition. No OQD is determined for each reference as a whole, if it contains data from more than condition. A single reference may evaluate only a limited number of conditions (*e.g.*, use of only the neat compound). If all other methods and results are adequate, the study may be considered acceptable for certain conditions of use. However, the study may still be limited for use in the risk evaluation because it may not address other uses (*e.g.*, lower concentrations, certain solvents/diluents). Within the contents of this document, 1,1-dichloroethane may be referred to as the acronyms 1,1-DCA and 1,1-DCE. The acronyms 1,2-DCA, 1,2-DCE, and DCE refer to the chemical 1,2-dichloroethane. The acronyms 1,1,2-TCE, 1,1,2-TCA, and TCE refer to the chemical 1,1,2-trichloroethane. The acronym trans-1,2-DCE refers to the chemical trans-1,2-dichloroethylene. The acronym 1,2-DCP refers to the chemical 1,2-dichloropropane.

Table of Contents

HERO ID	Reference	Page
1,1-Dichloroethane		4
In vitro		
11396332	Labcorp Early Development, (2024). 1,1-Dichloroethane: Rates of penetration through human skin using a flow through in vitro system.	4
Isomer: 1,2-Dichloroethane		57
In vitro		
7490381	DuPont Haskell Lab, (2005). Ethylene dichloride: In vitro dermal absorption rate testing.	60
734137	Frasch, H. F., Barbero, A. M. (2009). A paired comparison between human skin and hairless guinea pig skin in vitro permeability and lag time measurements for 6 industrial chemicals. <i>Cutaneous and Ocular Toxicology</i> 28(3):107-113.	69
1313327	Frasch, H. F., Barbero, A. M., Alachkar, H., Mcdougal, J. N. (2007). Skin penetration and lag times of neat and aqueous diethyl phthalate, 1,1-Dichloroethane and naphthalene. <i>Cutaneous and Ocular Toxicology</i> 26(2):147-160.	77
4141541	Gajjar, R. M., Kasting, G. B. (2014). Absorption of ethanol, acetone, benzene and 1,1-Dichloroethane through human skin in vitro: a test of diffusion model predictions. <i>Toxicology and Applied Pharmacology</i> 281(1):109-117.	86
4940676	Schenk, L., Rauma, M., Fransson, M. N., Johanson, G. (2018). Percutaneous absorption of thirty-eight organic solvents in vitro using pig skin. <i>PLoS ONE</i> 13(10):e0205458.	96
In vivo - Animal		
94881	Jakobson, I., Wahlberg, J. E., Holmberg, B., Johansson, G. (1982). Uptake via the blood and elimination of 10 organic solvents following epicutaneous exposure of anesthetized guinea pigs. <i>Toxicology and Applied Pharmacology</i> 63(2):181-187.	104
200487	Morgan, D. L., Cooper, S. W., Carlock, D. L., Sykora, J. J., Sutton, B., Mattie, D. R., Mcdougal, J. N. (1991). Dermal absorption of neat and aqueous volatile organic chemicals in the Fischer 344 rat. <i>Environmental Research</i> 55(1):51-63.	112

Study Citation: Labcorp Early Development, (2024). 1,1-Dichloroethane: Rates of penetration through human skin using a flow through in vitro system.
HERO ID: 11396332

EXTRACTION**Parameter****Data**

Extraction ID; Chemical:	Infinite; 50% in IPM; 1,1-Dichloroethane-Parent compound
Skin Material/Species; Skin Preparation; Skin Thickness (um); Diffusion Cell Exposure Setup Type:	ex vivo human; Split thickness; 400; Flow-through; Notes: Not Reported
Occlusion Type; Donor Chamber Vehicle; Concentration of Test Substance in Vehicle (enter as percent):	Occluded; Isopropyl myristate; 50
Mass per Surface Area on Skin (mg/cm ²); Duration of Test Substance on Skin:	118; 24 hrs; Not Reported
Duration of Absorbance Measured; Frequency of Samples:	24 hrs; Samples were collected at 10 minutes, 30 minutes, 1 hour, 2 hours, and then every 2 hours; Notes: Not Reported
Time Skin was Washed and Method used; Radiolabel Presence:	Skin not washed; Yes
Total Recovery (percent); Dose Type:	Not Reported; Infinite
Percent Found in Skin Depot After Washing and Tape Stripping; Comments:	Not Reported; Notes: Not Reported
Percent Found in All Tape Strips, Excluding the Upper Two Strips; Comments:	Not Reported; Notes: Not Reported
Percent Found in Receptor Fluid and Receptor Fluid Rinse; Comments:	Not Reported; Notes: Not Reported
Total Percent Absorbed:	0
Steady State Permeability Coefficient (Kp) (cm/hr); Steady State Permeability Coefficient (Comments); Steady State Flux (ug/cm ² /hr); Steady State Flux (Comments); Maximum Permeability Coefficient (Kp) (cm/hr); Maximum Permeability Coefficient (Comments); Maximum Flux (ug/cm ² /hr); Maximum Flux (Comments):	0.00322; Notes: CV = 30.7%; Not Reported; Notes: Not Reported; Not Reported; Notes: Not Reported; 533; Notes: CV = 35.6%

EVALUATION

Domain	Metric	Rating	Comments
Domain 1: Test Substance			
	Metric 1:	High	Test substances were identified as non-radiolabeled 1,1-dichloroethane and radiolabeled 1,1- ¹⁴ C dichloroethane, CAS number: 75-34-3; the structure was reported for the radiolabeled substance noting the location of the radiolabel within the substance.
	Metric 2:	High	The non-radiolabeled, 1,1-dichloroethane was sourced from LGC standards; the radiolabeled 1,1-dichloroethane was sourced from Moravek Biochemical Inc. The study included certificates of analyses that included lot/batch numbers.
	Metric 3:	High	The radiochemical purity of the radiolabeled test substance was reported to be $\geq 98\%$ by GC-MS. The purity of the non-radiolabeled substance was 99.48% (batch 1) and 99.9% (batch 2).

Continued on next page ...

... continued from previous page

Study Citation:		Labcorp Early Development, (2024). 1,1-Dichloroethane: Rates of penetration through human skin using a flow through in vitro system.		
HERO ID:		11396332		
		EVALUATION		
Domain	Metric	Rating	Comments	
Domain 2: Test Design	Metric 4:	Reference compounds	Medium	Testosterone (non-labelled and radiolabeled) was used as a reference compound in accordance with OECD 28 guidelines, although the reference tests were only conducted using finite exposure conditions. 1 mg/ml was administered to the skin (0.011 mg/cm ²). Study authors report 92.1% recovery (CV = 2.57%). The CV values for some measurements were <25% (total dislodgeable dose, total unabsorbed dose, and the amount in skin washes), while others were >25% (i.e., total absorbed dose, dose in the skin, dose on surface tape and donor chambers etc.). It is unclear if these high CV values suggest any issues with the reliability of the tests conducted. However, this lab also cites previous publications that demonstrate the labs proficiency in performing dermal absorption studies with lipophilic compounds (Van de Sandt et al. 2004).
	Metric 5:	Assay procedures	Medium	This study was conducted according to OECD TG 428, OECD 28, and OECD 156. The assay procedures specified in the report were described in detail, although some information was missing. The flow through diffusion setup was sufficiently reported including a schematic drawing. The study did not report if there was an equilibration period after the skin was placed into the chamber (guidelines recommend a 30-minute equilibrium period). An infinite dose of 1,1-dichloroethane (neat, 50%, or 10% dissolved in either isopropyl myristate or 1,2-dichloroethane) was applied to human skin (at least 6 replicates; surface area of 0.64 cm ²). The volume applied was 64 – 640 uL per 0.64 cm ² skin area depending on the dilution. Higher volumes were necessary to maintain infinite dosing. Human skin samples were dermatomed; samples contained epidermis and some dermis. The thickness of the skin samples ranged from 200 to 400 um, exact thicknesses were not reported. The skin was exposed to a test substance under occluded conditions (PTFE screw lid) for 24 hours. The receptor solution was water fortified with 6% polyoxyethylene 20 oleyl ether. OECD 156 guidelines recommend <6% polyoxyethelene (20) oleyl ether in water for lipophilic compounds. The solubility of 1,1-dichloroethane was tested in the receptor fluid prior to the study and it was determined that dissolution of 1,1-dichloroethane was not rate-limiting. The skin membranes were maintained at 32 degrees C; humidity ranged from 30-70%. The flow rate of receptor fluid was 1.5 mL/hour. Receptor fluid samples (volume not reported) were collected at 10 minutes, 30 minutes, 1 hour, 2 hours, and every 2 hours henceforth for 24 hours (16 samples) post-application and analyzed for radioactivity. "Radioactivity in gross amounts less than twice the background level was considered to be below the limit of accurate determination." A steady state was reached. Details of tape stripping were provided.

Continued on next page ...

... continued from previous page

Study Citation:		Labcorp Early Development, (2024). 1,1-Dichloroethane: Rates of penetration through human skin using a flow through in vitro system.		
HERO ID:		11396332		
		EVALUATION		
Domain	Metric	Rating	Comments	
	Metric 6:	Standards for tests	Medium	The skin samples (abdomen) were obtained from female and male human donors following elective surgery. Samples were stored at -20oC until used. The integrity of the skin was determined by stable trans epidermal water loss (TEWL) prior to dose application and at the end of the experiment. A TEWL of ≤ 13 gm-2h-1 was considered acceptable. This is a slight deviation from guidelines which suggest viable skin has a TEWL reading of less than 10 grams/m2/hour. Skin that did not meet the criteria was allowed to dry and the TEWL was remeasured. Skins that failed to meet the criteria were not included in the analysis. TEWL readings were reported. The percentage of recovered test substances was not reported. However, recovery determination is not generally relevant for studies only determining a Kp. Coefficients of variation (CV) values were not reported but could be determined based on the SD relative to the mean. CVs were all > 25%. Further discussion of the CVs is provided in Metric 19.
Domain 3: Exposure Characterization				
	Metric 7:	Preparation and storage of test substance (chemical)	Medium	The preparation of the test substance was adequately reported. Dilutions were magnetically stirred if deemed necessary by study authors (criteria not reported). The activity and homogeneity of the diluted doses were assessed from the top, middle, and bottom of the solution while being stirred. All doses were considered homogenous (data not shown). Storage conditions were not reported; it is unclear how far in advanced solutions were made, and how they were stored. However, the study included only a single application, and dosing was confirmed by analysis of QC samples taken during the dosing of the cells, therefore, the lack of storage information is not likely to have an impact on the study results. The solubility of the test substance in the receptor fluid was tested, (895 ug/mL) and was appropriate (>15-fold higher than observed absorption); solubility was considered non rate-limiting.
	Metric 8:	Consistency of exposure administration	Low	The study did not use the same volume across all samples (ranging from 64ul – 640ul); however, this was done to maintain an infinite dose. The skin surface area of 0.64 cm2 was consistent across groups. However, the skin thickness was reported as a range (200 to 400 uM). It is unclear if the variation in thicknesses was consistent across groups, and this may have contributed to some of the endpoint variations (and subsequently high CVs) observed.
	Metric 9:	Reporting of concentrations	High	The applied dose is reported as dpm, mg per cell, and mg/cm2. Individual cells are reported independently. Nominal and analytical doses are reported.
	Metric 10:	Exposure frequency	High	Exposure duration was reported and appropriate for Kp determination, a 24-hour exposure was used and a steady state was reached.
	Metric 11:	Number of exposure groups and concentration spacing	High	There were 3 dose groups tested in a wide range of concentrations (neat to 10%). Dilutions of the test substance were performed in two different vehicles (isopropyl myristate or 1,2-dichloroethane). The study authors state “The dose levels selected in this study were selected to be representative of potential in-life exposure levels”.
Domain 4: Test Model				
Continued on next page ...				

... continued from previous page

Study Citation:		Labcorp Early Development, (2024). 1,1-Dichloroethane: Rates of penetration through human skin using a flow through in vitro system.		
HERO ID:		11396332		
		EVALUATION		
Domain	Metric	Rating	Comments	
	Metric 12:	Test model (skin)	High	The test model and descriptive information were reported. Samples of full thickness skin samples were obtained from males and females following elective surgery. Abdominal skin samples from 27 people (race not reported) ranging in age from 30-70 years old were studied. The samples were stored frozen at -20 degrees C +/-10 degrees C. Prior to use, the skin samples were thawed, wiped to remove residual fat and blood, re-hydrated in purified water, and dermatomed using a mini-dermatome. The thickness ranged from 200 to 400 uM. The skin contained epidermis and some dermis. The OECD guidelines split thickness (dermatomed) skin is preferred. Membrane integrity was determined by measuring trans epidermal water loss prior to/and upon completion of the experiment.
	Metric 13:	Number/Replicates per group	Medium	The number of replicates was appropriate as per OECD 428. The study used 6 replicates in order to ensure a minimum of 4 intact replicates could be obtained if skin integrity was not maintained.
Domain 5: Outcome Assessment				
	Metric 14:	Outcome assessment methodology	High	The outcome assessment methodology addressed the intended outcome of interest and was sensitive for the outcome. The test followed OECD guidelines 428 and 28. Measurement techniques and timing were reported and appropriate. An infinite dose of the test substance was used to determine the Kp and maximum absorption rate. A dose volume of >100 uL/cm2 was used.
	Metric 15:	Consistency of outcome assessment	High	Details of outcome assessment protocol were reported and outcomes were assessed consistently across study replicates. The same duration of exposure, receptor fluid used, and sampling period were consistent across replicates.
	Metric 16:	Sampling adequacy and sensitivity	Medium	The study reported adequate sampling for the outcomes of interest; measurement sensitivity was sufficient. The sampling intervals were adequate to allow for a steady-state portion of the absorption profile to be obtained. Methods for the determination of radioactivity are reported. "Radioactivity in gross amounts less than twice the background level was considered to be below the limit of accurate determination." Scintillation counts were not shown. A graphical representation of absorption over time is shown.
Domain 6: Confounding/Variable Control				
	Metric 17:	Confounding variables in test design and procedures	Medium	Two batches of radiolabeled and non-radiolabeled 1,1-dichloroethane were used; it is unclear if batches were combined or if different batches were used for different experiments. Abdominal skin was obtained from predominantly females (24 females and 2 males) ranging in age from 30-70 years old. The large age range and sex differences may influence results. Skin integrity was confirmed by TEWL and only skin meeting inclusion criteria (≤ 13 gm-2h-1) was included in the analysis. The sample sizes were sufficient even with the exclusion of any outliers.
	Metric 18:	Confounding variables in outcomes unrelated to exposure	High	There were no reported differences among the study replicates that were unrelated to exposure; the test substance was demonstrated to be soluble in the receptor fluid.
Domain 7: Data Presentation and Analysis				

Continued on next page ...

... continued from previous page

Study Citation: Labcorp Early Development, (2024). 1,1-Dichloroethane: Rates of penetration through human skin using a flow through in vitro system.
HERO ID: 11396332

EVALUATION

Domain	Metric	Rating	Comments
	Metric 19: Data analysis	Uninformative	Means and standard deviations were calculated and reported. Mathematical calculations used to determine Kp were reported and were appropriate. The Kp was based on the steady-state part of the absorption curve. The CVs for Kp and maximum absorption rates were >25% for all dose groups/conditions. The high CV values make the study uninformative.
	Metric 20: Data interpretation	High	The Kp and mean maximum absorption rates were derived from an appropriate exposure condition (infinite dose). Recovery was not reported but this determination is not relevant for infinite dose applications.
	Metric 21: Reporting of data	High	Data for all relevant endpoints were reported quantitatively as means ± SD. Individual cell data are shown.

Overall Quality Determination

Uninformative

Study Citation: Labcorp Early Development, (2024). 1,1-Dichloroethane: Rates of penetration through human skin using a flow through in vitro system.
HERO ID: 11396332

EXTRACTION

Parameter	Data
Extraction ID; Chemical:	Infinite; 10% in IPM; 1,1-Dichloroethane-Parent compound
Skin Material/Species; Skin Preparation; Skin Thickness (um); Diffusion Cell Exposure Setup Type:	ex vivo human; Split thickness; 400; Flow-through; Notes: Not Reported
Occlusion Type; Donor Chamber Vehicle; Concentration of Test Substance in Vehicle (enter as percent):	Occluded; Isopropyl myristate; 10
Mass per Surface Area on Skin (mg/cm2); Duration of Test Substance on Skin:	109; 24 hrs; Not Reported
Duration of Absorbance Measured; Frequency of Samples:	24 hrs; Samples were collected at 10 minutes, 30 minutes, 1 hour, 2 hours, and then every 2 hours; Notes: Not Reported
Time Skin was Washed and Method used; Radiolabel Presence:	Skin not washed; Yes
Total Recovery (percent); Dose Type:	Not Reported; Infinite
Percent Found in Skin Depot After Washing and Tape Stripping; Comments:	Not Reported; Notes: Not Reported
Percent Found in All Tape Strips, Excluding the Upper Two Strips; Comments:	Not Reported; Notes: Not Reported
Percent Found in Receptor Fluid and Receptor Fluid Rinse; Comments:	Not Reported; Notes: Not Reported
Total Percent Absorbed:	0
Steady State Permeability Coefficient (Kp) (cm/hr); Steady State Permeability Coefficient (Comments); Steady State Flux (ug/cm2/hr); Steady State Flux (Comments); Maximum Permeability Coefficient (Kp) (cm/hr); Maximum Permeability Coefficient (Comments); Maximum Flux (ug/cm2/hr); Maximum Flux (Comments):	0.0003; Notes: CV = 46.7%; Not Reported; Notes: Not Reported; Not Reported; Notes: Not Reported; 50.1; Notes: CV = 61.9%

EVALUATION

Domain	Metric	Rating	Comments
Domain 1: Test Substance			
	Metric 1:	High	Test substances were identified as non-radiolabeled 1,1-dichloroethane and radiolabeled 1,1-14C dichloroethane, CAS number: 75-34-3; the structure was reported for the radiolabeled substance noting the location of the radiolabel within the substance.
	Metric 2:	High	The non-radiolabeled, 1,1-dichloroethane was sourced from LGC standards; the radiolabeled 1,1-dichloroethane was sourced from Moravex Biochemical Inc. The study included certificates of analyses that included lot/batch numbers.
	Metric 3:	High	The radiochemical purity of the radiolabeled test substance was reported to be ≥98% by GC-MS. The purity of the non-radiolabeled substance was 99.48% (batch 1) and 99.9% (batch 2).

Domain 2: Test Design

Continued on next page ...

... continued from previous page

Study Citation:		Labcorp Early Development, (2024). 1,1-Dichloroethane: Rates of penetration through human skin using a flow through in vitro system.		
HERO ID:		11396332		
		EVALUATION		
Domain	Metric	Rating	Comments	
	Metric 4:	Reference compounds	Medium	Testosterone (non-labelled and radiolabeled) was used as a reference compound in accordance with OECD 28 guidelines, although the reference tests were only conducted using finite exposure conditions. 1 mg/ml was administered to the skin (0.011 mg/cm ²). Study authors report 92.1% recovery (CV = 2.57%). The CV values for some measurements were <25% (total dislodgeable dose, total unabsorbed dose, and the amount in skin washes), while others were >25% (i.e., total absorbed dose, dose in the skin, dose on surface tape and donor chambers etc.). It is unclear if these high CV values suggest any issues with the reliability of the tests conducted. However, this lab also cites previous publications that demonstrate the labs proficiency in performing dermal absorption studies with lipophilic compounds (Van de Sandt et al. 2004).
	Metric 5:	Assay procedures	Medium	This study was conducted according to OECD TG 428, OECD 28, and OECD 156. The assay procedures specified in the report were described in detail, although some information was missing. The flow through diffusion setup was sufficiently reported including a schematic drawing. The study did not report if there was an equilibration period after the skin was placed into the chamber (guidelines recommend a 30-minute equilibrium period). An infinite dose of 1,1-dichloroethane (neat, 50%, or 10% dissolved in either isopropyl myristate or 1,2-dichloroethane) was applied to human skin (at least 6 replicates; surface area of 0.64 cm ²). The volume applied was 64 – 640 uL per 0.64 cm ² skin area depending on the dilution. Higher volumes were necessary to maintain infinite dosing. Human skin samples were dermatomed; samples contained epidermis and some dermis. The thickness of the skin samples ranged from 200 to 400 um, exact thicknesses were not reported. The skin was exposed to a test substance under occluded conditions (PTFE screw lid) for 24 hours. The receptor solution was water fortified with 6% polyoxyethylene 20 oleyl ether. OECD 156 guidelines recommend <6% polyoxyethelene (20) oleyl ether in water for lipophilic compounds. The solubility of 1,1-dichloroethane was tested in the receptor fluid prior to the study and it was determined that dissolution of 1,1-dichloroethane was not rate-limiting. The skin membranes were maintained at 32 degrees C; humidity ranged from 30-70%. The flow rate of receptor fluid was 1.5 mL/hour. Receptor fluid samples (volume not reported) were collected at 10 minutes, 30 minutes, 1 hour, 2 hours, and every 2 hours henceforth for 24 hours (16 samples) post-application and analyzed for radioactivity. "Radioactivity in gross amounts less than twice the background level was considered to be below the limit of accurate determination." A steady state was reached. Details of tape stripping were provided.
	Metric 6:	Standards for tests	Medium	The skin samples (abdomen) were obtained from female and male human donors following elective surgery. Samples were stored at -20oC until used. The integrity of the skin was determined by stable trans epidermal water loss (TEWL) prior to dose application and at the end of the experiment. A TEWL of ≤ 13 gm-2h-1 was considered acceptable. This is a slight deviation from guidelines which suggest viable skin has a TEWL reading of less than 10 grams/m ² /hour. Skin that did not meet the criteria was allowed to dry and the TEWL was remeasured. Skins that failed to meet the criteria were not included in the analysis. TEWL readings were reported. The percentage of recovered test substances was not reported. However, recovery determination is not generally relevant for studies only determining a Kp. Coefficients of variation (CV) values were not reported but could be determined based on the SD relative to the mean. CVs were all > 25%. Further discussion of the CVs is provided in Metric 19.
Domain 3: Exposure Characterization				

Continued on next page ...

... continued from previous page

Study Citation:		Labcorp Early Development, (2024). 1,1-Dichloroethane: Rates of penetration through human skin using a flow through in vitro system.		
HERO ID:		11396332		
		EVALUATION		
Domain	Metric	Rating	Comments	
	Metric 7:	Preparation and storage of test substance (chemical)	Medium	The preparation of the test substance was adequately reported. Dilutions were magnetically stirred if deemed necessary by study authors (criteria not reported). The activity and homogeneity of the diluted doses were assessed from the top, middle, and bottom of the solution while being stirred. All doses were considered homogenous (data not shown). Storage conditions were not reported; it is unclear how far in advanced solutions were made, and how they were stored. However, the study included only a single application, and dosing was confirmed by analysis of QC samples taken during the dosing of the cells, therefore, the lack of storage information is not likely to have an impact on the study results. The solubility of the test substance in the receptor fluid was tested, (895 ug/mL) and was appropriate (>15-fold higher than observed absorption); solubility was considered non rate-limiting.
	Metric 8:	Consistency of exposure administration	Low	The study did not use the same volume across all samples (ranging from 64ul – 640ul); however, this was done to maintain an infinite dose. The skin surface area of 0.64 cm ² was consistent across groups. However, the skin thickness was reported as a range (200 to 400 uM). It is unclear if the variation in thicknesses was consistent across groups, and this may have contributed to some of the endpoint variations (and subsequently high CVs) observed.
	Metric 9:	Reporting of concentrations	High	The applied dose is reported as dpm, mg per cell, and mg/cm ² . Individual cells are reported independently. Nominal and analytical doses are reported.
	Metric 10:	Exposure frequency	High	Exposure duration was reported and appropriate for K _p determination, a 24-hour exposure was used and a steady state was reached.
	Metric 11:	Number of exposure groups and concentration spacing	High	There were 3 dose groups tested in a wide range of concentrations (neat to 10%). Dilutions of the test substance were performed in two different vehicles (isopropyl myristate or 1,2-dichloroethane). The study authors state “The dose levels selected in this study were selected to be representative of potential in-life exposure levels”.
Domain 4: Test Model				
	Metric 12:	Test model (skin)	High	The test model and descriptive information were reported. Samples of full thickness skin samples were obtained from males and females following elective surgery. Abdominal skin samples from 27 people (race not reported) ranging in age from 30-70 years old were studied. The samples were stored frozen at -20 degrees C +/-10 degrees C. Prior to use, the skin samples were thawed, wiped to remove residual fat and blood, re-hydrated in purified water, and dermatomed using a mini-dermatome. The thickness ranged from 200 to 400 uM. The skin contained epidermis and some dermis. The OECD guidelines split thickness (dermatomed) skin is preferred. Membrane integrity was determined by measuring trans epidermal water loss prior to/and upon completion of the experiment.
	Metric 13:	Number/Replicates per group	Medium	The number of replicates was appropriate as per OECD 428. The study used 6 replicates in order to ensure a minimum of 4 intact replicates could be obtained if skin integrity was not maintained.
Domain 5: Outcome Assessment				
	Metric 14:	Outcome assessment methodology	High	The outcome assessment methodology addressed the intended outcome of interest and was sensitive for the outcome. The test followed OECD guidelines 428 and 28. Measurement techniques and timing were reported and appropriate. An infinite dose of the test substance was used to determine the K _p and maximum absorption rate. A dose volume of >100 uL/cm ² was used.

Continued on next page ...

... continued from previous page

Study Citation:		Labcorp Early Development, (2024). 1,1-Dichloroethane: Rates of penetration through human skin using a flow through in vitro system.		
HERO ID:		11396332		
		EVALUATION		
Domain	Metric	Rating	Comments	
	Metric 15:	Consistency of outcome assessment	High	Details of outcome assessment protocol were reported and outcomes were assessed consistently across study replicates. The same duration of exposure, receptor fluid used, and sampling period were consistent across replicates.
	Metric 16:	Sampling adequacy and sensitivity	Medium	The study reported adequate sampling for the outcomes of interest; measurement sensitivity was sufficient. The sampling intervals were adequate to allow for a steady-state portion of the absorption profile to be obtained. Methods for the determination of radioactivity are reported. "Radioactivity in gross amounts less than twice the background level was considered to be below the limit of accurate determination." Scintillation counts were not shown. A graphical representation of absorption over time is shown.
Domain 6: Confounding/Variable Control				
	Metric 17:	Confounding variables in test design and procedures	Medium	Two batches of radiolabeled and non-radiolabeled 1,1-dichloroethane were used; it is unclear if batches were combined or if different batches were used for different experiments. Abdominal skin was obtained from predominantly females (24 females and 2 males) ranging in age from 30-70 years old. The large age range and sex differences may influence results. Skin integrity was confirmed by TEWL and only skin meeting inclusion criteria ($\leq 13 \text{ gm-2h-1}$) was included in the analysis. The sample sizes were sufficient even with the exclusion of any outliers.
	Metric 18:	Confounding variables in outcomes unrelated to exposure	High	There were no reported differences among the study replicates that were unrelated to exposure; the test substance was demonstrated to be soluble in the receptor fluid.
Domain 7: Data Presentation and Analysis				
	Metric 19:	Data analysis	Uninformative	Means and standard deviations were calculated and reported. Mathematical calculations used to determine Kp were reported and were appropriate. The Kp was based on the steady-state part of the absorption curve. The CVs for Kp and maximum absorption rates were $>25\%$ for all dose groups/conditions. The high CV values make the study uninformative.
	Metric 20:	Data interpretation	High	The Kp and mean maximum absorption rates were derived from an appropriate exposure condition (infinite dose). Recovery was not reported but this determination is not relevant for infinite dose applications.
	Metric 21:	Reporting of data	High	Data for all relevant endpoints were reported quantitatively as means \pm SD. Individual cell data are shown.

Overall Quality Determination

Uninformative

Study Citation: Labcorp Early Development, (2024). 1,1-Dichloroethane: Rates of penetration through human skin using a flow through in vitro system.
HERO ID: 11396332

EXTRACTION

Parameter	Data
Extraction ID; Chemical:	Infinite; neat; 1,1-Dichloroethane-Parent compound
Skin Material/Species; Skin Preparation; Skin Thickness (um); Diffusion Cell Exposure Setup Type:	ex vivo human; Split thickness; 400; Flow-through; Notes: Not Reported
Occlusion Type; Donor Chamber Vehicle; Concentration of Test Substance in Vehicle (enter as percent):	Occluded; Not Reported; 100
Mass per Surface Area on Skin (mg/cm2); Duration of Test Substance on Skin:	1177; 24 hrs; Not Reported
Duration of Absorbance Measured; Frequency of Samples:	24 hrs; Samples were collected at 10 minutes, 30 minutes, 1 hour, 2 hours, and then every 2 hours; Notes: Not Reported
Time Skin was Washed and Method used; Radiolabel Presence:	Skin not washed; Yes
Total Recovery (percent); Dose Type:	Not Reported; Infinite
Percent Found in Skin Depot After Washing and Tape Stripping; Comments:	Not Reported; Notes: Not Reported
Percent Found in All Tape Strips, Excluding the Upper Two Strips; Comments:	Not Reported; Notes: Not Reported
Percent Found in Receptor Fluid and Receptor Fluid Rinse; Comments:	Not Reported; Notes: Not Reported
Total Percent Absorbed:	0
Steady State Permeability Coefficient (Kp) (cm/hr); Steady State Permeability Coefficient (Comments); Steady State Flux (ug/cm2/hr); Steady State Flux (Comments); Maximum Permeability Coefficient (Kp) (cm/hr); Maximum Permeability Coefficient (Comments); Maximum Flux (ug/cm2/hr); Maximum Flux (Comments):	0.00229; Notes: CV=77.7%; Not Reported; Notes: Not Reported; Not Reported; Notes: Not Reported; 3344; Notes: CV = 67.8%

EVALUATION

Domain	Metric	Rating	Comments
Domain 1: Test Substance			
	Metric 1:	High	Test substances were identified as non-radiolabeled 1,1-dichloroethane and radiolabeled 1,1-14C dichloroethane, CAS number: 75-34-3; the structure was reported for the radiolabeled substance noting the location of the radiolabel within the substance.
	Metric 2:	High	The non-radiolabeled, 1,1-dichloroethane was sourced from LGC standards; the radiolabeled 1,1-dichloroethane was sourced from Moravex Biochemical Inc. The study included certificates of analyses that included lot/batch numbers.
	Metric 3:	High	The radiochemical purity of the radiolabeled test substance was reported to be ≥98% by GC-MS. The purity of the non-radiolabeled substance was 99.48% (batch 1) and 99.9% (batch 2).

Domain 2: Test Design

Continued on next page ...

... continued from previous page

Study Citation:		Labcorp Early Development, (2024). 1,1-Dichloroethane: Rates of penetration through human skin using a flow through in vitro system.		
HERO ID:		11396332		
		EVALUATION		
Domain	Metric	Rating	Comments	
	Metric 4:	Reference compounds	Medium	Testosterone (non-labelled and radiolabeled) was used as a reference compound in accordance with OECD 28 guidelines, although the reference tests were only conducted using finite exposure conditions. 1 mg/ml was administered to the skin (0.011 mg/cm ²). Study authors report 92.1% recovery (CV = 2.57%). The CV values for some measurements were <25% (total dislodgeable dose, total unabsorbed dose, and the amount in skin washes), while others were >25% (i.e., total absorbed dose, dose in the skin, dose on surface tape and donor chambers etc.). It is unclear if these high CV values suggest any issues with the reliability of the tests conducted. However, this lab also cites previous publications that demonstrate the labs proficiency in performing dermal absorption studies with lipophilic compounds (Van de Sandt et al. 2004).
	Metric 5:	Assay procedures	Medium	This study was conducted according to OECD TG 428, OECD 28, and OECD 156. The assay procedures specified in the report were described in detail, although some information was missing. The flow through diffusion setup was sufficiently reported including a schematic drawing. The study did not report if there was an equilibration period after the skin was placed into the chamber (guidelines recommend a 30-minute equilibrium period). An infinite dose of 1,1-dichloroethane (neat, 50%, or 10% dissolved in either isopropyl myristate or 1,2-dichloroethane) was applied to human skin (at least 6 replicates; surface area of 0.64 cm ²). The volume applied was 64 – 640 uL per 0.64 cm ² skin area depending on the dilution. Higher volumes were necessary to maintain infinite dosing. Human skin samples were dermatomed; samples contained epidermis and some dermis. The thickness of the skin samples ranged from 200 to 400 um, exact thicknesses were not reported. The skin was exposed to a test substance under occluded conditions (PTFE screw lid) for 24 hours. The receptor solution was water fortified with 6% polyoxyethylene 20 oleyl ether. OECD 156 guidelines recommend <6% polyoxyethelene (20) oleyl ether in water for lipophilic compounds. The solubility of 1,1-dichloroethane was tested in the receptor fluid prior to the study and it was determined that dissolution of 1,1-dichloroethane was not rate-limiting. The skin membranes were maintained at 32 degrees C; humidity ranged from 30-70%. The flow rate of receptor fluid was 1.5 mL/hour. Receptor fluid samples (volume not reported) were collected at 10 minutes, 30 minutes, 1 hour, 2 hours, and every 2 hours henceforth for 24 hours (16 samples) post-application and analyzed for radioactivity. “Radioactivity in gross amounts less than twice the background level was considered to be below the limit of accurate determination.” A steady state was reached. Details of tape stripping were provided.
	Metric 6:	Standards for tests	Medium	The skin samples (abdomen) were obtained from female and male human donors following elective surgery. Samples were stored at -20oC until used. The integrity of the skin was determined by stable trans epidermal water loss (TEWL) prior to dose application and at the end of the experiment. A TEWL of ≤ 13 gm-2h-1 was considered acceptable. This is a slight deviation from guidelines which suggest viable skin has a TEWL reading of less than 10 grams/m ² /hour. Skin that did not meet the criteria was allowed to dry and the TEWL was remeasured. Skins that failed to meet the criteria were not included in the analysis. TEWL readings were reported. The percentage of recovered test substances was not reported. However, recovery determination is not generally relevant for studies only determining a Kp. Coefficients of variation (CV) values were not reported but could be determined based on the SD relative to the mean. CVs were all > 25%. Further discussion of the CVs is provided in Metric 19.

Domain 3: Exposure Characterization

Continued on next page ...

... continued from previous page

Study Citation:		Labcorp Early Development, (2024). 1,1-Dichloroethane: Rates of penetration through human skin using a flow through in vitro system.		
HERO ID:		11396332		
		EVALUATION		
Domain	Metric	Rating	Comments	
	Metric 7:	Preparation and storage of test substance (chemical)	Medium	The preparation of the test substance was adequately reported. Dilutions were magnetically stirred if deemed necessary by study authors (criteria not reported). The activity and homogeneity of the diluted doses were assessed from the top, middle, and bottom of the solution while being stirred. All doses were considered homogenous (data not shown). Storage conditions were not reported; it is unclear how far in advanced solutions were made, and how they were stored. However, the study included only a single application, and dosing was confirmed by analysis of QC samples taken during the dosing of the cells, therefore, the lack of storage information is not likely to have an impact on the study results. The solubility of the test substance in the receptor fluid was tested, (895 ug/mL) and was appropriate (>15-fold higher than observed absorption); solubility was considered non rate-limiting.
	Metric 8:	Consistency of exposure administration	Low	The study did not use the same volume across all samples (ranging from 64ul – 640ul); however, this was done to maintain an infinite dose. The skin surface area of 0.64 cm ² was consistent across groups. However, the skin thickness was reported as a range (200 to 400 uM). It is unclear if the variation in thicknesses was consistent across groups, and this may have contributed to some of the endpoint variations (and subsequently high CVs) observed.
	Metric 9:	Reporting of concentrations	High	The applied dose is reported as dpm, mg per cell, and mg/cm ² . Individual cells are reported independently. Nominal and analytical doses are reported.
	Metric 10:	Exposure frequency	High	Exposure duration was reported and appropriate for K _p determination, a 24-hour exposure was used and a steady state was reached.
	Metric 11:	Number of exposure groups and concentration spacing	High	There were 3 dose groups tested in a wide range of concentrations (neat to 10%). Dilutions of the test substance were performed in two different vehicles (isopropyl myristate or 1,2-dichloroethane). The study authors state “The dose levels selected in this study were selected to be representative of potential in-life exposure levels”.
Domain 4: Test Model				
	Metric 12:	Test model (skin)	High	The test model and descriptive information were reported. Samples of full thickness skin samples were obtained from males and females following elective surgery. Abdominal skin samples from 27 people (race not reported) ranging in age from 30-70 years old were studied. The samples were stored frozen at -20 degrees C +/-10 degrees C. Prior to use, the skin samples were thawed, wiped to remove residual fat and blood, re-hydrated in purified water, and dermatomed using a mini-dermatome. The thickness ranged from 200 to 400 uM. The skin contained epidermis and some dermis. The OECD guidelines split thickness (dermatomed) skin is preferred. Membrane integrity was determined by measuring trans epidermal water loss prior to/and upon completion of the experiment.
	Metric 13:	Number/Replicates per group	Medium	The number of replicates was appropriate as per OECD 428. The study used 6 replicates in order to ensure a minimum of 4 intact replicates could be obtained if skin integrity was not maintained.
Domain 5: Outcome Assessment				
	Metric 14:	Outcome assessment methodology	High	The outcome assessment methodology addressed the intended outcome of interest and was sensitive for the outcome. The test followed OECD guidelines 428 and 28. Measurement techniques and timing were reported and appropriate. An infinite dose of the test substance was used to determine the K _p and maximum absorption rate. A dose volume of >100 uL/cm ² was used.

Continued on next page ...

... continued from previous page

Study Citation:		Labcorp Early Development, (2024). 1,1-Dichloroethane: Rates of penetration through human skin using a flow through in vitro system.		
HERO ID:		11396332		
		EVALUATION		
Domain	Metric	Rating	Comments	
	Metric 15:	Consistency of outcome assessment	High	Details of outcome assessment protocol were reported and outcomes were assessed consistently across study replicates. The same duration of exposure, receptor fluid used, and sampling period were consistent across replicates.
	Metric 16:	Sampling adequacy and sensitivity	Medium	The study reported adequate sampling for the outcomes of interest; measurement sensitivity was sufficient. The sampling intervals were adequate to allow for a steady-state portion of the absorption profile to be obtained. Methods for the determination of radioactivity are reported. "Radioactivity in gross amounts less than twice the background level was considered to be below the limit of accurate determination." Scintillation counts were not shown. A graphical representation of absorption over time is shown.
Domain 6: Confounding/Variable Control				
	Metric 17:	Confounding variables in test design and procedures	Medium	Two batches of radiolabeled and non-radiolabeled 1,1-dichloroethane were used; it is unclear if batches were combined or if different batches were used for different experiments. Abdominal skin was obtained from predominantly females (24 females and 2 males) ranging in age from 30-70 years old. The large age range and sex differences may influence results. Skin integrity was confirmed by TEWL and only skin meeting inclusion criteria ($\leq 13 \text{ gm-2h-1}$) was included in the analysis. The sample sizes were sufficient even with the exclusion of any outliers.
	Metric 18:	Confounding variables in outcomes unrelated to exposure	High	There were no reported differences among the study replicates that were unrelated to exposure; the test substance was demonstrated to be soluble in the receptor fluid.
Domain 7: Data Presentation and Analysis				
	Metric 19:	Data analysis	Uninformative	Means and standard deviations were calculated and reported. Mathematical calculations used to determine Kp were reported and were appropriate. The Kp was based on the steady-state part of the absorption curve. The CVs for Kp and maximum absorption rates were $>25\%$ for all dose groups/conditions. The high CV values make the study uninformative.
	Metric 20:	Data interpretation	High	The Kp and mean maximum absorption rates were derived from an appropriate exposure condition (infinite dose). Recovery was not reported but this determination is not relevant for infinite dose applications.
	Metric 21:	Reporting of data	High	Data for all relevant endpoints were reported quantitatively as means \pm SD. Individual cell data are shown.

Overall Quality Determination

Uninformative

Study Citation: Labcorp Early Development, (2024). 1,1-Dichloroethane: Rates of penetration through human skin using a flow through in vitro system.
HERO ID: 11396332

EXTRACTION**Parameter****Data**

Extraction ID; Chemical: Infinite; 50% in 1,2-dichloroethane; 1,1-Dichloroethane-Parent compound
 Skin Material/Species; Skin Preparation; Skin Thickness (um); Diffusion Cell Exposure Setup Type: ex vivo human; Split thickness; 400; Flow-through; Notes: Not Reported
 Occlusion Type; Donor Chamber Vehicle; Concentration of Test Substance in Vehicle (enter as percent): Occluded; 1,2-dichloroethane; 50
 Mass per Surface Area on Skin (mg/cm2); Duration of Test Substance on Skin: 277; 24 hrs; Not Reported
 Duration of Absorbance Measured; Frequency of Samples: 24 hrs; Samples were collected at 10 minutes, 30 minutes, 1 hour, 2 hours, and then every 2 hours; Notes: Not Reported
 Time Skin was Washed and Method used; Radiolabel Presence: Skin not washed; Yes
 Total Recovery (percent); Dose Type: Not Reported; Infinite
 Percent Found in Skin Depot After Washing and Tape Stripping; Comments: Not Reported; Notes: Not Reported
 Percent Found in All Tape Strips, Excluding the Upper Two Strips; Comments: Not Reported; Notes: Not Reported
 Percent Found in Receptor Fluid and Receptor Fluid Rinse; Comments: Not Reported; Notes: Not Reported
 Total Percent Absorbed: 0
 Steady State Permeability Coefficient (Kp) (cm/hr); Steady State Permeability Coefficient (Comments); Steady State Flux (ug/cm2/hr); Steady State Flux (Comments); Maximum Permeability Coefficient (Kp) (cm/hr); Maximum Permeability Coefficient (Comments); Maximum Flux (ug/cm2/hr); Maximum Flux (Comments): 0.00078; Notes: CV = 82.1%; Not Reported; Notes: Not Reported; Not Reported; Notes: Not Reported; 418; Notes: CV = 53.1%

EVALUATION

Domain	Metric	Rating	Comments
Domain 1: Test Substance			
	Metric 1:	High	Test substances were identified as non-radiolabeled 1,1-dichloroethane and radiolabeled 1,1-14C dichloroethane, CAS number: 75-34-3; the structure was reported for the radiolabeled substance noting the location of the radiolabel within the substance.
	Metric 2:	High	The non-radiolabeled, 1,1-dichloroethane was sourced from LGC standards; the radiolabeled 1,1-dichloroethane was sourced from Moravex Biochemical Inc. The study included certificates of analyses that included lot/batch numbers.
	Metric 3:	High	The radiochemical purity of the radiolabeled test substance was reported to be $\geq 98\%$ % by GC-MS. The purity of the non-radiolabeled substance was 99.48% (batch 1) and 99.9% (batch 2).

Domain 2: Test Design

Continued on next page ...

... continued from previous page

Study Citation:		Labcorp Early Development, (2024). 1,1-Dichloroethane: Rates of penetration through human skin using a flow through in vitro system.		
HERO ID:		11396332		
		EVALUATION		
Domain	Metric	Rating	Comments	
	Metric 4:	Reference compounds	Medium	Testosterone (non-labelled and radiolabeled) was used as a reference compound in accordance with OECD 28 guidelines, although the reference tests were only conducted using finite exposure conditions. 1 mg/ml was administered to the skin (0.011 mg/cm ²). Study authors report 92.1% recovery (CV = 2.57%). The CV values for some measurements were <25% (total dislodgeable dose, total unabsorbed dose, and the amount in skin washes), while others were >25% (i.e., total absorbed dose, dose in the skin, dose on surface tape and donor chambers etc.). It is unclear if these high CV values suggest any issues with the reliability of the tests conducted. However, this lab also cites previous publications that demonstrate the labs proficiency in performing dermal absorption studies with lipophilic compounds (Van de Sandt et al. 2004).
	Metric 5:	Assay procedures	Medium	This study was conducted according to OECD TG 428, OECD 28, and OECD 156. The assay procedures specified in the report were described in detail, although some information was missing. The flow through diffusion setup was sufficiently reported including a schematic drawing. The study did not report if there was an equilibration period after the skin was placed into the chamber (guidelines recommend a 30-minute equilibrium period). An infinite dose of 1,1-dichloroethane (neat, 50%, or 10% dissolved in either isopropyl myristate or 1,2-dichloroethane) was applied to human skin (at least 6 replicates; surface area of 0.64 cm ²). The volume applied was 64 – 640 uL per 0.64 cm ² skin area depending on the dilution. Higher volumes were necessary to maintain infinite dosing. Human skin samples were dermatomed; samples contained epidermis and some dermis. The thickness of the skin samples ranged from 200 to 400 um, exact thicknesses were not reported. The skin was exposed to a test substance under occluded conditions (PTFE screw lid) for 24 hours. The receptor solution was water fortified with 6% polyoxyethylene 20 oleyl ether. OECD 156 guidelines recommend <6% polyoxyethelene (20) oleyl ether in water for lipophilic compounds. The solubility of 1,1-dichloroethane was tested in the receptor fluid prior to the study and it was determined that dissolution of 1,1-dichloroethane was not rate-limiting. The skin membranes were maintained at 32 degrees C; humidity ranged from 30-70%. The flow rate of receptor fluid was 1.5 mL/hour. Receptor fluid samples (volume not reported) were collected at 10 minutes, 30 minutes, 1 hour, 2 hours, and every 2 hours henceforth for 24 hours (16 samples) post-application and analyzed for radioactivity. "Radioactivity in gross amounts less than twice the background level was considered to be below the limit of accurate determination." A steady state was reached. Details of tape stripping were provided.
	Metric 6:	Standards for tests	Medium	The skin samples (abdomen) were obtained from female and male human donors following elective surgery. Samples were stored at -20oC until used. The integrity of the skin was determined by stable trans epidermal water loss (TEWL) prior to dose application and at the end of the experiment. A TEWL of ≤ 13 gm-2h-1 was considered acceptable. This is a slight deviation from guidelines which suggest viable skin has a TEWL reading of less than 10 grams/m ² /hour. Skin that did not meet the criteria was allowed to dry and the TEWL was remeasured. Skins that failed to meet the criteria were not included in the analysis. TEWL readings were reported. The percentage of recovered test substances was not reported. However, recovery determination is not generally relevant for studies only determining a Kp. Coefficients of variation (CV) values were not reported but could be determined based on the SD relative to the mean. CVs were all > 25%. Further discussion of the CVs is provided in Metric 19.
Domain 3: Exposure Characterization				

Continued on next page ...

... continued from previous page

Study Citation:		Labcorp Early Development, (2024). 1,1-Dichloroethane: Rates of penetration through human skin using a flow through in vitro system.		
HERO ID:		11396332		
		EVALUATION		
Domain	Metric	Rating	Comments	
	Metric 7:	Preparation and storage of test substance (chemical)	Medium	The preparation of the test substance was adequately reported. Dilutions were magnetically stirred if deemed necessary by study authors (criteria not reported). The activity and homogeneity of the diluted doses were assessed from the top, middle, and bottom of the solution while being stirred. All doses were considered homogenous (data not shown). Storage conditions were not reported; it is unclear how far in advanced solutions were made, and how they were stored. However, the study included only a single application, and dosing was confirmed by analysis of QC samples taken during the dosing of the cells, therefore, the lack of storage information is not likely to have an impact on the study results. The solubility of the test substance in the receptor fluid was tested, (895 ug/mL) and was appropriate (>15-fold higher than observed absorption); solubility was considered non rate-limiting.
	Metric 8:	Consistency of exposure administration	Low	The study did not use the same volume across all samples (ranging from 64ul – 640ul); however, this was done to maintain an infinite dose. The skin surface area of 0.64 cm ² was consistent across groups. However, the skin thickness was reported as a range (200 to 400 uM). It is unclear if the variation in thicknesses was consistent across groups, and this may have contributed to some of the endpoint variations (and subsequently high CVs) observed.
	Metric 9:	Reporting of concentrations	High	The applied dose is reported as dpm, mg per cell, and mg/cm ² . Individual cells are reported independently. Nominal and analytical doses are reported.
	Metric 10:	Exposure frequency	High	Exposure duration was reported and appropriate for K _p determination, a 24-hour exposure was used and a steady state was reached.
	Metric 11:	Number of exposure groups and concentration spacing	High	There were 3 dose groups tested in a wide range of concentrations (neat to 10%). Dilutions of the test substance were performed in two different vehicles (isopropyl myristate or 1,2-dichloroethane). The study authors state “The dose levels selected in this study were selected to be representative of potential in-life exposure levels”.
Domain 4: Test Model				
	Metric 12:	Test model (skin)	High	The test model and descriptive information were reported. Samples of full thickness skin samples were obtained from males and females following elective surgery. Abdominal skin samples from 27 people (race not reported) ranging in age from 30-70 years old were studied. The samples were stored frozen at -20 degrees C +/-10 degrees C. Prior to use, the skin samples were thawed, wiped to remove residual fat and blood, re-hydrated in purified water, and dermatomed using a mini-dermatome. The thickness ranged from 200 to 400 uM. The skin contained epidermis and some dermis. The OECD guidelines split thickness (dermatomed) skin is preferred. Membrane integrity was determined by measuring trans epidermal water loss prior to/and upon completion of the experiment.
	Metric 13:	Number/Replicates per group	Medium	The number of replicates was appropriate as per OECD 428. The study used 6 replicates in order to ensure a minimum of 4 intact replicates could be obtained if skin integrity was not maintained.
Domain 5: Outcome Assessment				
	Metric 14:	Outcome assessment methodology	High	The outcome assessment methodology addressed the intended outcome of interest and was sensitive for the outcome. The test followed OECD guidelines 428 and 28. Measurement techniques and timing were reported and appropriate. An infinite dose of the test substance was used to determine the K _p and maximum absorption rate. A dose volume of >100 uL/cm ² was used.

Continued on next page ...

... continued from previous page

Study Citation:		Labcorp Early Development, (2024). 1,1-Dichloroethane: Rates of penetration through human skin using a flow through in vitro system.		
HERO ID:		11396332		
		EVALUATION		
Domain	Metric	Rating	Comments	
	Metric 15:	Consistency of outcome assessment	High	Details of outcome assessment protocol were reported and outcomes were assessed consistently across study replicates. The same duration of exposure, receptor fluid used, and sampling period were consistent across replicates.
	Metric 16:	Sampling adequacy and sensitivity	Medium	The study reported adequate sampling for the outcomes of interest; measurement sensitivity was sufficient. The sampling intervals were adequate to allow for a steady-state portion of the absorption profile to be obtained. Methods for the determination of radioactivity are reported. "Radioactivity in gross amounts less than twice the background level was considered to be below the limit of accurate determination." Scintillation counts were not shown. A graphical representation of absorption over time is shown.
Domain 6: Confounding/Variable Control				
	Metric 17:	Confounding variables in test design and procedures	Medium	Two batches of radiolabeled and non-radiolabeled 1,1-dichloroethane were used; it is unclear if batches were combined or if different batches were used for different experiments. Abdominal skin was obtained from predominantly females (24 females and 2 males) ranging in age from 30-70 years old. The large age range and sex differences may influence results. Skin integrity was confirmed by TEWL and only skin meeting inclusion criteria (≤ 13 gm-2h-1) was included in the analysis. The sample sizes were sufficient even with the exclusion of any outliers.
	Metric 18:	Confounding variables in outcomes unrelated to exposure	High	There were no reported differences among the study replicates that were unrelated to exposure; the test substance was demonstrated to be soluble in the receptor fluid.
Domain 7: Data Presentation and Analysis				
	Metric 19:	Data analysis	Uninformative	Means and standard deviations were calculated and reported. Mathematical calculations used to determine Kp were reported and were appropriate. The Kp was based on the steady-state part of the absorption curve. The CVs for Kp and maximum absorption rates were $>25\%$ for all dose groups/conditions. The high CV values make the study uninformative.
	Metric 20:	Data interpretation	High	The Kp and mean maximum absorption rates were derived from an appropriate exposure condition (infinite dose). Recovery was not reported but this determination is not relevant for infinite dose applications.
	Metric 21:	Reporting of data	High	Data for all relevant endpoints were reported quantitatively as means \pm SD. Individual cell data are shown.

Overall Quality Determination**Uninformative**

Study Citation: Labcorp Early Development, (2024). 1,1-Dichloroethane: Rates of penetration through human skin using a flow through in vitro system.
HERO ID: 11396332

EXTRACTION**Parameter****Data**

Extraction ID; Chemical: Infinite; 10% in 1,2-dichloroethane; 1,1-Dichloroethane-Parent compound
 Skin Material/Species; Skin Preparation; Skin Thickness (um); Diffusion Cell Exposure Setup Type: ex vivo human; Split thickness; 400; Flow-through; Notes: Not Reported
 Occlusion Type; Donor Chamber Vehicle; Concentration of Test Substance in Vehicle (enter as percent): Occluded; 1,2-dichloroethane; 10
 Mass per Surface Area on Skin (mg/cm²); Duration of Test Substance on Skin: 249; 24 hrs; Not Reported
 Duration of Absorbance Measured; Frequency of Samples: 24 hrs; Samples were collected at 10 minutes, 30 minutes, 1 hour, 2 hours, and then every 2 hours; Notes: Not Reported
 Time Skin was Washed and Method used; Radiolabel Presence: Skin not washed; Yes
 Total Recovery (percent); Dose Type: Not Reported; Infinite
 Percent Found in Skin Depot After Washing and Tape Stripping; Comments: Not Reported; Notes: Not Reported
 Percent Found in All Tape Strips, Excluding the Upper Two Strips; Comments: Not Reported; Notes: Not Reported
 Percent Found in Receptor Fluid and Receptor Fluid Rinse; Comments: Not Reported; Notes: Not Reported
 Total Percent Absorbed: 0
 Steady State Permeability Coefficient (Kp) (cm/hr); Steady State Permeability Coefficient (Comments); Steady State Flux (ug/cm²/hr); Steady State Flux (Comments); Maximum Permeability Coefficient (Kp) (cm/hr); Maximum Permeability Coefficient (Comments); Maximum Flux (ug/cm²/hr); Maximum Flux (Comments): 0.0002; Notes: CV = 60%; Not Reported; Notes: Not Reported; Not Reported; Notes: Not Reported; 77.8; Notes: CV = 64.7%

EVALUATION

Domain	Metric	Rating	Comments
Domain 1: Test Substance			
	Metric 1:	High	Test substances were identified as non-radiolabeled 1,1-dichloroethane and radiolabeled 1,1- ¹⁴ C dichloroethane, CAS number: 75-34-3; the structure was reported for the radiolabeled substance noting the location of the radiolabel within the substance.
	Metric 2:	High	The non-radiolabeled, 1,1-dichloroethane was sourced from LGC standards; the radiolabeled 1,1-dichloroethane was sourced from Moravek Biochemical Inc. The study included certificates of analyses that included lot/batch numbers.
	Metric 3:	High	The radiochemical purity of the radiolabeled test substance was reported to be ≥98% by GC-MS. The purity of the non-radiolabeled substance was 99.48% (batch 1) and 99.9% (batch 2).

Domain 2: Test Design

Continued on next page ...

... continued from previous page

Study Citation:		Labcorp Early Development, (2024). 1,1-Dichloroethane: Rates of penetration through human skin using a flow through in vitro system.		
HERO ID:		11396332		
		EVALUATION		
Domain	Metric	Rating	Comments	
	Metric 4:	Reference compounds	Medium	Testosterone (non-labelled and radiolabeled) was used as a reference compound in accordance with OECD 28 guidelines, although the reference tests were only conducted using finite exposure conditions. 1 mg/ml was administered to the skin (0.011 mg/cm ²). Study authors report 92.1% recovery (CV = 2.57%). The CV values for some measurements were <25% (total dislodgeable dose, total unabsorbed dose, and the amount in skin washes), while others were >25% (i.e., total absorbed dose, dose in the skin, dose on surface tape and donor chambers etc.). It is unclear if these high CV values suggest any issues with the reliability of the tests conducted. However, this lab also cites previous publications that demonstrate the labs proficiency in performing dermal absorption studies with lipophilic compounds (Van de Sandt et al. 2004).
	Metric 5:	Assay procedures	Medium	This study was conducted according to OECD TG 428, OECD 28, and OECD 156. The assay procedures specified in the report were described in detail, although some information was missing. The flow through diffusion setup was sufficiently reported including a schematic drawing. The study did not report if there was an equilibration period after the skin was placed into the chamber (guidelines recommend a 30-minute equilibrium period). An infinite dose of 1,1-dichloroethane (neat, 50%, or 10% dissolved in either isopropyl myristate or 1,2-dichloroethane) was applied to human skin (at least 6 replicates; surface area of 0.64 cm ²). The volume applied was 64 – 640 uL per 0.64 cm ² skin area depending on the dilution. Higher volumes were necessary to maintain infinite dosing. Human skin samples were dermatomed; samples contained epidermis and some dermis. The thickness of the skin samples ranged from 200 to 400 um, exact thicknesses were not reported. The skin was exposed to a test substance under occluded conditions (PTFE screw lid) for 24 hours. The receptor solution was water fortified with 6% polyoxyethylene 20 oleyl ether. OECD 156 guidelines recommend <6% polyoxyethelene (20) oleyl ether in water for lipophilic compounds. The solubility of 1,1-dichloroethane was tested in the receptor fluid prior to the study and it was determined that dissolution of 1,1-dichloroethane was not rate-limiting. The skin membranes were maintained at 32 degrees C; humidity ranged from 30-70%. The flow rate of receptor fluid was 1.5 mL/hour. Receptor fluid samples (volume not reported) were collected at 10 minutes, 30 minutes, 1 hour, 2 hours, and every 2 hours henceforth for 24 hours (16 samples) post-application and analyzed for radioactivity. "Radioactivity in gross amounts less than twice the background level was considered to be below the limit of accurate determination." A steady state was reached. Details of tape stripping were provided.
	Metric 6:	Standards for tests	Medium	The skin samples (abdomen) were obtained from female and male human donors following elective surgery. Samples were stored at -20oC until used. The integrity of the skin was determined by stable trans epidermal water loss (TEWL) prior to dose application and at the end of the experiment. A TEWL of ≤ 13 gm-2h-1 was considered acceptable. This is a slight deviation from guidelines which suggest viable skin has a TEWL reading of less than 10 grams/m ² /hour. Skin that did not meet the criteria was allowed to dry and the TEWL was remeasured. Skins that failed to meet the criteria were not included in the analysis. TEWL readings were reported. The percentage of recovered test substances was not reported. However, recovery determination is not generally relevant for studies only determining a Kp. Coefficients of variation (CV) values were not reported but could be determined based on the SD relative to the mean. CVs were all > 25%. Further discussion of the CVs is provided in Metric 19.
Domain 3: Exposure Characterization				

Continued on next page ...

... continued from previous page

Study Citation:		Labcorp Early Development, (2024). 1,1-Dichloroethane: Rates of penetration through human skin using a flow through in vitro system.		
HERO ID:		11396332		
		EVALUATION		
Domain	Metric	Rating	Comments	
	Metric 7:	Preparation and storage of test substance (chemical)	Medium	The preparation of the test substance was adequately reported. Dilutions were magnetically stirred if deemed necessary by study authors (criteria not reported). The activity and homogeneity of the diluted doses were assessed from the top, middle, and bottom of the solution while being stirred. All doses were considered homogenous (data not shown). Storage conditions were not reported; it is unclear how far in advanced solutions were made, and how they were stored. However, the study included only a single application, and dosing was confirmed by analysis of QC samples taken during the dosing of the cells, therefore, the lack of storage information is not likely to have an impact on the study results. The solubility of the test substance in the receptor fluid was tested, (895 ug/mL) and was appropriate (>15-fold higher than observed absorption); solubility was considered non rate-limiting.
	Metric 8:	Consistency of exposure administration	Low	The study did not use the same volume across all samples (ranging from 64ul – 640ul); however, this was done to maintain an infinite dose. The skin surface area of 0.64 cm ² was consistent across groups. However, the skin thickness was reported as a range (200 to 400 uM). It is unclear if the variation in thicknesses was consistent across groups, and this may have contributed to some of the endpoint variations (and subsequently high CVs) observed.
	Metric 9:	Reporting of concentrations	High	The applied dose is reported as dpm, mg per cell, and mg/cm ² . Individual cells are reported independently. Nominal and analytical doses are reported.
	Metric 10:	Exposure frequency	High	Exposure duration was reported and appropriate for K _p determination, a 24-hour exposure was used and a steady state was reached.
	Metric 11:	Number of exposure groups and concentration spacing	High	There were 3 dose groups tested in a wide range of concentrations (neat to 10%). Dilutions of the test substance were performed in two different vehicles (isopropyl myristate or 1,2-dichloroethane). The study authors state “The dose levels selected in this study were selected to be representative of potential in-life exposure levels”.
Domain 4: Test Model				
	Metric 12:	Test model (skin)	High	The test model and descriptive information were reported. Samples of full thickness skin samples were obtained from males and females following elective surgery. Abdominal skin samples from 27 people (race not reported) ranging in age from 30-70 years old were studied. The samples were stored frozen at -20 degrees C +/-10 degrees C. Prior to use, the skin samples were thawed, wiped to remove residual fat and blood, re-hydrated in purified water, and dermatomed using a mini-dermatome. The thickness ranged from 200 to 400 uM. The skin contained epidermis and some dermis. The OECD guidelines split thickness (dermatomed) skin is preferred. Membrane integrity was determined by measuring trans epidermal water loss prior to/and upon completion of the experiment.
	Metric 13:	Number/Replicates per group	Medium	The number of replicates was appropriate as per OECD 428. The study used 6 replicates in order to ensure a minimum of 4 intact replicates could be obtained if skin integrity was not maintained.
Domain 5: Outcome Assessment				
	Metric 14:	Outcome assessment methodology	High	The outcome assessment methodology addressed the intended outcome of interest and was sensitive for the outcome. The test followed OECD guidelines 428 and 28. Measurement techniques and timing were reported and appropriate. An infinite dose of the test substance was used to determine the K _p and maximum absorption rate. A dose volume of >100 uL/cm ² was used.

Continued on next page ...

... continued from previous page

Study Citation:		Labcorp Early Development, (2024). 1,1-Dichloroethane: Rates of penetration through human skin using a flow through in vitro system.		
HERO ID:		11396332		
		EVALUATION		
Domain	Metric	Rating	Comments	
	Metric 15:	Consistency of outcome assessment	High	Details of outcome assessment protocol were reported and outcomes were assessed consistently across study replicates. The same duration of exposure, receptor fluid used, and sampling period were consistent across replicates.
	Metric 16:	Sampling adequacy and sensitivity	Medium	The study reported adequate sampling for the outcomes of interest; measurement sensitivity was sufficient. The sampling intervals were adequate to allow for a steady-state portion of the absorption profile to be obtained. Methods for the determination of radioactivity are reported. "Radioactivity in gross amounts less than twice the background level was considered to be below the limit of accurate determination." Scintillation counts were not shown. A graphical representation of absorption over time is shown.
Domain 6: Confounding/Variable Control				
	Metric 17:	Confounding variables in test design and procedures	Medium	Two batches of radiolabeled and non-radiolabeled 1,1-dichloroethane were used; it is unclear if batches were combined or if different batches were used for different experiments. Abdominal skin was obtained from predominantly females (24 females and 2 males) ranging in age from 30-70 years old. The large age range and sex differences may influence results. Skin integrity was confirmed by TEWL and only skin meeting inclusion criteria ($\leq 13 \text{ gm-2h-1}$) was included in the analysis. The sample sizes were sufficient even with the exclusion of any outliers.
	Metric 18:	Confounding variables in outcomes unrelated to exposure	High	There were no reported differences among the study replicates that were unrelated to exposure; the test substance was demonstrated to be soluble in the receptor fluid.
Domain 7: Data Presentation and Analysis				
	Metric 19:	Data analysis	Uninformative	Means and standard deviations were calculated and reported. Mathematical calculations used to determine Kp were reported and were appropriate. The Kp was based on the steady-state part of the absorption curve. The CVs for Kp and maximum absorption rates were $>25\%$ for all dose groups/conditions. The high CV values make the study uninformative.
	Metric 20:	Data interpretation	High	The Kp and mean maximum absorption rates were derived from an appropriate exposure condition (infinite dose). Recovery was not reported but this determination is not relevant for infinite dose applications.
	Metric 21:	Reporting of data	High	Data for all relevant endpoints were reported quantitatively as means \pm SD. Individual cell data are shown.

Overall Quality Determination

Uninformative

Study Citation: Labcorp Early Development, (2024). 1,1-Dichloroethane: Rates of penetration through human skin using a flow through in vitro system.
HERO ID: 11396332

EXTRACTION

Parameter	Data
Extraction ID; Chemical:	Finite; 10% in 1,2-dichloroethane; 1,1-Dichloroethane-Parent compound
Skin Material/Species; Skin Preparation; Skin Thickness (um); Diffusion Cell Exposure Setup Type:	ex vivo human; Split thickness; 400; Flow-through; Notes: Not Reported
Occlusion Type; Donor Chamber Vehicle; Concentration of Test Substance in Vehicle (enter as percent):	Semi-occluded (e.g., vapor trap); 1,2-dichloroethane; 10
Mass per Surface Area on Skin (mg/cm ²); Duration of Test Substance on Skin:	0.93; 8 hrs; Not Reported
Duration of Absorbance Measured; Frequency of Samples:	24 hrs; Samples were collected at 10 minutes, 30 minutes, 1 hours, 2 hours, and then 2-hourly intervals; Notes: Not Reported
Time Skin was Washed and Method used; Radiolabel Presence:	"The surface of the membrane was rinsed three times (3 x 0.5 mL) with a mild solution of Dove™ soap in water (ca. 1%). The skin surface was swabbed twice with cotton buds soaked with the liquid soap solution. Finally, the membrane was rinsed with a small volume (0.5 mL) of water and a further cotton bud used to swab the surface of the membrane until dry."; Yes
Total Recovery (percent); Dose Type:	92.76; Finite
Percent Found in Skin Depot After Washing and Tape Stripping; Comments:	0; Notes: not detected
Percent Found in All Tape Strips, Excluding the Upper Two Strips; Comments:	0; Notes: not detected
Percent Found in Receptor Fluid and Receptor Fluid Rinse; Comments:	0.02; Notes: CV= 100%
Total Percent Absorbed:	0
Steady State Permeability Coefficient (Kp) (cm/hr); Steady State Permeability Coefficient (Comments); Steady State Flux (ug/cm ² /hr); Steady State Flux (Comments); Maximum Permeability Coefficient (Kp) (cm/hr); Maximum Permeability Coefficient (Comments); Maximum Flux (ug/cm ² /hr); Maximum Flux (Comments):	Not Reported; Notes: Not Reported; Not Reported; Notes: Not Reported; Not Reported; Notes: Not Reported; Not Reported; Notes: Not Reported

EVALUATION

Domain	Metric	Rating	Comments	
Domain 1: Test Substance				
	Metric 1:	Test substance identity	High	Test substances were identified as non-radiolabeled 1,1-dichloroethane and radiolabeled 1,1- ¹⁴ C dichloroethane, CAS number: 75-34-3; the structure was reported for the radiolabeled substance noting the location of the radiolabel within the substance.
	Metric 2:	Test substance source	High	The non-radiolabeled, 1,1-dichloroethane was sourced from LGC standards; the radiolabeled 1,1-dichloroethane was sourced from Moravek Biochemical Inc. The study included certificates of analyses that included lot/batch numbers.
	Metric 3:	Test substance purity	High	The radiochemical purity of the radiolabeled test substance was reported to be ≥98% by GC-MS. The purity of the non-radiolabeled substance was 99.48% (batch 1) and 99.9% (batch 2).

Continued on next page ...

... continued from previous page

Study Citation:		Labcorp Early Development, (2024). 1,1-Dichloroethane: Rates of penetration through human skin using a flow through in vitro system.		
HERO ID:		11396332		
		EVALUATION		
Domain	Metric	Rating	Comments	
Domain 2: Test Design	Metric 4:	Reference compounds	Medium	Testosterone (non-labelled and radiolabeled) was used as a reference compound in accordance with OECD 28 guidelines. 1 mg/ml was administered to the skin (0.011 mg/cm ²). Study authors report 92.1% recovery (CV = 2.57%). The CV values for some measurements were <25% (total dislodgeable dose, total unabsorbed dose, and the amount in skin washes), while others were >25% (i.e., total absorbed dose, dose in the skin, dose on surface tape and donor chambers etc.). It is unclear if these high CV values suggest any issues with the reliability of the tests conducted. However, this lab also cites previous publications that demonstrate the labs proficiency in performing dermal absorption studies with lipophilic compounds (Van de Sandt et al. 2004).
	Metric 5:	Assay procedures	Medium	This study was conducted according to OECD TG 428, OECD 28, and OECD 156. The assay procedures specified in the report were described in detail, although some information was missing. The flow through diffusion set-up was sufficiently reported including a schematic drawing. The study does not report if there was an equilibration period after the skin was placed into the chamber (guidelines recommend a 30-minute equilibrium period). Finite doses of 1,1-dichloroethane (neat, 50%, 10%, and 1% dissolved in either isopropyl myristate or 1,2-dichloroethane) were applied to human skin (at least 6 replicates; surface area of 0.64 cm ²). The volume applied (10uL/cm ²) applied to the skin surface was reported. The cells were semi-occluded by charcoal filters in the necks of the donor cells. The thickness of the skin samples ranged from 200 to 400 um, the exact thicknesses were not reported. The receptor solution was water fortified with 6% polyoxyethylene 20 oleyl ether. OECD 156 guidelines recommend <6% polyoxyethylene (20) oleyl ether in water for lipophilic compounds. The solubility of 1,1-dichloroethane was tested in the receptor fluid prior to the study and it was determined that the dissolution of 1,1-dichloroethane was not rate-limiting. The skin membranes were maintained at 32 degrees C; humidity ranged from 30-70%. The flow rate of receptor fluid was 1.5 mL/hour. After 8 hours skin was washed. "The surface of the membrane was rinsed three times (3 x 0.5 mL) with a mild solution of Dove™ soap in water (ca. 1%). The skin surface was swabbed twice with cotton buds soaked with the liquid soap solution. Finally, the membrane was rinsed with a small volume (0.5 mL) of water and a further cotton bud was used to swab the surface of the membrane until dry". For some groups, headspace above the skin was drawn into a syringe for analysis after 4 hours. Receptor fluid samples (volume not reported) were collected at 10 minutes, 30 minutes, 1 hour, 2 hours and every 2 hours henceforth for 24 hours (16 samples) and analyzed for radioactivity. After 24 hours, the skin was tape-stripped 5 times. "Radioactivity in gross amounts less than twice the background level was considered to be below the limit of accurate determination."

Continued on next page ...

... continued from previous page

Study Citation:		Labcorp Early Development, (2024). 1,1-Dichloroethane: Rates of penetration through human skin using a flow through in vitro system.		
HERO ID:		11396332		
		EVALUATION		
Domain	Metric	Rating	Comments	
	Metric 6:	Standards for tests	Low	The skin samples (abdomen) were obtained from female and male human donors following elective surgery (24 females and 2 males). Samples were stored at -20oC until used. The integrity of the skin was determined by stable trans epidermal water loss (TEWL) prior to dose application and at termination (prior to skin washing and tape stripping). A TEWL of ≤ 13 gm-2h-1 was considered acceptable. This is a slight deviation from guidelines which suggest viable skin has a TEWL reading of less than 10 grams/m2/hour. TEWL readings were reported. Coefficients of variation (CV) values were reported. CVs were all $> 25\%$. Further discussion of the CVs is provided in Metric 19. The percent recoveries were reported ($<60\%$) and were below the recommended $100 \pm 20\%$.
Domain 3: Exposure Characterization	Metric 7:	Preparation and storage of test substance (chemical)	Medium	The preparation of the test substance was adequately reported. Dilutions were magnetically stirred if deemed necessary by study authors (criteria not reported). The activity and homogeneity of the diluted doses were assessed from the top, middle, and bottom of the solution while being stirred. All doses were considered homogenous (data not shown). Storage conditions were not reported; it is unclear how far in advanced solutions were made, and how they were stored. However, the study included only a single application, and dosing was confirmed by analysis of QC samples taken during the dosing of the cells, therefore, the lack of storage information is not likely to have an impact on the study results. The solubility of the test substance in the receptor fluid was tested, (895 ug/mL) and was appropriate (>15 -fold higher than observed absorption) and was considered non-rate-limiting.
	Metric 8:	Consistency of exposure administration	Low	The volume of test substance applied to the skin was 10 uL/cm2 exposed skin (6.4 uL dose), which is in agreement with OECD guidelines. The skin surface area of 0.64 cm2 was consistent across groups. The skin thickness was reported as a range (200 to 400 uM). It is unclear if the variation in thicknesses was consistent across groups, and this may have contributed to some of the endpoint variations (and subsequently high CVs) observed.
	Metric 9:	Reporting of concentrations	High	The applied dose is reported as dpm, mg per cell, and mg/cm2. Individual cells are reported independently. Nominal and analytical doses are reported.
	Metric 10:	Exposure frequency	High	Exposure duration was reported and appropriate for determining absorption. The test substance was in contact with the skin for 8 hours prior to washing. Samples of receptor fluid were collected for a total of 24 hours.
	Metric 11:	Number of exposure groups and concentration spacing	High	There were 4 dose groups tested in a wide range of concentrations (neat to 1%). Dilutions of the test substance were performed in two different vehicles (isopropyl myristate or 1,2-dichloroethane). The study authors state "The dose levels selected in this study were selected to be representative of potential in-life exposure levels".
Domain 4: Test Model				
Continued on next page ...				

... continued from previous page

Study Citation:		Labcorp Early Development, (2024). 1,1-Dichloroethane: Rates of penetration through human skin using a flow through in vitro system.		
HERO ID:		11396332		
		EVALUATION		
Domain	Metric		Rating	Comments
	Metric 12:	Test model (skin)	High	The test model and descriptive information were reported. Samples of full-thickness skin samples were obtained from males and females following elective surgery. Abdominal skin samples from 26 people (race not reported) ranging in age from 30-70 years old were studied. The samples were stored frozen at -20 degrees C +/-10 degrees C. Prior to use, the skin samples were thawed, wiped to remove residual fat and blood, re-hydrated in purified water, and dermatomed using a mini-dermatome. The thickness ranged from 200 to 400 uM. The skin contained epidermis and some dermis. The OECD guidelines split thickness (dermatomed) skin is preferred. Membrane integrity was determined by measuring transepidermal water loss prior to/and upon completion of the experiment.
	Metric 13:	Number/Replicates per group	Medium	The number of replicates was appropriate as per OECD 428. The study used 6 replicates in order to ensure a minimum of 4 intact replicates could be obtained if skin integrity was not maintained.
Domain 5: Outcome Assessment				
	Metric 14:	Outcome assessment methodology	High	The outcome assessment methodology addressed the intended outcome of interest and was sensitive for the outcome. The test followed OECD guidelines 428 and 28. 10 uL/cm2 of the test substance was applied to skin samples. Measurement techniques and timing were reported and appropriate. A finite dose was used to determine absorption.
	Metric 15:	Consistency of outcome assessment	High	Details of outcome assessment protocol were reported and outcomes were assessed consistently across study replicates. The same duration of exposure, receptor fluid used, and sampling period were consistent across replicates.
	Metric 16:	Sampling adequacy and sensitivity	Medium	The study reported adequate sampling for the outcomes of interest; measurement sensitivity was sufficient. The sampling intervals were adequate (16 over 24 hours). Methods for the determination of radioactivity are reported. "Radioactivity in gross amounts less than twice the background level was considered to be below the limit of accurate determination." Scintillation counts were not shown. A graphical representation of absorption over time is shown.
Domain 6: Confounding/Variable Control				
	Metric 17:	Confounding variables in test design and procedures	Medium	Two batches of radiolabeled and non-radiolabeled 1,1-dichloroethane were used; it is unclear if batches were combined or if different batches were used for different experiments. Abdominal skin was obtained from males and females ranging in age from 30-70 years old. The large age range and sex differences may influence results. Skin integrity was confirmed by TEWL and only skin meeting inclusion criteria (≤ 13 gm-2h-1) were included in the analysis. This is a slight deviation from guidelines which suggest viable skin has a TEWL reading of less than 10 grams/m2/hour.
	Metric 18:	Confounding variables in outcomes unrelated to exposure	High	There were no reported differences among the study replicates that were unrelated to exposure; the test substance was demonstrated to be soluble in the receptor fluid.
Domain 7: Data Presentation and Analysis				
	Metric 19:	Data analysis	Uninformative	Means and standard deviations were calculated and reported. The CV for total absorbed dose is >25% for all dose groups/conditions. The high CV values make the study uninformative.

Continued on next page ...

... continued from previous page

Study Citation: Labcorp Early Development, (2024). 1,1-Dichloroethane: Rates of penetration through human skin using a flow through in vitro system.
HERO ID: 11396332

EVALUATION

Domain	Metric	EVALUATION Rating	Comments	
	Metric 20:	Data interpretation	Uninformative	The study reported dose in receptor fluid, receptor chamber, exposed skin, in stratum corneum, dislodgeable dose, donor chamber, in trap, and in tape strips. Data were interpreted correctly. The percent recoveries were low; 58.42% (neat), 54.36% (50% 1,1-dichloroethane in 1,2-dichloroethane) and 55.42% (1% dichloroethane in 1,2-dichloroethane). These low recoveries of radioactivity make this metric uninformative. The authors conducted separate experiments to investigate losses of radioactivity. Decreased recovery in some groups was most likely due to the volatility of the test substance.
	Metric 21:	Reporting of data	High	Data for all relevant endpoints were reported quantitatively as means ± SD. Individual cell data are shown.

Overall Quality Determination

Uninformative

Study Citation: Labcorp Early Development, (2024). 1,1-Dichloroethane: Rates of penetration through human skin using a flow through in vitro system.
HERO ID: 11396332

EXTRACTION

Parameter	Data
Extraction ID; Chemical:	Finite; neat; 1,1-Dichloroethane-Parent compound
Skin Material/Species; Skin Preparation; Skin Thickness (um); Diffusion Cell Exposure Setup Type:	ex vivo human; Split thickness; 400; Flow-through; Notes: Not Reported
Occlusion Type; Donor Chamber Vehicle; Concentration of Test Substance in Vehicle (enter as percent):	Semi-occluded (e.g., vapor trap); Not Reported; 100
Mass per Surface Area on Skin (mg/cm ²); Duration of Test Substance on Skin:	11.68; 8 hrs; Not Reported
Duration of Absorbance Measured; Frequency of Samples:	24 hrs; Samples were collected at 10 minutes, 30 minutes, 1 hours, 2 hours, and then 2-hourly intervals; Notes: Not Reported
Time Skin was Washed and Method used; Radiolabel Presence:	"The surface of the membrane was rinsed three times (3 x 0.5 mL) with a mild solution of Dove™ soap in water (ca. 1%). The skin surface was swabbed twice with cotton buds soaked with the liquid soap solution. Finally, the membrane was rinsed with a small volume (0.5 mL) of water and a further cotton bud used to swab the surface of the membrane until dry."; Yes
Total Recovery (percent); Dose Type:	58.42; Finite
Percent Found in Skin Depot After Washing and Tape Stripping; Comments:	0; Notes: Reported as <0.01%
Percent Found in All Tape Strips, Excluding the Upper Two Strips; Comments:	0; Notes: Reported as <0.01%
Percent Found in Receptor Fluid and Receptor Fluid Rinse; Comments:	0.13; Notes: CV =38.5%
Total Percent Absorbed:	0
Steady State Permeability Coefficient (Kp) (cm/hr); Steady State Permeability Coefficient (Comments); Steady State Flux (ug/cm ² /hr); Steady State Flux (Comments); Maximum Permeability Coefficient (Kp) (cm/hr); Maximum Permeability Coefficient (Comments); Maximum Flux (ug/cm ² /hr); Maximum Flux (Comments):	Not Reported; Notes: Not Reported; Not Reported; Notes: Not Reported; Not Reported; Notes: Not Reported; Not Reported; Notes: Not Reported

EVALUATION

Domain	Metric	Rating	Comments	
Domain 1: Test Substance				
	Metric 1:	Test substance identity	High	Test substances were identified as non-radiolabeled 1,1-dichloroethane and radiolabeled 1,1-14C dichloroethane, CAS number: 75-34-3; the structure was reported for the radiolabeled substance noting the location of the radiolabel within the substance.
	Metric 2:	Test substance source	High	The non-radiolabeled, 1,1-dichloroethane was sourced from LGC standards; the radiolabeled 1,1-dichloroethane was sourced from Moravek Biochemical Inc. The study included certificates of analyses that included lot/batch numbers.
	Metric 3:	Test substance purity	High	The radiochemical purity of the radiolabeled test substance was reported to be ≥98% by GC-MS. The purity of the non-radiolabeled substance was 99.48% (batch 1) and 99.9% (batch 2).

Continued on next page ...

... continued from previous page

Study Citation:		Labcorp Early Development, (2024). 1,1-Dichloroethane: Rates of penetration through human skin using a flow through in vitro system.		
HERO ID:		11396332		
		EVALUATION		
Domain	Metric	Rating	Comments	
Domain 2: Test Design	Metric 4:	Reference compounds	Medium	Testosterone (non-labelled and radiolabeled) was used as a reference compound in accordance with OECD 28 guidelines. 1 mg/ml was administered to the skin (0.011 mg/cm ²). Study authors report 92.1% recovery (CV = 2.57%). The CV values for some measurements were <25% (total dislodgeable dose, total unabsorbed dose, and the amount in skin washes), while others were >25% (i.e., total absorbed dose, dose in the skin, dose on surface tape and donor chambers etc.). It is unclear if these high CV values suggest any issues with the reliability of the tests conducted. However, this lab also cites previous publications that demonstrate the labs proficiency in performing dermal absorption studies with lipophilic compounds (Van de Sandt et al. 2004).
	Metric 5:	Assay procedures	Medium	This study was conducted according to OECD TG 428, OECD 28, and OECD 156. The assay procedures specified in the report were described in detail, although some information was missing. The flow through diffusion set-up was sufficiently reported including a schematic drawing. The study does not report if there was an equilibration period after the skin was placed into the chamber (guidelines recommend a 30-minute equilibrium period). Finite doses of 1,1-dichloroethane (neat, 50%, 10%, and 1% dissolved in either isopropyl myristate or 1,2-dichloroethane) were applied to human skin (at least 6 replicates; surface area of 0.64 cm ²). The volume applied (10uL/cm ²) applied to the skin surface was reported. The cells were semi-occluded by charcoal filters in the necks of the donor cells. The thickness of the skin samples ranged from 200 to 400 um, the exact thicknesses were not reported. The receptor solution was water fortified with 6% polyoxyethylene 20 oleyl ether. OECD 156 guidelines recommend <6% polyoxyethylene (20) oleyl ether in water for lipophilic compounds. The solubility of 1,1-dichloroethane was tested in the receptor fluid prior to the study and it was determined that the dissolution of 1,1-dichloroethane was not rate-limiting. The skin membranes were maintained at 32 degrees C; humidity ranged from 30-70%. The flow rate of receptor fluid was 1.5 mL/hour. After 8 hours skin was washed. "The surface of the membrane was rinsed three times (3 x 0.5 mL) with a mild solution of Dove™ soap in water (ca. 1%). The skin surface was swabbed twice with cotton buds soaked with the liquid soap solution. Finally, the membrane was rinsed with a small volume (0.5 mL) of water and a further cotton bud was used to swab the surface of the membrane until dry". For some groups, headspace above the skin was drawn into a syringe for analysis after 4 hours. Receptor fluid samples (volume not reported) were collected at 10 minutes, 30 minutes, 1 hour, 2 hours and every 2 hours henceforth for 24 hours (16 samples) and analyzed for radioactivity. After 24 hours, the skin was tape-stripped 5 times. "Radioactivity in gross amounts less than twice the background level was considered to be below the limit of accurate determination."

Continued on next page ...

... continued from previous page

Study Citation:		Labcorp Early Development, (2024). 1,1-Dichloroethane: Rates of penetration through human skin using a flow through in vitro system.		
HERO ID:		11396332		
		EVALUATION		
Domain	Metric	Rating	Comments	
	Metric 6:	Standards for tests	Low	The skin samples (abdomen) were obtained from female and male human donors following elective surgery (24 females and 2 males). Samples were stored at -20oC until used. The integrity of the skin was determined by stable trans epidermal water loss (TEWL) prior to dose application and at termination (prior to skin washing and tape stripping). A TEWL of ≤ 13 gm-2h-1 was considered acceptable. This is a slight deviation from guidelines which suggest viable skin has a TEWL reading of less than 10 grams/m2/hour. TEWL readings were reported. Coefficients of variation (CV) values were reported. CVs were all $> 25\%$. Further discussion of the CVs is provided in Metric 19. The percent recoveries were reported ($<60\%$) and were below the recommended $100 \pm 20\%$.
Domain 3: Exposure Characterization	Metric 7:	Preparation and storage of test substance (chemical)	Medium	The preparation of the test substance was adequately reported. Dilutions were magnetically stirred if deemed necessary by study authors (criteria not reported). The activity and homogeneity of the diluted doses were assessed from the top, middle, and bottom of the solution while being stirred. All doses were considered homogenous (data not shown). Storage conditions were not reported; it is unclear how far in advanced solutions were made, and how they were stored. However, the study included only a single application, and dosing was confirmed by analysis of QC samples taken during the dosing of the cells, therefore, the lack of storage information is not likely to have an impact on the study results. The solubility of the test substance in the receptor fluid was tested, (895 ug/mL) and was appropriate (>15 -fold higher than observed absorption) and was considered non-rate-limiting.
	Metric 8:	Consistency of exposure administration	Low	The volume of test substance applied to the skin was 10 uL/cm2 exposed skin (6.4 uL dose), which is in agreement with OECD guidelines. The skin surface area of 0.64 cm2 was consistent across groups. The skin thickness was reported as a range (200 to 400 uM). It is unclear if the variation in thicknesses was consistent across groups, and this may have contributed to some of the endpoint variations (and subsequently high CVs) observed.
	Metric 9:	Reporting of concentrations	High	The applied dose is reported as dpm, mg per cell, and mg/cm2. Individual cells are reported independently. Nominal and analytical doses are reported.
	Metric 10:	Exposure frequency	High	Exposure duration was reported and appropriate for determining absorption. The test substance was in contact with the skin for 8 hours prior to washing. Samples of receptor fluid were collected for a total of 24 hours.
	Metric 11:	Number of exposure groups and concentration spacing	High	There were 4 dose groups tested in a wide range of concentrations (neat to 1%). Dilutions of the test substance were performed in two different vehicles (isopropyl myristate or 1,2-dichloroethane). The study authors state "The dose levels selected in this study were selected to be representative of potential in-life exposure levels".
Domain 4: Test Model				
Continued on next page ...				

... continued from previous page

Study Citation:		Labcorp Early Development, (2024). 1,1-Dichloroethane: Rates of penetration through human skin using a flow through in vitro system.		
HERO ID:		11396332		
		EVALUATION		
Domain	Metric		Rating	Comments
	Metric 12:	Test model (skin)	High	The test model and descriptive information were reported. Samples of full-thickness skin samples were obtained from males and females following elective surgery. Abdominal skin samples from 26 people (race not reported) ranging in age from 30-70 years old were studied. The samples were stored frozen at -20 degrees C +/-10 degrees C. Prior to use, the skin samples were thawed, wiped to remove residual fat and blood, re-hydrated in purified water, and dermatomed using a mini-dermatome. The thickness ranged from 200 to 400 uM. The skin contained epidermis and some dermis. The OECD guidelines split thickness (dermatomed) skin is preferred. Membrane integrity was determined by measuring transepidermal water loss prior to/and upon completion of the experiment.
	Metric 13:	Number/Replicates per group	Medium	The number of replicates was appropriate as per OECD 428. The study used 6 replicates in order to ensure a minimum of 4 intact replicates could be obtained if skin integrity was not maintained.
Domain 5: Outcome Assessment				
	Metric 14:	Outcome assessment methodology	High	The outcome assessment methodology addressed the intended outcome of interest and was sensitive for the outcome. The test followed OECD guidelines 428 and 28. 10 uL/cm2 of the test substance was applied to skin samples. Measurement techniques and timing were reported and appropriate. A finite dose was used to determine absorption.
	Metric 15:	Consistency of outcome assessment	High	Details of outcome assessment protocol were reported and outcomes were assessed consistently across study replicates. The same duration of exposure, receptor fluid used, and sampling period were consistent across replicates.
	Metric 16:	Sampling adequacy and sensitivity	Medium	The study reported adequate sampling for the outcomes of interest; measurement sensitivity was sufficient. The sampling intervals were adequate (16 over 24 hours). Methods for the determination of radioactivity are reported. "Radioactivity in gross amounts less than twice the background level was considered to be below the limit of accurate determination." Scintillation counts were not shown. A graphical representation of absorption over time is shown.
Domain 6: Confounding/Variable Control				
	Metric 17:	Confounding variables in test design and procedures	Medium	Two batches of radiolabeled and non-radiolabeled 1,1-dichloroethane were used; it is unclear if batches were combined or if different batches were used for different experiments. Abdominal skin was obtained from males and females ranging in age from 30-70 years old. The large age range and sex differences may influence results. Skin integrity was confirmed by TEWL and only skin meeting inclusion criteria (≤ 13 gm-2h-1) were included in the analysis. This is a slight deviation from guidelines which suggest viable skin has a TEWL reading of less than 10 grams/m2/hour.
	Metric 18:	Confounding variables in outcomes unrelated to exposure	High	There were no reported differences among the study replicates that were unrelated to exposure; the test substance was demonstrated to be soluble in the receptor fluid.
Domain 7: Data Presentation and Analysis				
	Metric 19:	Data analysis	Uninformative	Means and standard deviations were calculated and reported. The CV for total absorbed dose is >25% for all dose groups/conditions. The high CV values make the study uninformative.

Continued on next page ...

... continued from previous page

Study Citation:		Labcorp Early Development, (2024). 1,1-Dichloroethane: Rates of penetration through human skin using a flow through in vitro system.		
HERO ID:		11396332		
EVALUATION				
Domain	Metric	EVALUATION Rating	Comments	
	Metric 20:	Data interpretation	Uninformative	The study reported dose in receptor fluid, receptor chamber, exposed skin, in stratum corneum, dislodgeable dose, donor chamber, in trap, and in tape strips. Data were interpreted correctly. The percent recoveries were low; 58.42% (neat), 54.36% (50% 1,1-dichloroethane in 1,2-dichloroethane) and 55.42% (1% dichloroethane in 1,2-dichloroethane). These low recoveries of radioactivity make this metric uninformative. The authors conducted separate experiments to investigate losses of radioactivity. Decreased recovery in some groups was most likely due to the volatility of the test substance.
	Metric 21:	Reporting of data	High	Data for all relevant endpoints were reported quantitatively as means ± SD. Individual cell data are shown.

Overall Quality Determination	Uninformative
--------------------------------------	----------------------

Study Citation: Labcorp Early Development, (2024). 1,1-Dichloroethane: Rates of penetration through human skin using a flow through in vitro system.
HERO ID: 11396332

EXTRACTION

Parameter	Data
Extraction ID; Chemical:	Finite; 50% in 1,2-dichloroethane; 1,1-Dichloroethane-Parent compound
Skin Material/Species; Skin Preparation; Skin Thickness (um); Diffusion Cell Exposure Setup Type:	ex vivo human; Split thickness; 400; Flow-through; Notes: Not Reported
Occlusion Type; Donor Chamber Vehicle; Concentration of Test Substance in Vehicle (enter as percent):	Semi-occluded (e.g., vapor trap); 1,2-dichloroethane; 50
Mass per Surface Area on Skin (mg/cm ²); Duration of Test Substance on Skin:	5.61; 8 hrs; Not Reported
Duration of Absorbance Measured; Frequency of Samples:	24 hrs; Samples were collected at 10 minutes, 30 minutes, 1 hours, 2 hours, and then 2-hourly intervals; Notes: Not Reported
Time Skin was Washed and Method used; Radiolabel Presence:	"The surface of the membrane was rinsed three times (3 x 0.5 mL) with a mild solution of Dove™ soap in water (ca. 1%). The skin surface was swabbed twice with cotton buds soaked with the liquid soap solution. Finally, the membrane was rinsed with a small volume (0.5 mL) of water and a further cotton bud used to swab the surface of the membrane until dry."; Yes
Total Recovery (percent); Dose Type:	54.36; Finite
Percent Found in Skin Depot After Washing and Tape Stripping; Comments:	0; Notes: not detected
Percent Found in All Tape Strips, Excluding the Upper Two Strips; Comments:	0; Notes: not detected
Percent Found in Receptor Fluid and Receptor Fluid Rinse; Comments:	0.12; Notes: CV= 75%
Total Percent Absorbed:	0
Steady State Permeability Coefficient (Kp) (cm/hr); Steady State Permeability Coefficient (Comments); Steady State Flux (ug/cm ² /hr); Steady State Flux (Comments); Maximum Permeability Coefficient (Kp) (cm/hr); Maximum Permeability Coefficient (Comments); Maximum Flux (ug/cm ² /hr); Maximum Flux (Comments):	Not Reported; Notes: Not Reported; Not Reported; Notes: Not Reported; Not Reported; Notes: Not Reported; Not Reported; Notes: Not Reported

EVALUATION

Domain	Metric	Rating	Comments
Domain 1: Test Substance			
	Metric 1:	High	Test substances were identified as non-radiolabeled 1,1-dichloroethane and radiolabeled 1,1-14C dichloroethane, CAS number: 75-34-3; the structure was reported for the radiolabeled substance noting the location of the radiolabel within the substance.
	Metric 2:	High	The non-radiolabeled, 1,1-dichloroethane was sourced from LGC standards; the radiolabeled 1,1-dichloroethane was sourced from Moravek Biochemical Inc. The study included certificates of analyses that included lot/batch numbers.
	Metric 3:	High	The radiochemical purity of the radiolabeled test substance was reported to be ≥98% by GC-MS. The purity of the non-radiolabeled substance was 99.48% (batch 1) and 99.9% (batch 2).

Continued on next page ...

... continued from previous page

Study Citation:		Labcorp Early Development, (2024). 1,1-Dichloroethane: Rates of penetration through human skin using a flow through in vitro system.		
HERO ID:		11396332		
		EVALUATION		
Domain	Metric	Rating	Comments	
Domain 2: Test Design	Metric 4:	Reference compounds	Medium	Testosterone (non-labelled and radiolabeled) was used as a reference compound in accordance with OECD 28 guidelines. 1 mg/ml was administered to the skin (0.011 mg/cm ²). Study authors report 92.1% recovery (CV = 2.57%). The CV values for some measurements were <25% (total dislodgeable dose, total unabsorbed dose, and the amount in skin washes), while others were >25% (i.e., total absorbed dose, dose in the skin, dose on surface tape and donor chambers etc.). It is unclear if these high CV values suggest any issues with the reliability of the tests conducted. However, this lab also cites previous publications that demonstrate the labs proficiency in performing dermal absorption studies with lipophilic compounds (Van de Sandt et al. 2004).
	Metric 5:	Assay procedures	Medium	This study was conducted according to OECD TG 428, OECD 28, and OECD 156. The assay procedures specified in the report were described in detail, although some information was missing. The flow through diffusion set-up was sufficiently reported including a schematic drawing. The study does not report if there was an equilibration period after the skin was placed into the chamber (guidelines recommend a 30-minute equilibrium period). Finite doses of 1,1-dichloroethane (neat, 50%, 10%, and 1% dissolved in either isopropyl myristate or 1,2-dichloroethane) were applied to human skin (at least 6 replicates; surface area of 0.64 cm ²). The volume applied (10uL/cm ²) applied to the skin surface was reported. The cells were semi-occluded by charcoal filters in the necks of the donor cells. The thickness of the skin samples ranged from 200 to 400 um, the exact thicknesses were not reported. The receptor solution was water fortified with 6% polyoxyethylene 20 oleyl ether. OECD 156 guidelines recommend <6% polyoxyethylene (20) oleyl ether in water for lipophilic compounds. The solubility of 1,1-dichloroethane was tested in the receptor fluid prior to the study and it was determined that the dissolution of 1,1-dichloroethane was not rate-limiting. The skin membranes were maintained at 32 degrees C; humidity ranged from 30-70%. The flow rate of receptor fluid was 1.5 mL/hour. After 8 hours skin was washed. "The surface of the membrane was rinsed three times (3 x 0.5 mL) with a mild solution of Dove™ soap in water (ca. 1%). The skin surface was swabbed twice with cotton buds soaked with the liquid soap solution. Finally, the membrane was rinsed with a small volume (0.5 mL) of water and a further cotton bud was used to swab the surface of the membrane until dry". For some groups, headspace above the skin was drawn into a syringe for analysis after 4 hours. Receptor fluid samples (volume not reported) were collected at 10 minutes, 30 minutes, 1 hour, 2 hours and every 2 hours henceforth for 24 hours (16 samples) and analyzed for radioactivity. After 24 hours, the skin was tape-stripped 5 times. "Radioactivity in gross amounts less than twice the background level was considered to be below the limit of accurate determination."

Continued on next page ...

... continued from previous page

Study Citation:		Labcorp Early Development, (2024). 1,1-Dichloroethane: Rates of penetration through human skin using a flow through in vitro system.		
HERO ID:		11396332		
		EVALUATION		
Domain	Metric	Rating	Comments	
	Metric 6:	Standards for tests	Low	The skin samples (abdomen) were obtained from female and male human donors following elective surgery (24 females and 2 males). Samples were stored at -20oC until used. The integrity of the skin was determined by stable trans epidermal water loss (TEWL) prior to dose application and at termination (prior to skin washing and tape stripping). A TEWL of ≤ 13 gm-2h-1 was considered acceptable. This is a slight deviation from guidelines which suggest viable skin has a TEWL reading of less than 10 grams/m2/hour. TEWL readings were reported. Coefficients of variation (CV) values were reported. CVs were all $> 25\%$. Further discussion of the CVs is provided in Metric 19. The percent recoveries were reported ($<60\%$) and were below the recommended 100 +/-20%.
Domain 3: Exposure Characterization	Metric 7:	Preparation and storage of test substance (chemical)	Medium	The preparation of the test substance was adequately reported. Dilutions were magnetically stirred if deemed necessary by study authors (criteria not reported). The activity and homogeneity of the diluted doses were assessed from the top, middle, and bottom of the solution while being stirred. All doses were considered homogenous (data not shown). Storage conditions were not reported; it is unclear how far in advanced solutions were made, and how they were stored. However, the study included only a single application, and dosing was confirmed by analysis of QC samples taken during the dosing of the cells, therefore, the lack of storage information is not likely to have an impact on the study results. The solubility of the test substance in the receptor fluid was tested, (895 ug/mL) and was appropriate (>15 -fold higher than observed absorption) and was considered non-rate-limiting.
	Metric 8:	Consistency of exposure administration	Low	The volume of test substance applied to the skin was 10 uL/cm2 exposed skin (6.4 uL dose), which is in agreement with OECD guidelines. The skin surface area of 0.64 cm2 was consistent across groups. The skin thickness was reported as a range (200 to 400 uM). It is unclear if the variation in thicknesses was consistent across groups, and this may have contributed to some of the endpoint variations (and subsequently high CVs) observed.
	Metric 9:	Reporting of concentrations	High	The applied dose is reported as dpm, mg per cell, and mg/cm2. Individual cells are reported independently. Nominal and analytical doses are reported.
	Metric 10:	Exposure frequency	High	Exposure duration was reported and appropriate for determining absorption. The test substance was in contact with the skin for 8 hours prior to washing. Samples of receptor fluid were collected for a total of 24 hours.
	Metric 11:	Number of exposure groups and concentration spacing	High	There were 4 dose groups tested in a wide range of concentrations (neat to 1%). Dilutions of the test substance were performed in two different vehicles (isopropyl myristate or 1,2-dichloroethane). The study authors state "The dose levels selected in this study were selected to be representative of potential in-life exposure levels".
Domain 4: Test Model				
Continued on next page ...				

... continued from previous page

Study Citation:		Labcorp Early Development, (2024). 1,1-Dichloroethane: Rates of penetration through human skin using a flow through in vitro system.		
HERO ID:		11396332		
		EVALUATION		
Domain	Metric		Rating	Comments
	Metric 12:	Test model (skin)	High	The test model and descriptive information were reported. Samples of full-thickness skin samples were obtained from males and females following elective surgery. Abdominal skin samples from 26 people (race not reported) ranging in age from 30-70 years old were studied. The samples were stored frozen at -20 degrees C +/-10 degrees C. Prior to use, the skin samples were thawed, wiped to remove residual fat and blood, re-hydrated in purified water, and dermatomed using a mini-dermatome. The thickness ranged from 200 to 400 uM. The skin contained epidermis and some dermis. The OECD guidelines split thickness (dermatomed) skin is preferred. Membrane integrity was determined by measuring transepidermal water loss prior to/and upon completion of the experiment.
	Metric 13:	Number/Replicates per group	Medium	The number of replicates was appropriate as per OECD 428. The study used 6 replicates in order to ensure a minimum of 4 intact replicates could be obtained if skin integrity was not maintained.
Domain 5: Outcome Assessment				
	Metric 14:	Outcome assessment methodology	High	The outcome assessment methodology addressed the intended outcome of interest and was sensitive for the outcome. The test followed OECD guidelines 428 and 28. 10 uL/cm2 of the test substance was applied to skin samples. Measurement techniques and timing were reported and appropriate. A finite dose was used to determine absorption.
	Metric 15:	Consistency of outcome assessment	High	Details of outcome assessment protocol were reported and outcomes were assessed consistently across study replicates. The same duration of exposure, receptor fluid used, and sampling period were consistent across replicates.
	Metric 16:	Sampling adequacy and sensitivity	Medium	The study reported adequate sampling for the outcomes of interest; measurement sensitivity was sufficient. The sampling intervals were adequate (16 over 24 hours). Methods for the determination of radioactivity are reported. "Radioactivity in gross amounts less than twice the background level was considered to be below the limit of accurate determination." Scintillation counts were not shown. A graphical representation of absorption over time is shown.
Domain 6: Confounding/Variable Control				
	Metric 17:	Confounding variables in test design and procedures	Medium	Two batches of radiolabeled and non-radiolabeled 1,1-dichloroethane were used; it is unclear if batches were combined or if different batches were used for different experiments. Abdominal skin was obtained from males and females ranging in age from 30-70 years old. The large age range and sex differences may influence results. Skin integrity was confirmed by TEWL and only skin meeting inclusion criteria (≤ 13 gm-2h-1) were included in the analysis. This is a slight deviation from guidelines which suggest viable skin has a TEWL reading of less than 10 grams/m2/hour.
	Metric 18:	Confounding variables in outcomes unrelated to exposure	High	There were no reported differences among the study replicates that were unrelated to exposure; the test substance was demonstrated to be soluble in the receptor fluid.
Domain 7: Data Presentation and Analysis				
	Metric 19:	Data analysis	Uninformative	Means and standard deviations were calculated and reported. The CV for total absorbed dose is >25% for all dose groups/conditions. The high CV values make the study uninformative.

Continued on next page ...

... continued from previous page

Study Citation: Labcorp Early Development, (2024). 1,1-Dichloroethane: Rates of penetration through human skin using a flow through in vitro system.
HERO ID: 11396332

EVALUATION

Domain	Metric	Rating	Comments
	Metric 20: Data interpretation	Uninformative	The study reported dose in receptor fluid, receptor chamber, exposed skin, in stratum corneum, dislodgeable dose, donor chamber, in trap, and in tape strips. Data were interpreted correctly. The percent recoveries were low; 58.42% (neat), 54.36% (50% 1,1-dichloroethane in 1,2-dichloroethane) and 55.42% (1% dichloroethane in 1,2-dichloroethane). These low recoveries of radioactivity make this metric uninformative. The authors conducted separate experiments to investigate losses of radioactivity. Decreased recovery in some groups was most likely due to the volatility of the test substance.
	Metric 21: Reporting of data	High	Data for all relevant endpoints were reported quantitatively as means ± SD. Individual cell data are shown.

Overall Quality Determination

Uninformative

Study Citation: Labcorp Early Development, (2024). 1,1-Dichloroethane: Rates of penetration through human skin using a flow through in vitro system.
HERO ID: 11396332

EXTRACTION

Parameter	Data
Extraction ID; Chemical:	Finite; 1% in IPM; 1,1-Dichloroethane-Parent compound
Skin Material/Species; Skin Preparation; Skin Thickness (um); Diffusion Cell Exposure Setup Type:	ex vivo human; Split thickness; 400; Flow-through; Notes: Not Reported
Occlusion Type; Donor Chamber Vehicle; Concentration of Test Substance in Vehicle (enter as percent):	Semi-occluded (e.g., vapor trap); isopropyl myristate; 1
Mass per Surface Area on Skin (mg/cm ²); Duration of Test Substance on Skin:	0.114; 8 hrs; Not Reported
Duration of Absorbance Measured; Frequency of Samples:	24 hrs; Samples were collected at 10 minutes, 30 minutes, 1 hours, 2 hours, and then 2-hourly intervals; Notes: Not Reported
Time Skin was Washed and Method used; Radiolabel Presence:	"The surface of the membrane was rinsed three times (3 x 0.5 mL) with a mild solution of Dove™ soap in water (ca. 1%). The skin surface was swabbed twice with cotton buds soaked with the liquid soap solution. Finally, the membrane was rinsed with a small volume (0.5 mL) of water and a further cotton bud used to swab the surface of the membrane until dry."; Yes
Total Recovery (percent); Dose Type:	88.87; Finite
Percent Found in Skin Depot After Washing and Tape Stripping; Comments:	0; Notes: not detected
Percent Found in All Tape Strips, Excluding the Upper Two Strips; Comments:	0; Notes: not detected
Percent Found in Receptor Fluid and Receptor Fluid Rinse; Comments:	0.02; Notes: CV= 200%
Total Percent Absorbed:	0
Steady State Permeability Coefficient (Kp) (cm/hr); Steady State Permeability Coefficient (Comments); Steady State Flux (ug/cm ² /hr); Steady State Flux (Comments); Maximum Permeability Coefficient (Kp) (cm/hr); Maximum Permeability Coefficient (Comments); Maximum Flux (ug/cm ² /hr); Maximum Flux (Comments):	Not Reported; Notes: Not Reported; Not Reported; Notes: Not Reported; Not Reported; Notes: Not Reported; Not Reported; Notes: Not Reported

EVALUATION

Domain	Metric	Rating	Comments	
Domain 1: Test Substance				
	Metric 1:	Test substance identity	High	Test substances were identified as non-radiolabeled 1,1-dichloroethane and radiolabeled 1,1-14C dichloroethane, CAS number: 75-34-3; the structure was reported for the radiolabeled substance noting the location of the radiolabel within the substance.
	Metric 2:	Test substance source	High	The non-radiolabeled, 1,1-dichloroethane was sourced from LGC standards; the radiolabeled 1,1-dichloroethane was sourced from Moravek Biochemical Inc. The study included certificates of analyses that included lot/batch numbers.
	Metric 3:	Test substance purity	High	The radiochemical purity of the radiolabeled test substance was reported to be ≥98% by GC-MS. The purity of the non-radiolabeled substance was 99.48% (batch 1) and 99.9% (batch 2).

Continued on next page ...

... continued from previous page

Study Citation:		Labcorp Early Development, (2024). 1,1-Dichloroethane: Rates of penetration through human skin using a flow through in vitro system.		
HERO ID:		11396332		
		EVALUATION		
Domain	Metric	Rating	Comments	
Domain 2: Test Design	Metric 4:	Reference compounds	Medium	Testosterone (non-labelled and radiolabeled) was used as a reference compound in accordance with OECD 28 guidelines. 1 mg/ml was administered to the skin (0.011 mg/cm ²). Study authors report 92.1% recovery (CV = 2.57%). The CV values for some measurements were <25% (total dislodgeable dose, total unabsorbed dose, and the amount in skin washes), while others were >25% (i.e., total absorbed dose, dose in the skin, dose on surface tape and donor chambers etc.). It is unclear if these high CV values suggest any issues with the reliability of the tests conducted. However, this lab also cites previous publications that demonstrate the labs proficiency in performing dermal absorption studies with lipophilic compounds (Van de Sandt et al. 2004).
	Metric 5:	Assay procedures	Medium	This study was conducted according to OECD TG 428, OECD 28, and OECD 156. The assay procedures specified in the report were described in detail, although some information was missing. The flow through diffusion set-up was sufficiently reported including a schematic drawing. The study does not report if there was an equilibration period after the skin was placed into the chamber (guidelines recommend a 30-minute equilibrium period). Finite doses of 1,1-dichloroethane (neat, 50%, 10%, and 1% dissolved in either isopropyl myristate or 1,2-dichloroethane) were applied to human skin (at least 6 replicates; surface area of 0.64 cm ²). The volume applied to the skin surface was 10uL/cm ² . The cells were semi-occluded by a charcoal filter in the neck of the donor cell. The thickness of the skin samples ranged from 200 to 400 um, exact thicknesses were not reported. The receptor solution was water fortified with 6% polyoxyethelene 20 oleyl ether. OECD 156 guidelines recommend <6% polyoxyethelene (20) oleyl ether in water for lipophilic compounds. The solubility of 1,1-dichloroethane was tested in the receptor fluid prior to the study and determined dissolution of 1,1-dichloroethane was not rate-limiting. The skin membranes were maintained at 32 degrees C; humidity ranged from 30-70%. The flow rate of receptor fluid was 1.5 mL/hour. After 8 hours skin was washed. "The surface of the membrane was rinsed three times (3 x 0.5 mL) with a mild solution of Dove™ soap in water (ca. 1%). The skin surface was swabbed twice with cotton buds soaked with the liquid soap solution. Finally, the membrane was rinsed with a small volume (0.5 mL) of water and a further cotton bud was used to swab the surface of the membrane until dry". For some groups, headspace above the skin was drawn into a syringe for analysis after 4 hours. Receptor fluid samples (volume not reported) were collected at 10 minutes, 30 minutes, 1 hour, 2 hours and every 2 hours henceforth for 24 hours (16 samples) and analyzed for radioactivity. After 24 hours, the skin was tap-stripped 5 times. "Radioactivity in gross amounts less than twice the background level was considered to be below the limit of accurate determination."
	Metric 6:	Standards for tests	Medium	The skin samples (abdomen) were obtained from 24 female and 2 male human donors following elective surgery. Samples were stored at -20oC until used. The integrity of the skin was determined by stable trans epidermal water loss (TEWL) prior to dose application and at termination (prior to skin washing and tape stripping). A TEWL of ≤ 13 gm-2h-1 was considered acceptable. This is a slight deviation from guidelines which suggest viable skin has a TEWL reading of less than 10 grams/m2/hour. TEWL readings were reported. Percent recovery was >80%, which is acceptable for this volatile substance. Coefficients of variation (CV) values were reported. CVs were all > 25%. Further discussion of the CVs is provided in Metric 19.

Continued on next page ...

... continued from previous page

Study Citation:		Labcorp Early Development, (2024). 1,1-Dichloroethane: Rates of penetration through human skin using a flow through in vitro system.		
HERO ID:		11396332		
		EVALUATION		
Domain	Metric	Rating	Comments	
Domain 3: Exposure Characterization				
	Metric 7:	Preparation and storage of test substance (chemical)	Medium	The preparation of the test substance was adequately reported. Dilutions were magnetically stirred if deemed necessary by study authors (criteria not reported). The activity and homogeneity of the diluted doses were assessed from the top, middle, and bottom of the solution while being stirred. All doses were considered homogenous (data not shown). Storage conditions are not reported; it is unclear how far in advanced solutions were made, and how they were stored. However, the study included only a single application, and dosing was confirmed by analysis of QC samples taken during the dosing of the cells, therefore, the lack of storage information is not likely to have an impact on the study results. The solubility of the test substance in the receptor fluid was tested, (895 ug/mL) and was appropriate (>15-fold higher than observed absorption) and was considered non-rate-limiting.
	Metric 8:	Consistency of exposure administration	Low	The volume of test substance applied to the skin was 10 uL/cm2 exposed skin (6.4 uL dose), which is in agreement with OECD guidelines. The skin surface area of 0.64 cm2 was consistent across groups. The skin thickness was reported as a range (200 to 400 uM). It is unclear if the variation in thicknesses was consistent across groups, and this may have contributed to some of the endpoint variations (and subsequently high CVs) observed.
	Metric 9:	Reporting of concentrations	High	The applied dose is reported as dpm, mg per cell, and mg/cm2. Individual cells are reported independently. Nominal and analytical doses are reported.
	Metric 10:	Exposure frequency	High	Exposure duration was reported and appropriate for determining absorption. Test substance was in contact with the skin for 8 hours prior to washing. Samples of receptor fluid were collected for a total of 24 hours.
	Metric 11:	Number of exposure groups and concentration spacing	High	There were 4 dose groups tested in a wide range of concentrations (neat to 1%). Dilutions of test substance was performed in two different vehicles (isopropyl myristate or 1,2-dichloroethane). Study authors state "The dose levels selected in this study were selected to be representative of potential in-life exposure levels".
Domain 4: Test Model				
	Metric 12:	Test model (skin)	High	The test model and descriptive information were reported. Samples of full thickness skin samples were obtained from males and females following elective surgery. Abdominal skin samples from 26 people (race not reported) ranging in age from 30-70 years old were studied. The samples were stored frozen at -20 degrees C +/- 10 degrees C. Prior to use, the skin samples were thawed, wiped to remove residual fat and blood, re-hydrated in purified water, and dermatomed using a mini-dermatome. The thickness ranged from 200 to 400 uM. The skin contained epidermis and some dermis. The OECD guidelines split thickness (dermatomed) skin is preferred. Membrane integrity was determined by measuring trans epidermal water loss prior to/and upon completion of the experiment.
	Metric 13:	Number/Replicates per group	Medium	The number of replicates was appropriate as per OECD 428. The study used 6 replicates to ensure a minimum of 4 intact replicates could be obtained if skin integrity was not maintained.
Domain 5: Outcome Assessment				
Continued on next page ...				

... continued from previous page

Study Citation: HERO ID:		Labcorp Early Development, (2024). 1,1-Dichloroethane: Rates of penetration through human skin using a flow through in vitro system. 11396332		
		EVALUATION		
Domain	Metric	Rating	Comments	
	Metric 14:	Outcome assessment methodology	High	The outcome assessment methodology addressed the intended outcome of interest and was sensitive for the outcome. The test followed OECD guidelines 428 and 28. 10 uL/cm ² of test substance was applied to skin samples. Measurement techniques and timing were reported and appropriate. A finite dose was used to determine absorption.
	Metric 15:	Consistency of outcome assessment	High	Details of outcome assessment protocol were reported and outcomes were assessed consistently across study replicates. The same duration of exposure, receptor fluid used, and sampling period were consistent across replicates.
	Metric 16:	Sampling adequacy and sensitivity	Medium	The study reported adequate sampling for the outcomes of interest; measurement sensitivity was sufficient. The sampling intervals were adequate (16 over 24 hours). Methods for the determination of radioactivity are reported. "Radioactivity in gross amounts less than twice the background level was considered to be below the limit of accurate determination." Scintillation counts were not shown. A graphical representation of absorption over time is shown.
Domain 6: Confounding/Variable Control				
	Metric 17:	Confounding variables in test design and procedures	Medium	Two batches of radiolabeled and non-radiolabeled 1,1-dichloroethane were used; it is unclear if batches were combined or if different batches were used for different experiments. Abdominal skin was obtained from males and females ranging in age from 30-70 years old. The large age range and sex differences may influence results. Skin integrity was confirmed by TEWL and only skin meeting inclusion criteria (≤ 13 gm-2h-1) was included in the analysis. This is a slight deviation from guidelines which suggest viable skin has a TEWL reading of less than 10 grams/m ² /hour.
	Metric 18:	Confounding variables in outcomes unrelated to exposure	High	There were no reported differences among the study replicates that were unrelated to exposure; the test substance was demonstrated to be soluble in the receptor fluid.
Domain 7: Data Presentation and Analysis				
	Metric 19:	Data analysis	Uninformative	Means and standard deviations were calculated and reported. The CV for total absorbed dose is $>25\%$ for all dose groups/conditions. The high CV values make the study uninformative.
	Metric 20:	Data interpretation	High	The study reported doses in receptor fluid, receptor chamber, exposed skin, in stratum corneum, dislodgeable dose, donor chamber, in trap, and in tape strips. Data were interpreted correctly. Decreased recovery in some groups was most likely due to the volatility of the test substance, although total recoveries were $>80\%$.
	Metric 21:	Reporting of data	High	Data for all relevant endpoints were reported quantitatively as means \pm SD. Individual cell data are shown.
Overall Quality Determination		Uninformative		

Study Citation: Labcorp Early Development, (2024). 1,1-Dichloroethane: Rates of penetration through human skin using a flow through in vitro system.
HERO ID: 11396332

EXTRACTION

Parameter	Data
Extraction ID; Chemical:	Finite; 50% in IPM; 1,1-Dichloroethane-Parent compound
Skin Material/Species; Skin Preparation; Skin Thickness (um); Diffusion Cell Exposure Setup Type:	ex vivo human; Split thickness; 400; Flow-through; Notes: Not Reported
Occlusion Type; Donor Chamber Vehicle; Concentration of Test Substance in Vehicle (enter as percent):	Semi-occluded (e.g., vapor trap); isopropyl myristate; 50
Mass per Surface Area on Skin (mg/cm ²); Duration of Test Substance on Skin:	5.28; 8 hrs; Not Reported
Duration of Absorbance Measured; Frequency of Samples:	24 hrs; Samples were collected at 10 minutes, 30 minutes, 1 hours, 2 hours, and then 2-hourly intervals; Notes: Not Reported
Time Skin was Washed and Method used; Radiolabel Presence:	"The surface of the membrane was rinsed three times (3 x 0.5 mL) with a mild solution of Dove™ soap in water (ca. 1%). The skin surface was swabbed twice with cotton buds soaked with the liquid soap solution. Finally, the membrane was rinsed with a small volume (0.5 mL) of water and a further cotton bud used to swab the surface of the membrane until dry."; Yes
Total Recovery (percent); Dose Type:	87.66; Finite
Percent Found in Skin Depot After Washing and Tape Stripping; Comments:	0.01; Notes: CV= 200%
Percent Found in All Tape Strips, Excluding the Upper Two Strips; Comments:	0.01; Notes: CV= 200%
Percent Found in Receptor Fluid and Receptor Fluid Rinse; Comments:	0.05; Notes: CV= 60%
Total Percent Absorbed:	0
Steady State Permeability Coefficient (Kp) (cm/hr); Steady State Permeability Coefficient (Comments); Steady State Flux (ug/cm ² /hr); Steady State Flux (Comments); Maximum Permeability Coefficient (Kp) (cm/hr); Maximum Permeability Coefficient (Comments); Maximum Flux (ug/cm ² /hr); Maximum Flux (Comments):	Not Reported; Notes: Not Reported; Not Reported; Notes: Not Reported; Not Reported; Notes: Not Reported; Not Reported; Notes: Not Reported

EVALUATION

Domain	Metric	Rating	Comments	
Domain 1: Test Substance				
	Metric 1:	Test substance identity	High	Test substances were identified as non-radiolabeled 1,1-dichloroethane and radiolabeled 1,1-14C dichloroethane, CAS number: 75-34-3; the structure was reported for the radiolabeled substance noting the location of the radiolabel within the substance.
	Metric 2:	Test substance source	High	The non-radiolabeled, 1,1-dichloroethane was sourced from LGC standards; the radiolabeled 1,1-dichloroethane was sourced from Moravek Biochemical Inc. The study included certificates of analyses that included lot/batch numbers.
	Metric 3:	Test substance purity	High	The radiochemical purity of the radiolabeled test substance was reported to be ≥98% by GC-MS. The purity of the non-radiolabeled substance was 99.48% (batch 1) and 99.9% (batch 2).

Continued on next page ...

... continued from previous page

Study Citation:		Labcorp Early Development, (2024). 1,1-Dichloroethane: Rates of penetration through human skin using a flow through in vitro system.		
HERO ID:		11396332		
		EVALUATION		
Domain	Metric	Rating	Comments	
Domain 2: Test Design	Metric 4:	Reference compounds	Medium	Testosterone (non-labelled and radiolabeled) was used as a reference compound in accordance with OECD 28 guidelines. 1 mg/ml was administered to the skin (0.011 mg/cm ²). Study authors report 92.1% recovery (CV = 2.57%). The CV values for some measurements were <25% (total dislodgeable dose, total unabsorbed dose, and the amount in skin washes), while others were >25% (i.e., total absorbed dose, dose in the skin, dose on surface tape and donor chambers etc.). It is unclear if these high CV values suggest any issues with the reliability of the tests conducted. However, this lab also cites previous publications that demonstrate the labs proficiency in performing dermal absorption studies with lipophilic compounds (Van de Sandt et al. 2004).
	Metric 5:	Assay procedures	Medium	This study was conducted according to OECD TG 428, OECD 28, and OECD 156. The assay procedures specified in the report were described in detail, although some information was missing. The flow through diffusion set-up was sufficiently reported including a schematic drawing. The study does not report if there was an equilibration period after the skin was placed into the chamber (guidelines recommend a 30-minute equilibrium period). Finite doses of 1,1-dichloroethane (neat, 50%, 10%, and 1% dissolved in either isopropyl myristate or 1,2-dichloroethane) were applied to human skin (at least 6 replicates; surface area of 0.64 cm ²). The volume applied to the skin surface was 10uL/cm ² . The cells were semi-occluded by a charcoal filter in the neck of the donor cell. The thickness of the skin samples ranged from 200 to 400 um, exact thicknesses were not reported. The receptor solution was water fortified with 6% polyoxyethelene 20 oleyl ether. OECD 156 guidelines recommend <6% polyoxyethelene (20) oleyl ether in water for lipophilic compounds. The solubility of 1,1-dichloroethane was tested in the receptor fluid prior to the study and determined dissolution of 1,1-dichloroethane was not rate-limiting. The skin membranes were maintained at 32 degrees C; humidity ranged from 30-70%. The flow rate of receptor fluid was 1.5 mL/hour. After 8 hours skin was washed. "The surface of the membrane was rinsed three times (3 x 0.5 mL) with a mild solution of Dove™ soap in water (ca. 1%). The skin surface was swabbed twice with cotton buds soaked with the liquid soap solution. Finally, the membrane was rinsed with a small volume (0.5 mL) of water and a further cotton bud was used to swab the surface of the membrane until dry". For some groups, headspace above the skin was drawn into a syringe for analysis after 4 hours. Receptor fluid samples (volume not reported) were collected at 10 minutes, 30 minutes, 1 hour, 2 hours and every 2 hours henceforth for 24 hours (16 samples) and analyzed for radioactivity. After 24 hours, the skin was tap-stripped 5 times. "Radioactivity in gross amounts less than twice the background level was considered to be below the limit of accurate determination."
	Metric 6:	Standards for tests	Medium	The skin samples (abdomen) were obtained from 24 female and 2 male human donors following elective surgery. Samples were stored at -20oC until used. The integrity of the skin was determined by stable trans epidermal water loss (TEWL) prior to dose application and at termination (prior to skin washing and tape stripping). A TEWL of ≤ 13 gm-2h-1 was considered acceptable. This is a slight deviation from guidelines which suggest viable skin has a TEWL reading of less than 10 grams/m2/hour. TEWL readings were reported. Percent recovery was >80%, which is acceptable for this volatile substance. Coefficients of variation (CV) values were reported. CVs were all > 25%. Further discussion of the CVs is provided in Metric 19.

Continued on next page ...

... continued from previous page

Study Citation:		Labcorp Early Development, (2024). 1,1-Dichloroethane: Rates of penetration through human skin using a flow through in vitro system.		
HERO ID:		11396332		
		EVALUATION		
Domain	Metric	Rating	Comments	
Domain 3: Exposure Characterization				
	Metric 7:	Preparation and storage of test substance (chemical)	Medium	The preparation of the test substance was adequately reported. Dilutions were magnetically stirred if deemed necessary by study authors (criteria not reported). The activity and homogeneity of the diluted doses were assessed from the top, middle, and bottom of the solution while being stirred. All doses were considered homogenous (data not shown). Storage conditions are not reported; it is unclear how far in advanced solutions were made, and how they were stored. However, the study included only a single application, and dosing was confirmed by analysis of QC samples taken during the dosing of the cells, therefore, the lack of storage information is not likely to have an impact on the study results. The solubility of the test substance in the receptor fluid was tested, (895 ug/mL) and was appropriate (>15-fold higher than observed absorption) and was considered non-rate-limiting.
	Metric 8:	Consistency of exposure administration	Low	The volume of test substance applied to the skin was 10 uL/cm ² exposed skin (6.4 uL dose), which is in agreement with OECD guidelines. The skin surface area of 0.64 cm ² was consistent across groups. The skin thickness was reported as a range (200 to 400 uM). It is unclear if the variation in thicknesses was consistent across groups, and this may have contributed to some of the endpoint variations (and subsequently high CVs) observed.
	Metric 9:	Reporting of concentrations	High	The applied dose is reported as dpm, mg per cell, and mg/cm ² . Individual cells are reported independently. Nominal and analytical doses are reported.
	Metric 10:	Exposure frequency	High	Exposure duration was reported and appropriate for determining absorption. Test substance was in contact with the skin for 8 hours prior to washing. Samples of receptor fluid were collected for a total of 24 hours.
	Metric 11:	Number of exposure groups and concentration spacing	High	There were 4 dose groups tested in a wide range of concentrations (neat to 1%). Dilutions of test substance was performed in two different vehicles (isopropyl myristate or 1,2-dichloroethane). Study authors state "The dose levels selected in this study were selected to be representative of potential in-life exposure levels".
Domain 4: Test Model				
	Metric 12:	Test model (skin)	High	The test model and descriptive information were reported. Samples of full thickness skin samples were obtained from males and females following elective surgery. Abdominal skin samples from 26 people (race not reported) ranging in age from 30-70 years old were studied. The samples were stored frozen at -20 degrees C +/- 10 degrees C. Prior to use, the skin samples were thawed, wiped to remove residual fat and blood, re-hydrated in purified water, and dermatomed using a mini-dermatome. The thickness ranged from 200 to 400 uM. The skin contained epidermis and some dermis. The OECD guidelines split thickness (dermatomed) skin is preferred. Membrane integrity was determined by measuring trans epidermal water loss prior to/and upon completion of the experiment.
	Metric 13:	Number/Replicates per group	Medium	The number of replicates was appropriate as per OECD 428. The study used 6 replicates to ensure a minimum of 4 intact replicates could be obtained if skin integrity was not maintained.
Domain 5: Outcome Assessment				
Continued on next page ...				

... continued from previous page

Study Citation: HERO ID:		Labcorp Early Development, (2024). 1,1-Dichloroethane: Rates of penetration through human skin using a flow through in vitro system. 11396332		
		EVALUATION		
Domain	Metric		Rating	Comments
	Metric 14:	Outcome assessment methodology	High	The outcome assessment methodology addressed the intended outcome of interest and was sensitive for the outcome. The test followed OECD guidelines 428 and 28. 10 uL/cm ² of test substance was applied to skin samples. Measurement techniques and timing were reported and appropriate. A finite dose was used to determine absorption.
	Metric 15:	Consistency of outcome assessment	High	Details of outcome assessment protocol were reported and outcomes were assessed consistently across study replicates. The same duration of exposure, receptor fluid used, and sampling period were consistent across replicates.
	Metric 16:	Sampling adequacy and sensitivity	Medium	The study reported adequate sampling for the outcomes of interest; measurement sensitivity was sufficient. The sampling intervals were adequate (16 over 24 hours). Methods for the determination of radioactivity are reported. "Radioactivity in gross amounts less than twice the background level was considered to be below the limit of accurate determination." Scintillation counts were not shown. A graphical representation of absorption over time is shown.
Domain 6: Confounding/Variable Control				
	Metric 17:	Confounding variables in test design and procedures	Medium	Two batches of radiolabeled and non-radiolabeled 1,1-dichloroethane were used; it is unclear if batches were combined or if different batches were used for different experiments. Abdominal skin was obtained from males and females ranging in age from 30-70 years old. The large age range and sex differences may influence results. Skin integrity was confirmed by TEWL and only skin meeting inclusion criteria (≤ 13 gm-2h-1) was included in the analysis. This is a slight deviation from guidelines which suggest viable skin has a TEWL reading of less than 10 grams/m ² /hour.
	Metric 18:	Confounding variables in outcomes unrelated to exposure	High	There were no reported differences among the study replicates that were unrelated to exposure; the test substance was demonstrated to be soluble in the receptor fluid.
Domain 7: Data Presentation and Analysis				
	Metric 19:	Data analysis	Uninformative	Means and standard deviations were calculated and reported. The CV for total absorbed dose is $>25\%$ for all dose groups/conditions. The high CV values make the study uninformative.
	Metric 20:	Data interpretation	High	The study reported doses in receptor fluid, receptor chamber, exposed skin, in stratum corneum, dislodgeable dose, donor chamber, in trap, and in tape strips. Data were interpreted correctly. Decreased recovery in some groups was most likely due to the volatility of the test substance, although total recoveries were $>80\%$.
	Metric 21:	Reporting of data	High	Data for all relevant endpoints were reported quantitatively as means \pm SD. Individual cell data are shown.
Overall Quality Determination		Uninformative		

Study Citation: Labcorp Early Development, (2024). 1,1-Dichloroethane: Rates of penetration through human skin using a flow through in vitro system.
HERO ID: 11396332

EXTRACTION

Parameter	Data
Extraction ID; Chemical:	Finite; 10% in 1,2-dichloroethane; 1,1-Dichloroethane-Parent compound
Skin Material/Species; Skin Preparation; Skin Thickness (um); Diffusion Cell Exposure Setup Type:	ex vivo human; Split thickness; 400; Flow-through; Notes: Not Reported
Occlusion Type; Donor Chamber Vehicle; Concentration of Test Substance in Vehicle (enter as percent):	Semi-occluded (e.g., vapor trap); 1,2-dichloroethane; 10
Mass per Surface Area on Skin (mg/cm ²); Duration of Test Substance on Skin:	0.93; 8 hrs; Not Reported
Duration of Absorbance Measured; Frequency of Samples:	24 hrs; Samples were collected at 10 minutes, 30 minutes, 1 hours, 2 hours, and then 2-hourly intervals; Notes: Not Reported
Time Skin was Washed and Method used; Radiolabel Presence:	"The surface of the membrane was rinsed three times (3 x 0.5 mL) with a mild solution of Dove™ soap in water (ca. 1%). The skin surface was swabbed twice with cotton buds soaked with the liquid soap solution. Finally, the membrane was rinsed with a small volume (0.5 mL) of water and a further cotton bud used to swab the surface of the membrane until dry."; Yes
Total Recovery (percent); Dose Type:	92.76; Finite
Percent Found in Skin Depot After Washing and Tape Stripping; Comments:	0; Notes: not detected
Percent Found in All Tape Strips, Excluding the Upper Two Strips; Comments:	0; Notes: not detected
Percent Found in Receptor Fluid and Receptor Fluid Rinse; Comments:	0.02; Notes: CV= 100%
Total Percent Absorbed:	0
Steady State Permeability Coefficient (Kp) (cm/hr); Steady State Permeability Coefficient (Comments); Steady State Flux (ug/cm ² /hr); Steady State Flux (Comments); Maximum Permeability Coefficient (Kp) (cm/hr); Maximum Permeability Coefficient (Comments); Maximum Flux (ug/cm ² /hr); Maximum Flux (Comments):	Not Reported; Notes: Not Reported; Not Reported; Notes: Not Reported; Not Reported; Notes: Not Reported; Not Reported; Notes: Not Reported

EVALUATION

Domain	Metric	Rating	Comments
Domain 1: Test Substance			
	Metric 1: Test substance identity	High	Test substances were identified as non-radiolabeled 1,1-dichloroethane and radiolabeled 1,1-14C dichloroethane, CAS number: 75-34-3; the structure was reported for the radiolabeled substance noting the location of the radiolabel within the substance.
	Metric 2: Test substance source	High	The non-radiolabeled, 1,1-dichloroethane was sourced from LGC standards; the radiolabeled 1,1-dichloroethane was sourced from Moravek Biochemical Inc. The study included certificates of analyses that included lot/batch numbers.
	Metric 3: Test substance purity	High	The radiochemical purity of the radiolabeled test substance was reported to be ≥98% by GC-MS. The purity of the non-radiolabeled substance was 99.48% (batch 1) and 99.9% (batch 2).

Continued on next page ...

... continued from previous page

Study Citation:		Labcorp Early Development, (2024). 1,1-Dichloroethane: Rates of penetration through human skin using a flow through in vitro system.		
HERO ID:		11396332		
Domain	Metric	EVALUATION Rating	Comments	
Domain 2: Test Design	Metric 4:	Reference compounds	Medium	Testosterone (non-labelled and radiolabeled) was used as a reference compound in accordance with OECD 28 guidelines. 1 mg/ml was administered to the skin (0.011 mg/cm ²). Study authors report 92.1% recovery (CV = 2.57%). The CV values for some measurements were <25% (total dislodgeable dose, total unabsorbed dose, and the amount in skin washes), while others were >25% (i.e., total absorbed dose, dose in the skin, dose on surface tape and donor chambers etc.). It is unclear if these high CV values suggest any issues with the reliability of the tests conducted. However, this lab also cites previous publications that demonstrate the labs proficiency in performing dermal absorption studies with lipophilic compounds (Van de Sandt et al. 2004).
	Metric 5:	Assay procedures	Medium	This study was conducted according to OECD TG 428, OECD 28, and OECD 156. The assay procedures specified in the report were described in detail, although some information was missing. The flow through diffusion set-up was sufficiently reported including a schematic drawing. The study does not report if there was an equilibration period after the skin was placed into the chamber (guidelines recommend a 30-minute equilibrium period). Finite doses of 1,1-dichloroethane (neat, 50%, 10%, and 1% dissolved in either isopropyl myristate or 1,2-dichloroethane) were applied to human skin (at least 6 replicates; surface area of 0.64 cm ²). The volume applied to the skin surface was 10uL/cm ² . The cells were semi-occluded by a charcoal filter in the neck of the donor cell. The thickness of the skin samples ranged from 200 to 400 um, exact thicknesses were not reported. The receptor solution was water fortified with 6% polyoxyethelene 20 oleyl ether. OECD 156 guidelines recommend <6% polyoxyethelene (20) oleyl ether in water for lipophilic compounds. The solubility of 1,1-dichloroethane was tested in the receptor fluid prior to the study and determined dissolution of 1,1-dichloroethane was not rate-limiting. The skin membranes were maintained at 32 degrees C; humidity ranged from 30-70%. The flow rate of receptor fluid was 1.5 mL/hour. After 8 hours skin was washed. "The surface of the membrane was rinsed three times (3 x 0.5 mL) with a mild solution of Dove™ soap in water (ca. 1%). The skin surface was swabbed twice with cotton buds soaked with the liquid soap solution. Finally, the membrane was rinsed with a small volume (0.5 mL) of water and a further cotton bud was used to swab the surface of the membrane until dry". For some groups, headspace above the skin was drawn into a syringe for analysis after 4 hours. Receptor fluid samples (volume not reported) were collected at 10 minutes, 30 minutes, 1 hour, 2 hours and every 2 hours henceforth for 24 hours (16 samples) and analyzed for radioactivity. After 24 hours, the skin was tap-stripped 5 times. "Radioactivity in gross amounts less than twice the background level was considered to be below the limit of accurate determination."
	Metric 6:	Standards for tests	Medium	The skin samples (abdomen) were obtained from 24 female and 2 male human donors following elective surgery. Samples were stored at -20oC until used. The integrity of the skin was determined by stable trans epidermal water loss (TEWL) prior to dose application and at termination (prior to skin washing and tape stripping). A TEWL of ≤ 13 gm-2h-1 was considered acceptable. This is a slight deviation from guidelines which suggest viable skin has a TEWL reading of less than 10 grams/m2/hour. TEWL readings were reported. Percent recovery was >80%, which is acceptable for this volatile substance. Coefficients of variation (CV) values were reported. CVs were all > 25%. Further discussion of the CVs is provided in Metric 19.

Continued on next page ...

... continued from previous page

Study Citation:		Labcorp Early Development, (2024). 1,1-Dichloroethane: Rates of penetration through human skin using a flow through in vitro system.		
HERO ID:		11396332		
Domain	Metric	EVALUATION Rating	Comments	
Domain 3: Exposure Characterization				
Metric 7:	Preparation and storage of test substance (chemical)	Medium	The preparation of the test substance was adequately reported. Dilutions were magnetically stirred if deemed necessary by study authors (criteria not reported). The activity and homogeneity of the diluted doses were assessed from the top, middle, and bottom of the solution while being stirred. All doses were considered homogenous (data not shown). Storage conditions are not reported; it is unclear how far in advanced solutions were made, and how they were stored. However, the study included only a single application, and dosing was confirmed by analysis of QC samples taken during the dosing of the cells, therefore, the lack of storage information is not likely to have an impact on the study results. The solubility of the test substance in the receptor fluid was tested, (895 ug/mL) and was appropriate (>15-fold higher than observed absorption) and was considered non-rate-limiting.	
Metric 8:	Consistency of exposure administration	Low	The volume of test substance applied to the skin was 10 uL/cm2 exposed skin (6.4 uL dose), which is in agreement with OECD guidelines. The skin surface area of 0.64 cm2 was consistent across groups. The skin thickness was reported as a range (200 to 400 uM). It is unclear if the variation in thicknesses was consistent across groups, and this may have contributed to some of the endpoint variations (and subsequently high CVs) observed.	
Metric 9:	Reporting of concentrations	High	The applied dose is reported as dpm, mg per cell, and mg/cm2. Individual cells are reported independently. Nominal and analytical doses are reported.	
Metric 10:	Exposure frequency	High	Exposure duration was reported and appropriate for determining absorption. Test substance was in contact with the skin for 8 hours prior to washing. Samples of receptor fluid were collected for a total of 24 hours.	
Metric 11:	Number of exposure groups and concentration spacing	High	There were 4 dose groups tested in a wide range of concentrations (neat to 1%). Dilutions of test substance was performed in two different vehicles (isopropyl myristate or 1,2-dichloroethane). Study authors state "The dose levels selected in this study were selected to be representative of potential in-life exposure levels".	
Domain 4: Test Model				
Metric 12:	Test model (skin)	High	The test model and descriptive information were reported. Samples of full thickness skin samples were obtained from males and females following elective surgery. Abdominal skin samples from 26 people (race not reported) ranging in age from 30-70 years old were studied. The samples were stored frozen at -20 degrees C +/- 10 degrees C. Prior to use, the skin samples were thawed, wiped to remove residual fat and blood, re-hydrated in purified water, and dermatomed using a mini-dermatome. The thickness ranged from 200 to 400 uM. The skin contained epidermis and some dermis. The OECD guidelines split thickness (dermatomed) skin is preferred. Membrane integrity was determined by measuring trans epidermal water loss prior to/and upon completion of the experiment.	
Metric 13:	Number/Replicates per group	Medium	The number of replicates was appropriate as per OECD 428. The study used 6 replicates to ensure a minimum of 4 intact replicates could be obtained if skin integrity was not maintained.	
Domain 5: Outcome Assessment				
Continued on next page ...				

... continued from previous page

Study Citation: HERO ID:		Labcorp Early Development, (2024). 1,1-Dichloroethane: Rates of penetration through human skin using a flow through in vitro system. 11396332		
		EVALUATION		
Domain	Metric	Rating	Comments	
	Metric 14:	Outcome assessment methodology	High	The outcome assessment methodology addressed the intended outcome of interest and was sensitive for the outcome. The test followed OECD guidelines 428 and 28. 10 uL/cm ² of test substance was applied to skin samples. Measurement techniques and timing were reported and appropriate. A finite dose was used to determine absorption.
	Metric 15:	Consistency of outcome assessment	High	Details of outcome assessment protocol were reported and outcomes were assessed consistently across study replicates. The same duration of exposure, receptor fluid used, and sampling period were consistent across replicates.
	Metric 16:	Sampling adequacy and sensitivity	Medium	The study reported adequate sampling for the outcomes of interest; measurement sensitivity was sufficient. The sampling intervals were adequate (16 over 24 hours). Methods for the determination of radioactivity are reported. "Radioactivity in gross amounts less than twice the background level was considered to be below the limit of accurate determination." Scintillation counts were not shown. A graphical representation of absorption over time is shown.
Domain 6: Confounding/Variable Control				
	Metric 17:	Confounding variables in test design and procedures	Medium	Two batches of radiolabeled and non-radiolabeled 1,1-dichloroethane were used; it is unclear if batches were combined or if different batches were used for different experiments. Abdominal skin was obtained from males and females ranging in age from 30-70 years old. The large age range and sex differences may influence results. Skin integrity was confirmed by TEWL and only skin meeting inclusion criteria (≤ 13 gm-2h-1) was included in the analysis. This is a slight deviation from guidelines which suggest viable skin has a TEWL reading of less than 10 grams/m ² /hour.
	Metric 18:	Confounding variables in outcomes unrelated to exposure	High	There were no reported differences among the study replicates that were unrelated to exposure; the test substance was demonstrated to be soluble in the receptor fluid.
Domain 7: Data Presentation and Analysis				
	Metric 19:	Data analysis	Uninformative	Means and standard deviations were calculated and reported. The CV for total absorbed dose is >25% for all dose groups/conditions. The high CV values make the study uninformative.
	Metric 20:	Data interpretation	High	The study reported doses in receptor fluid, receptor chamber, exposed skin, in stratum corneum, dislodgeable dose, donor chamber, in trap, and in tape strips. Data were interpreted correctly. Decreased recovery in some groups was most likely due to the volatility of the test substance, although total recoveries were >80%.
	Metric 21:	Reporting of data	High	Data for all relevant endpoints were reported quantitatively as means \pm SD. Individual cell data are shown.
Overall Quality Determination		Uninformative		

Study Citation: Labcorp Early Development, (2024). 1,1-Dichloroethane: Rates of penetration through human skin using a flow through in vitro system.
HERO ID: 11396332

EXTRACTION

Parameter	Data
Extraction ID; Chemical:	Finite; 1% in 1,2-dichloroethane; 1,1-Dichloroethane-Parent compound
Skin Material/Species; Skin Preparation; Skin Thickness (um); Diffusion Cell Exposure Setup Type:	ex vivo human; Split thickness; 400; Flow-through; Notes: Not Reported
Occlusion Type; Donor Chamber Vehicle; Concentration of Test Substance in Vehicle (enter as percent):	Semi-occluded (e.g., vapor trap); 1,2-dichloroethane; 1
Mass per Surface Area on Skin (mg/cm ²); Duration of Test Substance on Skin:	0.14; 8 hrs; Not Reported
Duration of Absorbance Measured; Frequency of Samples:	24 hrs; Samples were collected at 10 minutes, 30 minutes, 1 hours, 2 hours, and then 2-hourly intervals; Notes: Not Reported
Time Skin was Washed and Method used; Radiolabel Presence:	"The surface of the membrane was rinsed three times (3 x 0.5 mL) with a mild solution of Dove™ soap in water (ca. 1%). The skin surface was swabbed twice with cotton buds soaked with the liquid soap solution. Finally, the membrane was rinsed with a small volume (0.5 mL) of water and a further cotton bud used to swab the surface of the membrane until dry."; Yes
Total Recovery (percent); Dose Type:	55.42; Finite
Percent Found in Skin Depot After Washing and Tape Stripping; Comments:	0; Notes: not detected
Percent Found in All Tape Strips, Excluding the Upper Two Strips; Comments:	0; Notes: not detected
Percent Found in Receptor Fluid and Receptor Fluid Rinse; Comments:	0.05; Notes: CV= 120%
Total Percent Absorbed:	0
Steady State Permeability Coefficient (Kp) (cm/hr); Steady State Permeability Coefficient (Comments); Steady State Flux (ug/cm ² /hr); Steady State Flux (Comments); Maximum Permeability Coefficient (Kp) (cm/hr); Maximum Permeability Coefficient (Comments); Maximum Flux (ug/cm ² /hr); Maximum Flux (Comments):	Not Reported; Notes: Not Reported; Not Reported; Notes: Not Reported; Not Reported; Notes: Not Reported; Not Reported; Notes: Not Reported

EVALUATION

Domain	Metric	Rating	Comments
Domain 1: Test Substance			
	Metric 1:	High	Test substances were identified as non-radiolabeled 1,1-dichloroethane and radiolabeled 1,1-14C dichloroethane, CAS number: 75-34-3; the structure was reported for the radiolabeled substance noting the location of the radiolabel within the substance.
	Metric 2:	High	The non-radiolabeled, 1,1-dichloroethane was sourced from LGC standards; the radiolabeled 1,1-dichloroethane was sourced from Moravek Biochemical Inc. The study included certificates of analyses that included lot/batch numbers.
	Metric 3:	High	The radiochemical purity of the radiolabeled test substance was reported to be ≥98% by GC-MS. The purity of the non-radiolabeled substance was 99.48% (batch 1) and 99.9% (batch 2).

Continued on next page ...

... continued from previous page

Study Citation:		Labcorp Early Development, (2024). 1,1-Dichloroethane: Rates of penetration through human skin using a flow through in vitro system.		
HERO ID:		11396332		
Domain	Metric	EVALUATION Rating	Comments	
Domain 2: Test Design	Metric 4:	Reference compounds	Medium	Testosterone (non-labelled and radiolabeled) was used as a reference compound in accordance with OECD 28 guidelines. 1 mg/ml was administered to the skin (0.011 mg/cm ²). Study authors report 92.1% recovery (CV = 2.57%). The CV values for some measurements were <25% (total dislodgeable dose, total unabsorbed dose, and the amount in skin washes), while others were >25% (i.e., total absorbed dose, dose in the skin, dose on surface tape and donor chambers etc.). It is unclear if these high CV values suggest any issues with the reliability of the tests conducted. However, this lab also cites previous publications that demonstrate the labs proficiency in performing dermal absorption studies with lipophilic compounds (Van de Sandt et al. 2004).
	Metric 5:	Assay procedures	Medium	This study was conducted according to OECD TG 428, OECD 28, and OECD 156. The assay procedures specified in the report were described in detail, although some information was missing. The flow through diffusion set-up was sufficiently reported including a schematic drawing. The study does not report if there was an equilibration period after the skin was placed into the chamber (guidelines recommend a 30-minute equilibrium period). Finite doses of 1,1-dichloroethane (neat, 50%, 10%, and 1% dissolved in either isopropyl myristate or 1,2-dichloroethane) were applied to human skin (at least 6 replicates; surface area of 0.64 cm ²). The volume applied to the skin surface was 10uL/cm ² . The cells were semi-occluded by a charcoal filter in the neck of the donor cell. The thickness of the skin samples ranged from 200 to 400 um, exact thicknesses were not reported. The receptor solution was water fortified with 6% polyoxyethelene 20 oleyl ether. OECD 156 guidelines recommend <6% polyoxyethelene (20) oleyl ether in water for lipophilic compounds. The solubility of 1,1-dichloroethane was tested in the receptor fluid prior to the study and determined dissolution of 1,1-dichloroethane was not rate-limiting. The skin membranes were maintained at 32 degrees C; humidity ranged from 30-70%. The flow rate of receptor fluid was 1.5 mL/hour. After 8 hours skin was washed. "The surface of the membrane was rinsed three times (3 x 0.5 mL) with a mild solution of Dove™ soap in water (ca. 1%). The skin surface was swabbed twice with cotton buds soaked with the liquid soap solution. Finally, the membrane was rinsed with a small volume (0.5 mL) of water and a further cotton bud was used to swab the surface of the membrane until dry". For some groups, headspace above the skin was drawn into a syringe for analysis after 4 hours. Receptor fluid samples (volume not reported) were collected at 10 minutes, 30 minutes, 1 hour, 2 hours and every 2 hours henceforth for 24 hours (16 samples) and analyzed for radioactivity. After 24 hours, the skin was tap-stripped 5 times. "Radioactivity in gross amounts less than twice the background level was considered to be below the limit of accurate determination."
	Metric 6:	Standards for tests	Medium	The skin samples (abdomen) were obtained from 24 female and 2 male human donors following elective surgery. Samples were stored at -20oC until used. The integrity of the skin was determined by stable trans epidermal water loss (TEWL) prior to dose application and at termination (prior to skin washing and tape stripping). A TEWL of ≤ 13 gm-2h-1 was considered acceptable. This is a slight deviation from guidelines which suggest viable skin has a TEWL reading of less than 10 grams/m2/hour. TEWL readings were reported. Percent recovery was >80%, which is acceptable for this volatile substance. Coefficients of variation (CV) values were reported. CVs were all > 25%. Further discussion of the CVs is provided in Metric 19.

Continued on next page ...

... continued from previous page

Study Citation:		Labcorp Early Development, (2024). 1,1-Dichloroethane: Rates of penetration through human skin using a flow through in vitro system.		
HERO ID:		11396332		
Domain	Metric	EVALUATION Rating	Comments	
Domain 3: Exposure Characterization				
	Metric 7:	Preparation and storage of test substance (chemical)	Medium	The preparation of the test substance was adequately reported. Dilutions were magnetically stirred if deemed necessary by study authors (criteria not reported). The activity and homogeneity of the diluted doses were assessed from the top, middle, and bottom of the solution while being stirred. All doses were considered homogenous (data not shown). Storage conditions are not reported; it is unclear how far in advanced solutions were made, and how they were stored. However, the study included only a single application, and dosing was confirmed by analysis of QC samples taken during the dosing of the cells, therefore, the lack of storage information is not likely to have an impact on the study results. The solubility of the test substance in the receptor fluid was tested, (895 ug/mL) and was appropriate (>15-fold higher than observed absorption) and was considered non-rate-limiting.
	Metric 8:	Consistency of exposure administration	Low	The volume of test substance applied to the skin was 10 uL/cm2 exposed skin (6.4 uL dose), which is in agreement with OECD guidelines. The skin surface area of 0.64 cm2 was consistent across groups. The skin thickness was reported as a range (200 to 400 uM). It is unclear if the variation in thicknesses was consistent across groups, and this may have contributed to some of the endpoint variations (and subsequently high CVs) observed.
	Metric 9:	Reporting of concentrations	High	The applied dose is reported as dpm, mg per cell, and mg/cm2. Individual cells are reported independently. Nominal and analytical doses are reported.
	Metric 10:	Exposure frequency	High	Exposure duration was reported and appropriate for determining absorption. Test substance was in contact with the skin for 8 hours prior to washing. Samples of receptor fluid were collected for a total of 24 hours.
	Metric 11:	Number of exposure groups and concentration spacing	High	There were 4 dose groups tested in a wide range of concentrations (neat to 1%). Dilutions of test substance was performed in two different vehicles (isopropyl myristate or 1,2-dichloroethane). Study authors state "The dose levels selected in this study were selected to be representative of potential in-life exposure levels".
Domain 4: Test Model				
	Metric 12:	Test model (skin)	High	The test model and descriptive information were reported. Samples of full thickness skin samples were obtained from males and females following elective surgery. Abdominal skin samples from 26 people (race not reported) ranging in age from 30-70 years old were studied. The samples were stored frozen at -20 degrees C +/- 10 degrees C. Prior to use, the skin samples were thawed, wiped to remove residual fat and blood, re-hydrated in purified water, and dermatomed using a mini-dermatome. The thickness ranged from 200 to 400 uM. The skin contained epidermis and some dermis. The OECD guidelines split thickness (dermatomed) skin is preferred. Membrane integrity was determined by measuring trans epidermal water loss prior to/and upon completion of the experiment.
	Metric 13:	Number/Replicates per group	Medium	The number of replicates was appropriate as per OECD 428. The study used 6 replicates to ensure a minimum of 4 intact replicates could be obtained if skin integrity was not maintained.
Domain 5: Outcome Assessment				
Continued on next page ...				

... continued from previous page

Study Citation: HERO ID:		Labcorp Early Development, (2024). 1,1-Dichloroethane: Rates of penetration through human skin using a flow through in vitro system. 11396332		
		EVALUATION		
Domain	Metric		Rating	Comments
	Metric 14:	Outcome assessment methodology	High	The outcome assessment methodology addressed the intended outcome of interest and was sensitive for the outcome. The test followed OECD guidelines 428 and 28. 10 uL/cm ² of test substance was applied to skin samples. Measurement techniques and timing were reported and appropriate. A finite dose was used to determine absorption.
	Metric 15:	Consistency of outcome assessment	High	Details of outcome assessment protocol were reported and outcomes were assessed consistently across study replicates. The same duration of exposure, receptor fluid used, and sampling period were consistent across replicates.
	Metric 16:	Sampling adequacy and sensitivity	Medium	The study reported adequate sampling for the outcomes of interest; measurement sensitivity was sufficient. The sampling intervals were adequate (16 over 24 hours). Methods for the determination of radioactivity are reported. "Radioactivity in gross amounts less than twice the background level was considered to be below the limit of accurate determination." Scintillation counts were not shown. A graphical representation of absorption over time is shown.
Domain 6: Confounding/Variable Control				
	Metric 17:	Confounding variables in test design and procedures	Medium	Two batches of radiolabeled and non-radiolabeled 1,1-dichloroethane were used; it is unclear if batches were combined or if different batches were used for different experiments. Abdominal skin was obtained from males and females ranging in age from 30-70 years old. The large age range and sex differences may influence results. Skin integrity was confirmed by TEWL and only skin meeting inclusion criteria (≤ 13 gm-2h-1) was included in the analysis. This is a slight deviation from guidelines which suggest viable skin has a TEWL reading of less than 10 grams/m ² /hour.
	Metric 18:	Confounding variables in outcomes unrelated to exposure	High	There were no reported differences among the study replicates that were unrelated to exposure; the test substance was demonstrated to be soluble in the receptor fluid.
Domain 7: Data Presentation and Analysis				
	Metric 19:	Data analysis	Uninformative	Means and standard deviations were calculated and reported. The CV for total absorbed dose is >25% for all dose groups/conditions. The high CV values make the study uninformative.
	Metric 20:	Data interpretation	High	The study reported doses in receptor fluid, receptor chamber, exposed skin, in stratum corneum, dislodgeable dose, donor chamber, in trap, and in tape strips. Data were interpreted correctly. Decreased recovery in some groups was most likely due to the volatility of the test substance, although total recoveries were >80%.
	Metric 21:	Reporting of data	High	Data for all relevant endpoints were reported quantitatively as means \pm SD. Individual cell data are shown.
Overall Quality Determination		Uninformative		

Study Citation: Labcorp Early Development, (2024). 1,1-Dichloroethane: Rates of penetration through human skin using a flow through in vitro system.
HERO ID: 11396332

EXTRACTION

Parameter	Data
Extraction ID; Chemical:	Finite; 10% in IPM; 1,1-Dichloroethane-Parent compound
Skin Material/Species; Skin Preparation; Skin Thickness (um); Diffusion Cell Exposure Setup Type:	ex vivo human; Split thickness; 400; Flow-through; Notes: Not Reported
Occlusion Type; Donor Chamber Vehicle; Concentration of Test Substance in Vehicle (enter as percent):	Semi-occluded (e.g., vapor trap); isopropyl myristate; 10
Mass per Surface Area on Skin (mg/cm ²); Duration of Test Substance on Skin:	1.04; 8 hrs; Not Reported
Duration of Absorbance Measured; Frequency of Samples:	24 hrs; Samples were collected at 10 minutes, 30 minutes, 1 hours, 2 hours, and then 2-hourly intervals; Notes: Not Reported
Time Skin was Washed and Method used; Radiolabel Presence:	”The surface of the membrane was rinsed three times (3 x 0.5 mL) with a mild solution of Dove™ soap in water (ca. 1%). The skin surface was swabbed twice with cotton buds soaked with the liquid soap solution. Finally, the membrane was rinsed with a small volume (0.5 mL) of water and a further cotton bud used to swab the surface of the membrane until dry.”; Yes
Total Recovery (percent); Dose Type:	87.68; Finite
Percent Found in Skin Depot After Washing and Tape Stripping; Comments:	0; Notes: not detected
Percent Found in All Tape Strips, Excluding the Upper Two Strips; Comments:	0; Notes: not detected
Percent Found in Receptor Fluid and Receptor Fluid Rinse; Comments:	0; Notes: <0.01%. Cell number 52 was not included in calculation due decreased membrane integrity (determined by TWEL measurements).
Total Percent Absorbed:	0
Steady State Permeability Coefficient (Kp) (cm/hr); Steady State Permeability Coefficient (Comments); Steady State Flux (ug/cm ² /hr); Steady State Flux (Comments); Maxium Permeability Coefficient (Kp) (cm/hr); Maxium Permeability Coefficient (Comments); Maximum Flux (ug/cm ² /hr); Maximum Flux (Comments):	Not Reported; Notes: Not Reported; Not Reported; Notes: Not Reported; Not Reported; Notes: Not Reported; Not Reported; Notes: Not Reported

EVALUATION

Domain	Metric	Rating	Comments
Domain 1: Test Substance			
	Metric 1: Test substance identity	High	Test substances were identified as non-radiolabeled 1,1-dichloroethane and radiolabeled 1,1-14C dichloroethane, CAS number: 75-34-3; the structure was reported for the radiolabeled substance noting the location of the radiolabel within the substance.
	Metric 2: Test substance source	High	The non-radiolabeled, 1,1-dichloroethane was sourced from LGC standards; the radiolabeled 1,1-dichloroethane was sourced from Moravex Biochemical Inc. The study included certificates of analyses that included lot/batch numbers.
	Metric 3: Test substance purity	High	The radiochemical purity of the radiolabeled test substance was reported to be ≥98% by GC-MS. The purity of the non-radiolabeled substance was 99.48% (batch 1) and 99.9% (batch 2).

Continued on next page ...

... continued from previous page

Study Citation: Labcorp Early Development, (2024). 1,1-Dichloroethane: Rates of penetration through human skin using a flow through in vitro system.
HERO ID: 11396332

		EVALUATION	
Domain	Metric	Rating	Comments
Domain 2: Test Design	Metric 4:	Reference compounds	Medium Testosterone (non-labelled and radiolabeled) was used as a reference compound in accordance with OECD 28 guidelines. 1 mg/ml was administered to the skin (0.011 mg/cm ²). Study authors report 92.1% recovery (CV = 2.57%). The CV values for some measurements were <25% (total dislodgeable dose, total unabsorbed dose, and the amount in skin washes), while others were >25% (i.e., total absorbed dose, dose in the skin, dose on surface tape and donor chambers etc.). It is unclear if these high CV values suggest any issues with the reliability of the tests conducted. However, this lab also cites previous publications that demonstrate the labs proficiency in performing dermal absorption studies with lipophilic compounds (Van de Sandt et al. 2004).
	Metric 5:	Assay procedures	Medium This study was conducted according to OECD TG 428, OECD 28, and OECD 156. The assay procedures specified in the report were described in detail, although some information was missing. The flow through diffusion set up was sufficiently reported including a schematic drawing. The study does not report if there was an equilibration period after the skin was placed into the chamber (guidelines recommend a 30-minute equilibrium period). Finite doses of 1,1-dichloroethane (neat, 50%, 10%, and 1% dissolved in either isopropyl myristate or 1,2-dichloroethane) were applied to human skin (at least 6 replicates; surface area of 0.64 cm ²). Volume applied (10uL/cm ²) was applied to skin surface. The cells were semi-occluded by a charcoal filter in the neck of the donor cell. The thickness of the skin samples ranged from 200 to 400 um, exact thicknesses were not reported. The receptor solution was water fortified with 6% polyoxyethelene 20 oleyl ether. OECD 156 guidelines recommend <6% polyoxyethelene (20) oleyl ether in water for lipophilic compounds. The solubility of 1,1-dichloroethane was tested in the receptor fluid prior to the study and determined dissolution of 1,1-dichloroethane was not rate-limiting. The skin membranes were maintained at 32 degrees C; humidity ranged from 30-70%. The flow rate of receptor fluid was 1.5 mL/hour. After 8 hours skin was washed. "The surface of the membrane was rinsed three times (3 x 0.5 mL) with a mild solution of Dove™ soap in water (ca. 1%). The skin surface was swabbed twice with cotton buds soaked with the liquid soap solution. Finally, the membrane was rinsed with a small volume (0.5 mL) of water and a further cotton bud was used to swab the surface of the membrane until dry". For some groups, headspace above the skin was drawn into a syringe for analysis after 4 hours. Receptor fluid samples (volume not reported) were collected at 10 minutes, 30 minutes, 1 hour, 2 hours and every 2 hours henceforth for 24 hours (16 samples) and analyzed for radioactivity. After 24 hours, the skin was tape-stripped 5 times. "Radioactivity in gross amounts less than twice the background level was considered to be below the limit of accurate determination."
	Metric 6:	Standards for tests	Medium The skin samples (abdomen) were obtained from 24 female and 2 male human donors following elective surgery. Samples were stored at -20oC until used. The integrity of the skin was determined by stable trans epidermal water loss (TEWL) prior to dose application and at termination (prior to skin washing and tape stripping). A TEWL of ≤ 13 gm-1h-1 was considered acceptable. This is a slight deviation from guidelines which suggest viable skin has a TEWL reading of less than 10 grams/m ² /hour. TEWL readings were reported. Percent recovery was >80%, which is acceptable for this volatile substance. Coefficients of variation (CV) values were reported. CVs were all > 25%. Further discussion of the CVs is provided in Metric 19.

Continued on next page ...

... continued from previous page

Study Citation: Labcorp Early Development, (2024). 1,1-Dichloroethane: Rates of penetration through human skin using a flow through in vitro system.
HERO ID: 11396332

		EVALUATION		
Domain	Metric	Rating		Comments
Domain 3: Exposure Characterization				
	Metric 7:	Preparation and storage of test substance (chemical)	Medium	The preparation of the test substance was adequately reported. Dilutions were magnetically stirred if deemed necessary by study authors (criteria not reported). The activity and homogeneity of the diluted doses were assessed from the top, middle, and bottom of the solution while being stirred. All doses were considered homogenous (data not shown). Storage conditions were not reported; it is unclear how far in advanced solutions were made, and how they were stored. However, the study included only a single application, and dosing was confirmed by analysis of QC samples taken during the dosing of the cells, therefore, the lack of storage information is not likely to have an impact on the study results. The solubility of the test substance in the receptor fluid was tested, (895 ug/mL) and was appropriate (>15-fold higher than observed absorption); solubility was considered non rate-limiting.
	Metric 8:	Consistency of exposure administration	Low	The volume of test substance applied to the skin was 10 uL/cm ² exposed skin (6.4 uL dose), which is in agreement with OECD guidelines. The skin surface area of 0.64 cm ² was consistent across groups. The skin thickness was reported as a range (200 to 400 uM). It is unclear if the variation in thicknesses was consistent across groups, and this may have contributed to some of the endpoint variations (and subsequently high CVs) observed.
	Metric 9:	Reporting of concentrations	High	The applied dose is reported as dpm, mg per cell, and mg/cm ² . Individual cells are reported independently. Nominal and analytical doses are reported.
	Metric 10:	Exposure frequency	High	Exposure duration was reported and appropriate for determining absorption. Test substance was in contact with the skin for 8 hours prior to washing. Samples of receptor fluid were collected for a total of 24 hours.
	Metric 11:	Number of exposure groups and concentration spacing	High	There were 4 dose groups tested in a wide range of concentrations (neat to 1%). Dilutions of test substance was performed in two different vehicles (isopropyl myristate or 1,2-dichloroethane). Study authors state "The dose levels selected in this study were selected to be representative of potential in-life exposure levels".
Domain 4: Test Model				
	Metric 12:	Test model (skin)	High	The test model and descriptive information were reported. Samples of full-thickness skin samples were obtained from males and females following elective surgery. Abdominal skin samples from 27 people (race not reported) ranging in age from 30-70 years old were studied. The samples were stored frozen at -20 degrees C +/-10 degrees C. Prior to use, the skin samples were thawed, wiped to remove residual fat and blood, re-hydrated in purified water, and dermatomed using a mini-dermatome. The thickness ranged from 200 to 400 uM. The skin contained epidermis and some dermis. The OECD guidelines split thickness (dermatomed) skin is preferred. Membrane integrity was determined by measuring transepidermal water loss prior to/and upon completion of the experiment.
	Metric 13:	Number/Replicates per group	Medium	The number of replicates was appropriate as per OECD 428. The study used 6 replicates in order to ensure a minimum of 4 intact replicates could be obtained if skin integrity was not maintained.
Domain 5: Outcome Assessment				

Continued on next page ...

... continued from previous page

Study Citation:		Labcorp Early Development, (2024). 1,1-Dichloroethane: Rates of penetration through human skin using a flow through in vitro system.		
HERO ID:		11396332		
		EVALUATION		
Domain	Metric	Rating	Comments	
	Metric 14:	Outcome assessment methodology	High	The outcome assessment methodology addressed the intended outcome of interest and was sensitive for the outcome. The test followed OECD guidelines 428 and 28. 10 uL/cm ² of the test substance was applied to skin samples. Measurement techniques and timing were reported and appropriate. A finite dose was used to determine absorption.
	Metric 15:	Consistency of outcome assessment	High	Details of outcome assessment protocol were reported and outcomes were assessed consistently across study replicates. The same duration of exposure, receptor fluid used, and sampling period was consistent across replicates.
	Metric 16:	Sampling adequacy and sensitivity	Medium	The study reported adequate sampling for the outcomes of interest; measurement sensitivity was sufficient. The sampling intervals were adequate (16 over 24 hours). Methods for the determination of radioactivity are reported. "Radioactivity in gross amounts less than twice the background level was considered to be below the limit of accurate determination." Scintillation counts were not shown. Graphical representation of absorption over time is shown.
Domain 6: Confounding/Variable Control				
	Metric 17:	Confounding variables in test design and procedures	Medium	Two batches of radiolabeled and non-radiolabeled 1,1-dichloroethane were used; it is unclear if batches were combined or if different batches were used for different experiments. Abdominal skin was obtained from males and females ranging in age from 30-70 years old. The large age range and sex differences may influence results. Skin integrity was confirmed by TEWL and only skin meeting inclusion criteria (≤ 13 gm-2h-1) were included in the analysis. This is a slight deviation from guidelines which suggest viable skin has a TEWL reading of less than 10 grams/m ² /hour.
	Metric 18:	Confounding variables in outcomes unrelated to exposure	High	There were no reported differences among the study replicates that were unrelated to exposure; the test substance was demonstrated to be soluble in the receptor fluid.
Domain 7: Data Presentation and Analysis				
	Metric 19:	Data analysis	High	Means and standard deviations were calculated and reported. <0.01% of the applied dose was absorbed. The CV values for total recovery, total unabsorbed, and dose in the trap extract were all 3.84%.
	Metric 20:	Data interpretation	High	The study reported dose in receptor fluid, receptor chamber, exposed skin, in stratum corneum, dislodgeable dose, donor chamber, in trap, and in tape strips. Data were interpreted correctly. Decreased recovery (87%) was most likely due to the volatility of the test substance.
	Metric 21:	Reporting of data	High	Data for all relevant endpoints were reported quantitatively as means \pm SD. Individual cell data are shown.
Overall Quality Determination		High		

Study Citation: DuPont Haskell Lab, (2005). Ethylene dichloride: In vitro dermal absorption rate testing.
HERO ID: 7490381

EXTRACTION

Parameter	Data
Extraction ID; Chemical:	Infinite dosing; 1,1-Dichloroethane-Parent compound
Skin Material/Species; Skin Preparation; Skin Thickness (um); Diffusion Cell Exposure Setup Type:	ex vivo human; Not Reported; 50; Static; Notes: Not Reported
Occlusion Type; Donor Chamber Vehicle; Concentration of Test Substance in Vehicle (enter as percent):	Occluded; None described; Not Reported
Mass per Surface Area on Skin (mg/cm ²); Duration of Test Substance on Skin:	1,482; 8 hrs; Not Reported
Duration of Absorbance Measured; Frequency of Samples:	8 hrs; Receptor fluid samples were taken at 0.5, 1, 2, 3, 4, 5, 6, 7, and 8 hours post application.; Notes: Not Reported
Time Skin was Washed and Method used; Radiolabel Presence:	Skin was washed with a 2% soap solution followed by rinsing with water.; Yes
Total Recovery (percent); Dose Type:	85.9; Infinite
Percent Found in Skin Depot After Washing and Tape Stripping; Comments:	0.01; Notes: <0.01% was recovered in the skin after washing.
Percent Found in All Tape Strips, Excluding the Upper Two Strips; Comments:	0.01; Notes: <0.01% was recovered in the skin after washing.
Percent Found in Receptor Fluid and Receptor Fluid Rinse; Comments:	0.6; Notes: 0.60 ± 0.21% (CV = 35%) was recovered in the receptor fluid.
Total Percent Absorbed:	1
Steady State Permeability Coefficient (Kp) (cm/hr); Steady State Permeability Coefficient (Comments); Steady State Flux (ug/cm ² /hr); Steady State Flux (Comments); Maximum Permeability Coefficient (Kp) (cm/hr); Maximum Permeability Coefficient (Comments); Maximum Flux (ug/cm ² /hr); Maximum Flux (Comments):	Not Reported; Notes: Not Reported; 0.000969; Notes: 9.69 x 10 ⁻⁴ ± 2.84 x 10 ⁻⁴ (CV = 29%); 1197.1; Notes: 1197.1 ± 351.1 (CV = 29%); Not Reported; Notes: Not Reported

EVALUATION

Domain	Metric	Rating	Comments	
Domain 1: Test Substance				
	Metric 1:	Test substance identity	High	Test substances were identified as non-radiolabeled 1,1-Dichloroethane and radiolabeled 1,2-14C EDC, CAS number: 107-06-2; the structure was reported for the radiolabeled substance noting the location of the radiolabel within the substance.
	Metric 2:	Test substance source	High	The non-radiolabeled, high-grade technical EDC was sourced from Occidental Chemical Corporation; the radiolabeled EDC was sourced from Perkin-Elmer Life Sciences. The study included certificates of analyses that included lot/batch numbers and verification by GC.

Continued on next page ...

... continued from previous page

Study Citation:		DuPont Haskell Lab, (2005). Ethylene dichloride: In vitro dermal absorption rate testing.		
HERO ID:		7490381		
		EVALUATION		
Domain	Metric		Rating	Comments
	Metric 3:	Test substance purity	High	The radiochemical purity of the radiolabeled test substance was 98.1% by HPLC; when mixed with technical EDC, radiochemical purity was > 98%. The purity of the non-radiolabeled substance was 99.94%. Impurity analysis was done via GC.
Domain 2: Test Design				
	Metric 4:	Reference compounds	Low	No reference compound was used and no history of test performance in the laboratory was reported.
	Metric 5:	Assay procedures	Medium	This study was conducted according to OECD TG 428 and OECD 28. The assay procedures specified in the report were described in detail, although some information was missing. After a 15 min equilibration period, an infinite dose of EDC (1,235 mg/cm ²) (based on a dose volume of 1200 uL/cm ² and a density of 1.2351g/mL) was applied to human skin (6 replicates; surface area of 0.64 cm ² , epidermal layer only, thickness ranged from 50 to 98 um) in a static diffusion cell model under occluded conditions (rubber stopper) for 8 hours. The specific activity of the applied substance was 0.02686uCi/mg. Receptor fluid (0.9% saline fortified with 6% polyethoxyoleate; 5 mL volume) was maintained at 32 degrees C; humidity was not reported. It was not specified whether the fluid was continuously stirred as per OECD 428 guidelines. Receptor fluid samples (50ul aliquots) were collected at 0.5, 1, 2, 3, 4, 5, 6, 7, and 8 hours post-application and analyzed for radioactivity. A steady state was reached. Tape stripping was not conducted, but this can be difficult with epidermal membranes. Radioactivity was measured in receptor fluid aliquots, in the excess dose removed at the end of exposure, in the stopper extract, the skin wash/rinse, the skin post-washing, and in the donor chamber rinse. The time and number of scintillation counts were reported. The scintillation time was 10 minutes for one sample and 5 minutes for all other samples, which the study authors deemed adequate. "The limit of detection (LOD) for the analysis of each sample was taken as twice the background disintegration rate obtained from the analysis of appropriate blank samples."
	Metric 6:	Standards for tests	Medium	The appropriate tests for quality control were conducted, and results were reported. The skin was allowed to equilibrate for 30 minutes before use. The integrity of the epidermal membranes was then tested prior to the start of the study via electrical impedance, and only membranes with an EI of $\geq 17 \text{ k}\Omega$ were used in the study. Skin integrity was also assessed via electrical impedance at the end of the experiment. A decrease in integrity from the values measured prior to the start of the study was noted. The percent recovery of 85.9% was reported and is acceptable for volatile substances as per OECD TG 28 (100 \pm 20%). However, recovery determination is not generally relevant for studies only determining a Kp. Coefficients of variation (CV) values were not reported but could be determined based on the SD relative to the mean. Except for data reporting total recovery, the CVs were all > 25%. Further discussion of the CVs is provided in Metric 19.
Domain 3: Exposure Characterization				
Continued on next page ...				

... continued from previous page

Study Citation:		DuPont Haskell Lab, (2005). Ethylene dichloride: In vitro dermal absorption rate testing.		
HERO ID:		7490381		
EVALUATION				
Domain	Metric	Rating	Comments	
	Metric 7:	Preparation and storage of test substance (chemical)	Medium	Information on preparation and storage was not thoroughly reported. The non-radiolabeled test substance was spiked with the radiolabeled test substance. The homogeneity and amount of radiolabeled EDC in the formulation was determined by radioanalysis via liquid scintillation counting. The concentration of EDC in the prepared solution was taken at its density, 1.2351 g/mL, and the homogeneity analysis was used to calculate the specific activity of the formulation. No further dilutions or use of a vehicle were described. No information on storage was provided, but it was indicated at the beginning of the report that the test substance was stable under the conditions of the study. The solubility of the test substance in the receptor fluid was tested, (9,690 ug/mL) and was appropriate. The authors also took measures during exposure to account for the volatility of the test substance (use of occluded conditions and a carbon trap (for short-term exposures reported in a separate form).
	Metric 8:	Consistency of exposure administration	Low	The study used the same volume (1,200 uL/cm ²) across all samples and the skin surface area of 0.64 cm ² was consistent. However, the skin thickness was reported as a range (50 to 98 uM). It is unclear of the variation in thicknesses was consistent across groups, and this may have contributed to some of the endpoint variations (and subsequently high CVs) observed.
	Metric 9:	Reporting of concentrations	High	The target rate was 1,200 uL/cm ² . For infinite dose experiments concentrations of 100 uL/cm ² or greater are appropriate as per OECD 28. "The actual dose applied to each replicate was determined by subjecting aliquots of the prepared solution to liquid scintillation counting. The total amount of EDC applied to the skin was determined by the total radioactivity applied and the verified specific activity." The verified specific activity for the prepared solution was 0.02686 uCi/mg, and the concentration in the applied formulation was taken at its density (1,235,1000 ug/cm ³)." With the information provided, the actual applied concentration in mg/cm ² can be calculated.
	Metric 10:	Exposure frequency	High	Exposure duration was reported and appropriate for Kp determination, an 8-hour exposure was used and a steady state was reached. The shorter duration was appropriate for a volatile test substance.
	Metric 11:	Number of exposure groups and concentration spacing	Low	Fewer than three concentrations were tested. This study only tested one concentration.
Domain 4: Test Model	Metric 12:	Test model (skin)	Low	The test model and descriptive information were reported. Samples of full thickness human cadaver skin were sourced from the National Disease Research Interchange. Abdominal skin samples from 3 Caucasian females were used. The samples were stored frozen at -20 degrees C. It was noted that the samples were removed from the donors within 24 hours of death, and were used within three months. Prior to use, the skin samples were thawed, soaked in water, and the epidermal layer was isolated then stored refrigerated (max duration not specified) until use. The thickness ranged from 50 to 98 uM. The OECD TG 156 specifies that use of epidermal membrane may, in some cases, overestimate human in vivo skin absorption due to the lack of a sufficient barrier function and split thickness (dermatomed) skin is preferred.
	Metric 13:	Number/Replicates per group	Medium	The study used 6 replicates (from 3 donors). The number of replicates was appropriate as per OECD 428.
Domain 5: Outcome Assessment				

Continued on next page ...

... continued from previous page

Study Citation:		DuPont Haskell Lab, (2005). Ethylene dichloride: In vitro dermal absorption rate testing.		
HERO ID:		7490381		
		EVALUATION		
Domain	Metric	Rating	Comments	
	Metric 14:	Outcome assessment methodology	High	The outcome assessment methodology addressed the intended outcome of interest and was sensitive for the outcome. The test followed OECD guidelines 428 and 28; FR volume 69, number 80; and European Commission Guidance Document on Dermal Absorption. Measurement techniques and timing were reported and appropriate. An infinite dose of the test substance was used to determine the Kp. A dose volume of 1200 uL/cm ² was used.
	Metric 15:	Consistency of outcome assessment	High	Details of outcome assessment protocol were reported and outcomes were assessed consistently across study replicates. The same duration of exposure, receptor fluid used, and sampling period was consistent across replicates.
	Metric 16:	Sampling adequacy and sensitivity	High	The study reported adequate sampling for the outcomes of interest; measurement sensitivity was sufficient. The sampling intervals were adequate to allow at least 4 data points at a steady state. The time and number of scintillation counts were reported for one of the receptor fluid samples, 10 minutes (160,000 disintegration counts). The other samples were counted for 5 minutes, which the authors deemed adequate. Analysis of a mock dose showed an appropriate signal-to-noise ratio.
Domain 6: Confounding/Variable Control				
	Metric 17:	Confounding variables in test design and procedures	Medium	There were no differences reported among study groups (test substance lot, number of skin samples, or quality of tissues) that could influence the outcome assessment. Skin integrity was confirmed by EI and only membranes with an EI of ≥ 17 k Ω were retained for use in the study. The study did indicate that, relative to pre-exposure EI results, there was a decrease in EI after exposure (the ratio of post-EI values to pre-EI values was 0.59). The study authors indicated that the decrease did not affect the results of the experiment. Scintillation counts were done for 5 minutes for most samples. The authors indicated that this deviated from the protocol, but that it had no impact on the interpretation of the study results.
	Metric 18:	Confounding variables in outcomes unrelated to exposure	High	There were no reported differences among the study replicates that were unrelated to exposure; the test substance was demonstrated to be soluble in the receptor fluid.
Domain 7: Data Presentation and Analysis				
	Metric 19:	Data analysis	Uninformative	The Kp was determined by first plotting the cumulative amount of ug equivalents in the receptor compartment at each serial collection time-point, adjusted for total receptor fluid volume, against time (hrs) to produce an absorption profile. The Kp was then calculated by dividing the penetration rate or slope of the line at steady-state, by the concentration of the applied chemical. The total recovery was determined and was the sum of the amount extracted from the rubber stopper, and what was detected in the receptor fluid, in skin washes, in the donor chamber, and in the skin. The permeability coefficient (Kp) was $9.69 \times 10^{-4} \pm 2.84 \times 10^{-4}$ (Mean \pm SD). The CV is $>25\%$. The CVs were also $>25\%$ for the cumulative amount penetrated (at each collection time point) and the steady-state penetration rate. The high CV values make the study uninformative.
	Metric 20:	Data interpretation	High	The Kp was derived from an appropriate exposure condition (infinite dose). Recovery determination is not relevant for infinite dose applications; however, the authors still reported total recovery, which fell within guideline specifications for a volatile test material (100% \pm 20%)
	Metric 21:	Reporting of data	High	Data for all relevant endpoints were reported quantitatively as means \pm SE.

Continued on next page ...

...continued from previous page

Study Citation: DuPont Haskell Lab, (2005). Ethylene dichloride: In vitro dermal absorption rate testing.
HERO ID: 7490381

EVALUATION

Domain

Metric

Rating

Comments

Overall Quality Determination

Uninformative

Study Citation: DuPont Haskell Lab, (2005). Ethylene dichloride: In vitro dermal absorption rate testing.
HERO ID: 7490381

EXTRACTION

Parameter	Data
Extraction ID; Chemical:	Finite dosing - 60 min; 1,1-Dichloroethane-Parent compound
Skin Material/Species; Skin Preparation; Skin Thickness (um); Diffusion Cell Exposure Setup Type:	ex vivo human; Not Reported; 50; Static; Notes: Not Reported
Occlusion Type; Donor Chamber Vehicle; Concentration of Test Substance in Vehicle (enter as percent):	Occluded; None described; Not Reported
Mass per Surface Area on Skin (mg/cm ²); Duration of Test Substance on Skin:	12.35; Other; 60 min
Duration of Absorbance Measured; Frequency of Samples:	Other; Collections were taken at a single time point 60 minutes after application.; Notes: 60 min
Time Skin was Washed and Method used; Radiolabel Presence:	Skin was washed with a 2% soap solution followed by rinsing with water.; Yes
Total Recovery (percent); Dose Type:	86.4; Finite
Percent Found in Skin Depot After Washing and Tape Stripping; Comments:	0.24; Notes: 0.24 ± 0.14% (CV = 58%) was recovered in the skin after washing.
Percent Found in All Tape Strips, Excluding the Upper Two Strips; Comments:	0.24; Notes: 0.24 ± 0.14% (CV = 58%) was recovered in the skin after washing.
Percent Found in Receptor Fluid and Receptor Fluid Rinse; Comments:	2.64; Notes: 2.64 ± 2.06% (CV = 78%) was recovered in the receptor fluid.
Total Percent Absorbed:	3
Steady State Permeability Coefficient (Kp) (cm/hr); Steady State Permeability Coefficient (Comments); Steady State Flux (ug/cm ² /hr); Steady State Flux (Comments); Maximum Permeability Coefficient (Kp) (cm/hr); Maximum Permeability Coefficient (Comments); Maximum Flux (ug/cm ² /hr); Maximum Flux (Comments):	Not Reported; Notes: Not Reported; Not Reported; Notes: Not Reported; Not Reported; Notes: Not Reported; 329.6; Notes: 329.6 ± 249.7 (CV = 76%)

EVALUATION

Domain	Metric	Rating	Comments
Domain 1: Test Substance			
	Metric 1:	High	Test substances were identified as non-radiolabeled 1,1-Dichloroethane and radiolabeled 1,2-14C EDC, CAS number: 107-06-2; the structure was reported for the radiolabeled substance noting the location of the radiolabel within the substance.
	Metric 2:	High	The non-radiolabeled, high-grade technical EDC was sourced from Occidental Chemical Corporation; the radiolabeled EDC was sourced from Perkin-Elmer Life Sciences. The study included certificates of analyses that included lot/batch numbers and verification by GC.
	Metric 3:	High	The radiochemical purity of the radiolabeled test substance was 98.1% by HPLC; when mixed with technical EDC, radiochemical purity was > 98%:. The purity of the non-radiolabeled substance was 99.94%. Impurity analysis was done via GC.

Continued on next page ...

... continued from previous page

Study Citation:		DuPont Haskell Lab, (2005). Ethylene dichloride: In vitro dermal absorption rate testing.		
HERO ID:		7490381		
		EVALUATION		
Domain	Metric	Rating	Comments	
Domain 2: Test Design	Metric 4:	Reference compounds	Low	No reference compound was used and no history of test performance in the laboratory was reported.
	Metric 5:	Assay procedures	Medium	This study was conducted according to OECD TG 428 and OECD 28. The assay procedures specified in the report were described in detail, although some information was missing. After a 15 min equilibration period, a finite dose of EDC (12.35 mg/cm ²) (based on a dose volume of 10 uL/cm ² and a density of 1.2351g/mL) was applied to human skin (12 replicates; surface area of 0.64 cm ² , epidermal layer only, thickness ranged from 50 to 98 um) in a static diffusion cell model under occluded conditions (charcoal trap). Absorption rates and recovery data were collected 10 minutes (6 replicates) and 60 minutes (6 replicates) post-dosing. During exposure, the receptor fluid (0.9% saline fortified with 6% polyethoxyoleate; 5 mL volume) was maintained at 32 degrees C; humidity was not reported. It was not specified whether the fluid was continuously stirred as per OECD 428 guidelines. Tape stripping was not conducted, but this can be difficult with epidermal membranes. Radioactivity was measured in receptor fluid aliquots, the skin wash/rinse, the skin post-washing, the donor chamber, and in the charcoal trap extraction. The time and number of scintillation counts were reported. The scintillation times (5 or 10 minutes) and counts were reported. "The limit of detection (LOD) for the analysis of each sample was taken as twice the background disintegration rate obtained from the analysis of appropriate blank samples."
	Metric 6:	Standards for tests	Medium	The appropriate tests for quality control were conducted, and results were reported. The skin was allowed to equilibrate for 30 minutes before use. The integrity of the epidermal membranes was then tested prior to the start of the study via electrical impedance, and only membranes with an EI of $\geq 17 \text{ k}\Omega$ were used in the study. Skin integrity was also assessed via electrical impedance at the end of the experiment and the ratios of the post-EI values to pre-EI values were 0.73 and 0.92 for the 10 min and 60 min exposures, respectively. The percent recoveries were 80.3% and 86.4%, for the 10 min and 60 min groups, respectively, and were acceptable for volatile substances as per OECD TG 28 (100 \pm 20%). Coefficients of variation (CV) values were not reported but could be determined based on the SD relative to the mean. For all endpoints (except total recovery), the CVs were $> 25\%$. Further discussion of the CVs is provided in Metric 19.
Domain 3: Exposure Characterization	Metric 7:	Preparation and storage of test substance (chemical)	Medium	Information on preparation and storage was not thoroughly reported. The non-radiolabeled test substance was spiked with the radiolabeled test substance. The homogeneity and amount of radiolabeled EDC in the formulation were determined by radioanalysis via liquid scintillation counting. The concentration of EDC in the prepared solution was taken at its density, 1.2351 g/mL, and the homogeneity analysis was used to calculate the specific activity of the formulation. No further dilutions or use of other vehicles were described. No information on storage was provided, but it was indicated at the beginning of the report that the test substance was stable under the conditions of the study. The solubility of the test substance in the receptor fluid was tested, (9,690 ug/mL) and was appropriate. The authors also took measures during exposure to account for the volatility of the test substance (use of occluded conditions and a carbon trap for short-term exposures).

Continued on next page ...

... continued from previous page

Study Citation: HERO ID:		DuPont Haskell Lab, (2005). Ethylene dichloride: In vitro dermal absorption rate testing. 7490381		
		EVALUATION		
Domain	Metric		Rating	Comments
	Metric 8:	Consistency of exposure administration	Low	The study used the same volume (10 uL/cm ²) across all samples and the skin surface area of 0.64 cm ² was consistent. However, the skin thickness was reported as a range (50 to 98 uM). It is unclear if the variation in thicknesses was consistent across groups, and this may have contributed to some of the endpoint variations (and subsequently high CVs) observed.
	Metric 9:	Reporting of concentrations	High	The target rate was 10 uL/cm ² , which for a finite dose experiment is consistent with OECD TG 28 recommendations. "The actual dose applied to each replicate was determined by subjecting aliquots of the prepared solution to liquid scintillation counting. The total amount of EDC applied to the skin was determined by the total radioactivity applied and the verified specific activity." The verified specific activity for the prepared solution was 0.02686 uCi/mg, and the concentration in the applied formulation was taken at its density (1,235,1000 ug/cm ³). With the information provided, the actual applied concentration in mg/cm ² can be calculated.
	Metric 10:	Exposure frequency	High	The study authors specifically wanted to determine short-term absorption rates, therefore the durations (10 and 60 min) were appropriate based on the purposes of the study.
	Metric 11:	Number of exposure groups and concentration spacing	Low	Fewer than three concentrations were tested. This study only tested one concentration.
Domain 4: Test Model	Metric 12:	Test model (skin)	Low	The test model and descriptive information were reported. Samples of full thickness human cadaver skin were sourced from the National Disease Research Interchange. Abdominal skin samples from 3 Caucasian females were used. The samples were stored frozen at -20 degrees C. It was noted that the samples were removed from the donors within 24 hours of death, and were used within three months. Prior to use, the skin samples were thawed, soaked in water, and the epidermal layer was isolated then stored refrigerated (max duration not specified) until use. The thickness ranged from 50 to 98 uM. The OECD TG 156 specifies that use of epidermal membrane may, in some cases, overestimate human in vivo skin absorption due to the lack of a sufficient barrier function and split thickness (dermatomed) skin is preferred.
	Metric 13:	Number/Replicates per group	Medium	The study used 12 replicates in total (4 samples x 3 donors). At each time point, 6 replicates (2 replicates per donor) were examined. The number of replicates was appropriate (>4 as per OECD 428)
Domain 5: Outcome Assessment	Metric 14:	Outcome assessment methodology	High	The outcome assessment methodology addressed the intended outcome of interest and was sensitive for the outcome. The test followed OECD guidelines 428 and 28; FR volume 69, number 80; and European Commission Guidance Document on Dermal Absorption. Measurement techniques and timing were reported and appropriate for the purposes of the study. A finite dose of the test substance was used to determine short-term maximum absorption rates. A dose volume of 10 uL/cm ² was used.
	Metric 15:	Consistency of outcome assessment	High	Details of outcome assessment protocol were reported and outcomes were assessed consistently across study replicates. The same duration of exposure, receptor fluid used, and sampling period was consistent across replicates.

Continued on next page ...

... continued from previous page

Study Citation:		DuPont Haskell Lab, (2005). Ethylene dichloride: In vitro dermal absorption rate testing.		
HERO ID:		7490381		
		EVALUATION		
Domain	Metric	Rating	Comments	
	Metric 16:	Sampling adequacy and sensitivity	High	The study reported adequate sampling for the outcomes of interest; measurement sensitivity was sufficient. The time and number of scintillation counts were reported. Analysis of a mock dose showed an appropriate signal-to-noise ratio.
Domain 6: Confounding/Variable Control	Metric 17:	Confounding variables in test design and procedures	Medium	There were no differences reported among study groups (test substance lot, number of skin samples, or quality of tissues) that could influence the outcome assessment. Skin integrity was confirmed by EI and only membranes with an EI of $\geq 17 \text{ k}\Omega$ were retained for use in the study. Scintillation counts were done for 5 minutes for most samples. The authors indicated that this deviated from the protocol, but that it had no impact on the interpretation of the study results.
	Metric 18:	Confounding variables in outcomes unrelated to exposure	High	There were no reported differences among the study replicates that were unrelated to exposure; the test substance was demonstrated to be soluble in the receptor fluid.
Domain 7: Data Presentation and Analysis	Metric 19:	Data analysis	Uninformative	The short-term absorption rates were calculated by dividing the sum of the ug equivalents in the receptor fluid by the skin area and exposure time. The total recovery was determined and was the sum of the amount extracted from the charcoal trap, and what was detected in the receptor fluid, in skin washes, in the donor chamber, and in the skin. The coefficients of variation were all $>25\%$ making this study uninformative.
	Metric 20:	Data interpretation	High	Short-term maximum absorption rates were derived from appropriate exposure conditions (finite dose). Recovery of the applied test substance was adequate and fell within guideline specifications for a volatile test material ($100\% \pm 20\%$)
	Metric 21:	Reporting of data	High	Data for all relevant endpoints were reported quantitatively as means \pm SE.
Overall Quality Determination		Uninformative		

Study Citation: Frasch, H. F., Barbero, A. M. (2009). A paired comparison between human skin and hairless guinea pig skin in vitro permeability and lag time measurements for 6 industrial chemicals. *Cutaneous and Ocular Toxicology* 28(3):107-113.

HERO ID: 734137

EXTRACTION**Parameter****Data**

Extraction ID; Chemical: Permeability coefficient (kp) in human skin; 1,1-Dichloroethane-Parent compound

Skin Material/Species; Skin Preparation; Skin Thickness (um); Diffusion Cell Exposure Setup Type: ex vivo human; Split thickness; Not Reported; Static; Notes: Not Reported

Occlusion Type; Donor Chamber Vehicle; Concentration of Test Substance in Vehicle (enter as percent): Occluded; HEPES-buffered Hanks' balanced salt solution; Not Reported

Mass per Surface Area on Skin (mg/cm2); Duration of Test Substance on Skin: 5.3; Other; 2 hrs

Duration of Absorbance Measured; Frequency of Samples: Other; 0.25, 0.5, 0.75, 1.0, 1.5, and 2.0 hours; Notes: 2 hrs

Time Skin was Washed and Method used; Radiolabel Presence: Not Reported; No

Total Recovery (percent); Dose Type: Not Reported; Infinite

Percent Found in Skin Depot After Washing and Tape Stripping; Comments: Not Reported; Notes: Not reported

Percent Found in All Tape Strips, Excluding the Upper Two Strips; Comments: Not Reported; Notes: Not reported

Percent Found in Receptor Fluid and Receptor Fluid Rinse; Comments: Not Reported; Notes: Not Reported

Total Percent Absorbed: 0

Steady State Permeability Coefficient (Kp) (cm/hr); Steady State Permeability Coefficient (Comments); Steady State Flux (ug/cm2/hr); Steady State Flux (Comments); Maximum Permeability Coefficient (Kp) (cm/hr); Maximum Permeability Coefficient (Comments); Maximum Flux (ug/cm2/hr); Maximum Flux (Comments): 0.259; Notes: 27%; Not Reported; Notes: Not Reported; Not Reported; Notes: Not Reported; Not Reported; Notes: Not Reported

EVALUATION

Domain	Metric	Rating	Comments
Domain 1: Test Substance			
	Metric 1: Test substance identity	High	The test substance was identified as 1,1-Dichloroethane with CAS No. 107-06-2.
	Metric 2: Test substance source	High	The source of the test substance was Sigma-Aldrich (St. Louis, MO, USA). Lot or batch number were not provided, but not required for a rating of high.
	Metric 3: Test substance purity	High	The purity of the test substance was reported to be $\geq 99.8\%$.
Domain 2: Test Design			

Continued on next page ...

... continued from previous page

Study Citation:		Frasch, H. F., Barbero, A. M. (2009). A paired comparison between human skin and hairless guinea pig skin in vitro permeability and lag time measurements for 6 industrial chemicals. <i>Cutaneous and Ocular Toxicology</i> 28(3):107-113.		
HERO ID:		734137		
Domain	Metric	EVALUATION Rating	Comments	
	Metric 4:	Reference compounds	Low	A concurrent reference compound was not tested along with the test substance. The study also tested and report data for 5 other chemical (aniline, benzene, diethyl phthalate, naphthalene, and tetrachloroethylene); however, these are not considered reference compounds.
	Metric 5:	Assay procedures	Low	The assay procedure was partially described. The study used a vertical static diffusion set up. The temperature of the skin surface was maintained at 32oC; humidity was not reported. The donor compartment was filled with 0.5 ml of saturated donor solution, the receptor compartment was filled with 5 ml of warmed HEPES-buffered Hanks' balanced salt solution. The study did not report if the receptor compartment fluid was stirred constantly (OECD guidelines recommends constant stirring of receptor fluid). The exposed skin surface area was 9mm. The donor compartment was covered with parafilm, donor solutions were replaced periodically to maintain "infinite" dose conditions. The integrity of the skin was verified under a dissecting microscope before testing for obvious holes, but not by any other means (i.e. electrical resistance, penetration with tritiated water, or checking g for trans-epidermal water loss).
	Metric 6:	Standards for tests	Low	Human skin was obtained from breast reduction or panniculectomies and stored at -85oC the day of surgery. The skin was used for these tests within 191 days. According to OECD Guidance No 28, skin should not be stored at -80oC because it can enhance permeability. The skin should be tested for integrity by either 1) checking electrical resistance; 2) checking trans-epidermal water loss is in normal range; or 3) measure penetration characteristics of a reference material (e.g. tritiated water). None of the three suggestions were done in this study. The study used a dissecting microscope to visually inspect the skin for obvious defects (holes). The coefficient of variance for permeability (kp) was 27%.
Domain 3: Exposure Characterization				
	Metric 7:	Preparation and storage of test substance (chemical)	Medium	Preparation of test substance is adequately reported. Saturated solutions were made by adding excess test chemical to buffer and vortexing overnight at room temperature, and then warming solution to 32oC. When added to donor cell, a small excess of chemical remained to maintain saturation. The storage conditions were not reported. It is not clear if test solutions were prepared immediately before each exposure test or if solutions were stored. Given that the test solution was a saturated solution, with small excess in donor cells, lack of storage information is unlikely to substantially impact results.
	Metric 8:	Consistency of exposure administration	Low	The diameter of skin studied was consistent (9mm) and volume of donor and receptor compartments were consistent. The study did not report the thickness of the skin. It states the epidermis was teased from the dermis using cotton swabs but does not report the thickness. Therefore, it is not known how identical each of the samples were. Permeability and lag time are reported with variance. Coefficient of variation was 27% for permeability and 42% for lag time suggesting substantial variation between donors. It is difficult to know if this is due to human donor variability or lack of consistency in preparation of skin samples.
	Metric 9:	Reporting of concentrations	High	The concentration in the donor solution was sampled in triplicate from each experiment and analyzed using gas chromatography. Concentration is presented with standard deviation.

Continued on next page ...

... continued from previous page

Study Citation:		Frasch, H. F., Barbero, A. M. (2009). A paired comparison between human skin and hairless guinea pig skin in vitro permeability and lag time measurements for 6 industrial chemicals. <i>Cutaneous and Ocular Toxicology</i> 28(3):107-113.		
HERO ID:		734137		
Domain	Metric	EVALUATION Rating	Comments	
	Metric 10:	Exposure frequency	High	The exposure duration of 2 hours was appropriate for outcome of interest (Kp/flux and lag time).
	Metric 11:	Number of exposure groups and concentration spacing	High	The number of exposure groups was appropriate for study design. An infinite dose was studied to determine Kp, in agreement with OECD guidelines.
Domain 4: Test Model	Metric 12:	Test model (skin)	Uninformative	Human skin was obtained from breast reduction or panniculectomies and stored at -85oC the day of surgery. The skin was used for these tests within 191 days. According to OECD 28, skin should not be stored at very low temperatures (with an example of -80oC) because it can enhance permeability. The skin should be tested for integrity by either 1) checking electrical resistance; 2) checking trans-epidermal water loss is in normal range; or 3) measure penetration characteristics of a reference material (e.g. tritiated water). None of the three approaches were taken in this study. The study used a dissecting microscope to visually inspect the skin for obvious defects (holes). Also, the thickness of the skin was not reported. The lack of information and verification of the skin integrity makes this study uninformative for this metric.
	Metric 13:	Number/Replicates per group	Medium	The number of replicates/group (n=4) and replicates per donor (3) were appropriate according to OECD guidelines.
Domain 5: Outcome Assessment	Metric 14:	Outcome assessment methodology	High	Outcome assessment methodology was appropriate. 0.5 ml of an infinite dose was added to the donor compartment. The diameter of the exposed skin was 9mm. The area of the skin was calculated by the reviewer to be 0.63 cm ² , therefore a 793 uL/cm ² of saturated solution was studied and in line with guidance.
	Metric 15:	Consistency of outcome assessment	High	The outcome assessment was carried out consistently across study groups. 0.2 ml samples were taken from the receptor compartment at specified and consistent timepoints. Receptor volume was replaced with the same volume of buffer. Assessment of concentration in receptor fluid was analyzed consistently with gas chromatography.
	Metric 16:	Sampling adequacy and sensitivity	High	The sampling intervals were adequate (0.25, 0.5, 0.75, 1.0, 1.5, and 2.0 hours). The study reports receptor fluid concentration was less than 10% of measured saturation quantities.
Domain 6: Confounding/Variable Control	Metric 17:	Confounding variables in test design and procedures	Low	The study did not measure skin integrity by any an acceptable measure (checking electrical resistance; checking trans-epidermal water loss is in normal range; or measure penetration characteristics of a reference material). The standard deviation between the 4-donor measure for permeability measurements was 27%. (0.259 +/-0.070 cm/h). From each donor, 3 skin disks were studied. The study does not provide data on the variance between the 3 disks from the same donor.
	Metric 18:	Confounding variables in outcomes unrelated to exposure	Medium	Solubility in the receptor fluid was not demonstrated, however study does state the concentration in the receptor fluid was less than 10% of saturated concentration, therefore is in not likely to be an issue.
Domain 7: Data Presentation and Analysis				

Continued on next page ...

...continued from previous page

Study Citation: Frasch, H. F., Barbero, A. M. (2009). A paired comparison between human skin and hairless guinea pig skin in vitro permeability and lag time measurements for 6 industrial chemicals. *Cutaneous and Ocular Toxicology* 28(3):107-113.

HERO ID: 734137

EVALUATION

Domain	Metric	Rating	Comments
	Metric 19: Data analysis	High	Statistical methods were clearly described. The study calculated Kp and lag time (τ) using a mathematical equation. The study states "Use of equation 1 is mathematically equivalent to calculating kp from the slope of the steady-state accumulation curve and τ as the intercept of this asymptote with the time axis."
	Metric 20: Data interpretation	High	Data were interpreted correctly. Permeability (Kp) was calculated using infinite concentration.
	Metric 21: Reporting of data	Medium	Study reports calculated permeability (kp) and lag time but does not provide any raw data. Information on variance between donors is reported, however variance between experiments within the same donor are not reported (4 donors, 3 skin disks from each donor were tested). Data is presented as mean +/- SD of the four donors.

Overall Quality Determination

Uninformative

Study Citation: Frasch, H. F., Barbero, A. M. (2009). A paired comparison between human skin and hairless guinea pig skin in vitro permeability and lag time measurements for 6 industrial chemicals. *Cutaneous and Ocular Toxicology* 28(3):107-113.

HERO ID: 734137

EXTRACTION

Parameter

Data

Extraction ID; Chemical:	Permeability coefficient (kp) in hairless guinea pig skin; 1,1-Dichloroethane-Parent compound
Skin Material/Species; Skin Preparation; Skin Thickness (um); Diffusion Cell Exposure Setup Type:	Other; Split thickness; Not Reported; Static; Notes: guinea pig
Occlusion Type; Donor Chamber Vehicle; Concentration of Test Substance in Vehicle (enter as percent):	Occluded; HEPES-buffered Hanks' balanced salt solution; Not Reported
Mass per Surface Area on Skin (mg/cm ²); Duration of Test Substance on Skin:	5.3; Other; 2 hrs
Duration of Absorbance Measured; Frequency of Samples:	Other; 0.25, 0.5, 0.75, 1.0, 1.5, and 2.0 hours; Notes: 2 hrs
Time Skin was Washed and Method used; Radiolabel Presence:	Not Reported; No
Total Recovery (percent); Dose Type:	Not Reported; Infinite
Percent Found in Skin Depot After Washing and Tape Stripping; Comments:	Not Reported; Notes: Not Reported
Percent Found in All Tape Strips, Excluding the Upper Two Strips; Comments:	Not Reported; Notes: Not Reported
Percent Found in Receptor Fluid and Receptor Fluid Rinse; Comments:	Not Reported; Notes: Not Reported
Total Percent Absorbed:	0
Steady State Permeability Coefficient (Kp) (cm/hr); Steady State Permeability Coefficient (Comments); Steady State Flux (ug/cm ² /hr); Steady State Flux (Comments); Maximum Permeability Coefficient (Kp) (cm/hr); Maximum Permeability Coefficient (Comments); Maximum Flux (ug/cm ² /hr); Maximum Flux (Comments):	0.295; Notes: 32%; Not Reported; Notes: Not Reported; Not Reported; Notes: Not Reported; Not Reported; Notes: Not Reported

EVALUATION

Domain	Metric	Rating	Comments
Domain 1: Test Substance			
	Metric 1:	High	The test substance was identified as 1,1-Dichloroethane with CAS No. 107-06-2.
	Metric 2:	High	The source of the test substance was Sigma-Aldrich (St. Louis, MO, USA). Lot or batch number were not provided, but not required for a rating of high.
	Metric 3:	High	The purity of the test substance was reported to be $\geq 99.8\%$.
Domain 2: Test Design			
	Metric 4:	Low	A concurrent reference compound was not tested along with the test substance. The study also tested and report data for 5 other chemical (aniline, benzene, diethyl phthalate, naphthalene, and tetrachloroethylene); however, these are not considered reference compounds.

Continued on next page ...

... continued from previous page

Study Citation:		Frasch, H. F., Barbero, A. M. (2009). A paired comparison between human skin and hairless guinea pig skin in vitro permeability and lag time measurements for 6 industrial chemicals. <i>Cutaneous and Ocular Toxicology</i> 28(3):107-113.		
HERO ID:		734137		
Domain	Metric	EVALUATION Rating	Comments	
	Metric 5:	Assay procedures	Low	The assay procedure was partially described. The study used a vertical static diffusion set up. The temperature of the skin surface was maintained at 32oC; humidity was not reported. The donor compartment was filled with 0.5 ml of saturated donor solution, the receptor compartment was filled with 5 ml of warmed HEPES-buffered Hanks' balanced salt solution. The test substance is non-polar and lipophilic, therefore there is the potential for inadequate solubility in receptor fluid. To address this the study should have added a modifier to the solution (6% polyethylene glycol 20 oleyl ether or 5% bovine serum albumin as described in OECD guidance document 28). The study did not report if the receptor compartment fluid was stirred constantly (OECD guidelines recommends constant stirring of receptor fluid). The exposed skin surface area was 9mm. The donor compartment was covered with parafilm, donor solutions were replaced periodically to maintain "infinite" dose conditions. Abdominal skin was harvested and used the same day.
	Metric 6:	Standards for tests	Low	Abdominal skin from hairless guinea pigs was harvested and used the same day. Since the skin was not stored, determining integrity by electrical resistance, trans-epidermal water loss or use of a reference material is not required. A dissecting microscope was used to visually inspect the skin for obvious defects (holes). The coefficient of variance for permeability (kp) was 32%.
Domain 3: Exposure Characterization				
	Metric 7:	Preparation and storage of test substance (chemical)	Medium	Preparation of test substance is adequately reported. Saturated solutions were made by adding excess test chemical to buffer and vortexing overnight at room temperature, and then warming solution to 32oC. When added to donor cell, a small excess of chemical remained to maintain saturation. The storage conditions were not reported. It is not clear if test solutions were prepared immediately before each exposure test or if solutions were stored. Given that the test solution was a saturated solution, with small excess in donor cells, lack of storage information is unlikely to substantially impact results.
	Metric 8:	Consistency of exposure administration	Low	The diameter of skin studied was consistent (9mm) and volume of donor and receptor compartments were consistent. The study did not report the thickness of the skin. It states the epidermis was teased from the dermis using cotton swabs but does not report the thickness. Therefore, it is not known how identical each of the samples were. Permeability and lag time are reported with variance. Coefficient of variation was 32% for permeability and 67% for lag time suggesting substantial variation between donors. It is difficult to know if this is due to lack of consistency in preparation of skin samples.
	Metric 9:	Reporting of concentrations	High	The concentration in the donor solution was sampled in triplicate from each experiment and analyzed using gas chromatography. Concentration is presented with standard deviation.
	Metric 10:	Exposure frequency	High	The exposure duration of 2 hours was appropriate for outcome of interest (Kp/flux and lag time).
	Metric 11:	Number of exposure groups and concentration spacing	High	The number of exposure groups was appropriate for study design. An infinite dose was studied to determine Kp, in agreement with OECD guidelines.
Domain 4: Test Model				

Continued on next page ...

... continued from previous page

Study Citation:		Frasch, H. F., Barbero, A. M. (2009). A paired comparison between human skin and hairless guinea pig skin in vitro permeability and lag time measurements for 6 industrial chemicals. <i>Cutaneous and Ocular Toxicology</i> 28(3):107-113.		
HERO ID:		734137		
Domain	Metric	EVALUATION Rating	Comments	
	Metric 12:	Test model (skin)	Low	Abdominal skin from hairless guinea pigs was harvested and used the same day. Heat separated epiderma membranes were used. Skin disks were visually inspected with a dissecting microscope for obvious defects. The thickness of the skin was not reported.
	Metric 13:	Number/Replicates per group	Medium	The number of replicates/group (n=4) and concentration studies (infinite) were appropriate according to OECD guidelines.
Domain 5: Outcome Assessment				
	Metric 14:	Outcome assessment methodology	High	Outcome assessment methodology was appropriate. 0.5 ml of an infinite dose was added to the donor compartment. The diameter of the exposed skin was 9mm. The area of the skin was calculated by the reviewer to be 0.63 cm ² , therefore a 793 uL/cm ² of saturated solution was studied and in line with guidance. The timing of sampling the receptor compartment (0.25, 0.5, 0.75, 1.0, 1.5 and 2.0 hours) was appropriate.
	Metric 15:	Consistency of outcome assessment	High	The outcome assessment was carried out consistently across study groups. 0.2 ml samples were taken from the receptor compartment at specified and consistent timepoints. Receptor volume was replaced with the same volume of buffer. Assessment of concentration in receptor fluid was analyzed consistently with gas chromatography.
	Metric 16:	Sampling adequacy and sensitivity	High	The sampling intervals were adequate (0.25, 0.5, 0.75, 1.0, 1.5, and 2.0 hours). The study reports receptor fluid concentration was less than 10% of measured saturation quantities.
Domain 6: Confounding/Variable Control				
	Metric 17:	Confounding variables in test design and procedures	Low	The thickness of the skin was not reported. Although not required, since skin was used the same day as harvest, integrity was not measured. The coefficient of variation between the 4-donor measure for permeability measurements was 32% (0.295 +/- 0.093 cm/h). From each donor, 3 skin disks were studied. The study does not provide data on the variance between the 3 disks from the same donor.
	Metric 18:	Confounding variables in outcomes unrelated to exposure	Medium	Solubility in the receptor fluid was not demonstrated, however study does state the concentration in the receptor fluid was less than 10% of saturated concentration, therefore is in not likely to be an issue.
Domain 7: Data Presentation and Analysis				
	Metric 19:	Data analysis	High	Statistical methods were clearly described. The study calculated Kp and lag time (τ) using a mathematical equation. The study states "Use of equation 1 is mathematically equivalent to calculating kp from the slope of the steady-state accumulation curve and τ as the intercept of this asymptote with the time axis."
	Metric 20:	Data interpretation	High	Data were interpreted correctly. Permeability (Kp) was calculated using infinite concentration.
	Metric 21:	Reporting of data	Medium	Study reports calculated permeability (kp) and lag time but does not provide any raw data. Information on variance between donors is reported, however variance between experiments within the same donor are not reported (4 donors, 3 skin disks from each donor were tested). Data is presented as mean +/- SD of the four donors.

Overall Quality Determination**Medium**

Continued on next page ...

...continued from previous page

Study Citation: Frasch, H. F., Barbero, A. M. (2009). A paired comparison between human skin and hairless guinea pig skin in vitro permeability and lag time measurements for 6 industrial chemicals. *Cutaneous and Ocular Toxicology* 28(3):107-113.

HERO ID: 734137

EVALUATION

Domain	Metric	Rating	Comments
--------	--------	--------	----------

Study Citation: Frasch, H. F., Barbero, A. M., Alachkar, H., Mcdougal, J. N. (2007). Skin penetration and lag times of neat and aqueous diethyl phthalate, 1,1-Dichloroethane and naphthalene. Cutaneous and Ocular Toxicology 26(2):147-160.

HERO ID: 1313327

EXTRACTION

Parameter	Data
Extraction ID; Chemical:	Neat from the Frasch lab; 1,1-Dichloroethane-Parent compound
Skin Material/Species; Skin Preparation; Skin Thickness (um); Diffusion Cell Exposure Setup Type:	Other; Split thickness; 315; Static; Notes: Hairless guinea pig
Occlusion Type; Donor Chamber Vehicle; Concentration of Test Substance in Vehicle (enter as percent):	Occluded; Not Reported; 99.8
Mass per Surface Area on Skin (mg/cm ²); Duration of Test Substance on Skin:	971; Other; 1.5 hrs
Duration of Absorbance Measured; Frequency of Samples:	Other; 0, 10, 20, 30, 45, 60, 90 minutes; Notes: 1.5 hours
Time Skin was Washed and Method used; Radiolabel Presence:	Not Reported; No
Total Recovery (percent); Dose Type:	Not Reported; Infinite
Percent Found in Skin Depot After Washing and Tape Stripping; Comments:	Not Reported; Notes: no skin depot - flux only
Percent Found in All Tape Strips, Excluding the Upper Two Strips; Comments:	Not Reported; Notes: no skin depot - flux only
Percent Found in Receptor Fluid and Receptor Fluid Rinse; Comments:	Not Reported; Notes: no rinse
Total Percent Absorbed:	0
Steady State Permeability Coefficient (Kp) (cm/hr); Steady State Permeability Coefficient (Comments); Steady State Flux (ug/cm ² /hr); Steady State Flux (Comments); Maximum Permeability Coefficient (Kp) (cm/hr); Maximum Permeability Coefficient (Comments); Maximum Flux (ug/cm ² /hr); Maximum Flux (Comments):	0.00501; Notes: 22%; Not Reported; Notes: Not Reported; 6280; Notes: 22%; Not Reported; Notes: Not Reported

EVALUATION

Domain	Metric	Rating	Comments
Domain 1: Test Substance	Metric 1: Test substance identity	High	Test substance was identified as 1,1-Dichloroethane; CAS RN 107-06-2.
	Metric 2: Test substance source	High	The source of the test substance was Sigma-Aldrich (St. Louis, MO, USA). Lot or batch number were not provided, but this is not required for a rating of high.
	Metric 3: Test substance purity	High	The purity of the test substance was reported to be 99.8%.
Domain 2: Test Design	Metric 4: Reference compounds	Low	A concurrent reference compound was not tested along with the test substance.

Continued on next page ...

... continued from previous page

Study Citation:		Frasch, H. F., Barbero, A. M., Alachkar, H., Mcdougal, J. N. (2007). Skin penetration and lag times of neat and aqueous diethyl phthalate, 1,1-Dichloroethane and naphthalene. Cutaneous and Ocular Toxicology 26(2):147-160.		
HERO ID:		1313327		
EVALUATION				
Domain	Metric		Rating	Comments
	Metric 5:	Assay procedures	Medium	The assay procedure was partially described. The study used a Franz static water-jetted cell apparatus with a 9 mm diameter opening for the skin (0.636 cm ² exposure area). The temperature of the receptor compartment was maintained at 37oC, and the donor compartment at 32oC; humidity was not reported. The donor compartment was 0.5 ml of neat solution. The receptor compartment was filled with 5 ml HEPES-buffered Hanks' balanced salt solution). The test substance is non-polar and lipophilic, therefore there is the potential for inadequate solubility in receptor fluid. To address this the study should have added a modifier to the solution (6% polyethylene glycol 20 oleyl ether or 5% bovine serum albumin as described in OECD guidance document 28). The receptor compartment fluid was stirred at ~1000 rpm. The donor compartment was covered with parafilm, donor solutions were replaced periodically to maintain "infinite" dose conditions. Concentration in receptor fluid was analyzed using gas chromatography and method adequately reported. Although not explicitly stated, from methods it can be inferred harvested skin was studied the same day.
	Metric 6:	Standards for tests	Low	Harvested skin was not stored and used the same day. The integrity of the skin was not tested prior to testing (via electrical resistance, trans-epidermal water loss is in normal range, or tritiated water). The Coefficient of variation was <25%.
Domain 3: Exposure Characterization				
	Metric 7:	Preparation and storage of test substance (chemical)	Medium	Neat test substance was added to the donor cell and covered with parafilm. Storage conditions of stock solution was not reported.
	Metric 8:	Consistency of exposure administration	Medium	The diameter of skin studied was consistent (9mm) and volume of donor and receptor compartments were consistent. Dorsal skin dermatomed at 315 um (after the removal of underlying muscle and fat). Coefficient of variation was <25% for permeability.
	Metric 9:	Reporting of concentrations	Medium	The concentration of neat test substance is not measured, rather the study uses the density of the substance (1.235 ug/cm ³) for calculations. Since the test substance is used neat this is unlikely to substantially impact results.
	Metric 10:	Exposure frequency	Low	The duration of 1.5 hours was much shorter than typical occupational exposure scenarios and may be too short to establish a reliable steady-state flux, although the shallow absorption curve does appear relatively linear by visual inspection.
	Metric 11:	Number of exposure groups and concentration spacing	High	The number of exposure groups was appropriate for study design. An infinite dose was studied to determine Kp, in agreement with OECD guidelines.
Domain 4: Test Model				
	Metric 12:	Test model (skin)	Low	Dorsal skin from a hairless guinea pig skin was harvested, dermatomed to 315 um and used directly. Skin integrity was not measured. The model and skin thickness were appropriate. Lack of measuring integrity could impact results.
	Metric 13:	Number/Replicates per group	Medium	The number of replicates/group (n=8) and concentration studies (infinite) were appropriate according to OECD guidelines.
Domain 5: Outcome Assessment				

Continued on next page ...

...continued from previous page

Study Citation:	Frasch, H. F., Barbero, A. M., Alachkar, H., Mcdougal, J. N. (2007). Skin penetration and lag times of neat and aqueous diethyl phthalate, 1,1-Dichloroethane and naphthalene. Cutaneous and Ocular Toxicology 26(2):147-160.			
HERO ID:	1313327			
		EVALUATION		
Domain	Metric	Rating	Comments	
	Metric 14:	Outcome assessment methodology	High	Outcome assessment methodology was appropriate. 0.5 ml of an infinite dose was added to the donor compartment. The diameter of the exposed skin was 9mm. The area of the skin was calculated by the reviewer to be 0.636 cm ² , therefore a 786 uL/cm ² of saturated solution was studied and in line with guidance. The timing of sampling the receptor compartment (0, 10, 20, 30, 45, 60, and 90 minutes) was appropriate. 100 uL of receptor solution was removed and replaced with the same volume fresh buffer.
	Metric 15:	Consistency of outcome assessment	High	The outcome assessment was carried out consistently across study groups. 100 ul samples were taken from the receptor compartment at specified and consistent timepoints. Receptor volume was replaced with the same volume of buffer. Assessment of concentration in receptor fluid was analyzed consistently with gas chromatography.
	Metric 16:	Sampling adequacy and sensitivity	High	The sampling intervals were adequate to allow for accurate graphical representation of results (0, 10, 20, 30, 45, 60, and 90 minutes).
Domain 6: Confounding/Variable Control				
	Metric 17:	Confounding variables in test design and procedures	Low	The thickness of the skin was consistent. The study did not measure skin integrity by any acceptable measure (checking electrical resistance; checking trans-epidermal water loss is in normal range; or measure penetration characteristics of a reference material).
	Metric 18:	Confounding variables in outcomes unrelated to exposure	High	Solubility in the receptor fluid was not expected to substantially impact results. The receptor compartment volume was 5 ml and the donor compartment volume was 0.5 ml. The solubility is reported to be 5,347 ug/ml at 22oC (determined by this lab), therefore is in not likely to be a limiting factor in the receptor fluid.
Domain 7: Data Presentation and Analysis				
	Metric 19:	Data analysis	High	Statistical methods and data analysis were clearly described.
	Metric 20:	Data interpretation	Medium	Permeability (Kp) was calculated using infinite concentration, however a non-linear regression was used for deriving Kp/flux values. The study also does not report what timepoints were used for Kp/flux calculation, although presumably it was the timepoints beginning from the end of lag time through the end of the experiment.
	Metric 21:	Reporting of data	High	Study reports calculated permeability (kp), steady state flux, and lag time. Mass accumulation vs time plots are shown. Data is presented as mean +/- SD (n=8).

Overall Quality Determination

Medium

Study Citation: Frasch, H. F., Barbero, A. M., Alachkar, H., McDougal, J. N. (2007). Skin penetration and lag times of neat and aqueous diethyl phthalate, 1,1-Dichloroethane and naphthalene. Cutaneous and Ocular Toxicology 26(2):147-160.
HERO ID: 1313327

EXTRACTION

Parameter	Data
Extraction ID; Chemical:	Neat from the McDougal lab; 1,1-Dichloroethane-Parent compound
Skin Material/Species; Skin Preparation; Skin Thickness (um); Diffusion Cell Exposure Setup Type:	Other; Split thickness; 350; Static; Notes: Hairless guinea pig
Occlusion Type; Donor Chamber Vehicle; Concentration of Test Substance in Vehicle (enter as percent):	Occluded; Not Reported; 99.5
Mass per Surface Area on Skin (mg/cm2); Duration of Test Substance on Skin:	1235; Other; 2 hrs
Duration of Absorbance Measured; Frequency of Samples:	Other; 0.25, 0.5, 0.75, 1.0, 1.25, 1.5, 1.75, 2.0 hrs; Notes: 2
Time Skin was Washed and Method used; Radiolabel Presence:	Not Reported; No
Total Recovery (percent); Dose Type:	Not Reported; Infinite
Percent Found in Skin Depot After Washing and Tape Stripping; Comments:	Not Reported; Notes: no skin depot - flux only
Percent Found in All Tape Strips, Excluding the Upper Two Strips; Comments:	Not Reported; Notes: no skin depot - flux only
Percent Found in Receptor Fluid and Receptor Fluid Rinse; Comments:	Not Reported; Notes: no rinse
Total Percent Absorbed:	0
Steady State Permeability Coefficient (Kp) (cm/hr); Steady State Permeability Coefficient (Comments); Steady State Flux (ug/cm2/hr); Steady State Flux (Comments); Maximum Permeability Coefficient (Kp) (cm/hr); Maximum Permeability Coefficient (Comments); Maximum Flux (ug/cm2/hr); Maximum Flux (Comments):	0.00307; Notes: 18.50%; Not Reported; Notes: Not Reported; 3842; Notes: 18.50%; Not Reported; Notes: Not Reported

EVALUATION

Domain	Metric	Rating	Comments
Domain 1: Test Substance			
	Metric 1: Test substance identity	High	Test substance was identified as 1,1-Dichloroethane; CAS RN 107-06-2.
	Metric 2: Test substance source	High	The source of the test substance was Sigma-Aldrich (St. Louis, MO, USA). Lot or batch number were not provided, but this is not required for a rating of high.
	Metric 3: Test substance purity	High	The purity of the test substance was reported to be 99.5%.
Domain 2: Test Design			
	Metric 4: Reference compounds	Low	A concurrent reference compound was not tested along with the test substance.

Continued on next page ...

... continued from previous page

Study Citation:		Frasch, H. F., Barbero, A. M., Alachkar, H., Mcdougal, J. N. (2007). Skin penetration and lag times of neat and aqueous diethyl phthalate, 1,1-Dichloroethane and naphthalene. Cutaneous and Ocular Toxicology 26(2):147-160.		
HERO ID:		1313327		
		EVALUATION		
Domain	Metric	Rating	Comments	
	Metric 5:	Assay procedures	Medium	The assay procedure was partially described. The study used a Franz static water-jetted cell apparatus with a 11.28 mm diameter opening for the skin (1.0 cm ² exposure area). The temperature of the receptor compartment was maintained at 37°C, and the donor compartment at 32°C; humidity was not reported. The donor compartment was 1 ml of neat solution. The receptor compartment was filled with 8 ml HEPES-buffered Hanks' balanced salt solution. The test substance is non-polar and lipophilic, therefore there is the potential for inadequate solubility in receptor fluid. To address this the study should have added a modifier to the solution (6% polyethylene glycol 20 oleyl ether or 5% bovine serum albumin as described in OECD guidance document 28). The receptor compartment fluid was stirred at ~800 rpm. The donor compartment was covered with parafilm, donor solutions were replaced periodically to maintain "infinite" dose conditions. Concentration in receptor fluid was analyzed using gas chromatography and method adequately reported. Although not explicitly stated, from methods it can be inferred harvested skin was studied the same day.
	Metric 6:	Standards for tests	Low	Harvested skin was not stored and used the same day. The integrity of the skin was not tested prior to testing (via electrical resistance, trans-epidermal water loss is in normal range, or tritiated water). The Coefficient of variation was <25%.
Domain 3: Exposure Characterization				
	Metric 7:	Preparation and storage of test substance (chemical)	Medium	Neat test substance was added to the donor cell and covered with parafilm. Storage conditions of stock solution was not reported.
	Metric 8:	Consistency of exposure administration	Medium	The diameter of skin studied was consistent (11.28 mm) and volume of donor and receptor compartments were consistent. Skin from the midscapular area of the guinea pigs back was excised and dermatomed to 350 um. Coefficient of variation was <25% for permeability.
	Metric 9:	Reporting of concentrations	Medium	The concentration of neat test substance is not measured, rather the study uses the density of the substance (1.235 ug/cm ³) for calculations. Since the test substance is used neat this is unlikely to substantially impact results.
	Metric 10:	Exposure frequency	Low	The duration of 2 hours was much shorter than typical occupational exposure scenarios and may be too short to establish a reliable steady-state flux, although the shallow absorption curve does appear relatively linear by visual inspection.
	Metric 11:	Number of exposure groups and concentration spacing	High	The number of exposure groups was appropriate for study design. An infinite dose was studied to determine K _p , in agreement with OECD guidelines.
Domain 4: Test Model				
	Metric 12:	Test model (skin)	Low	Skin from the midscapular area from a hairless guinea pig skin was harvested, dermatomed to 350 um and used directly. Skin integrity was not measured. The model and skin thickness were appropriate. Lack of measuring integrity could impact results.
	Metric 13:	Number/Replicates per group	Medium	The number of replicates/group (n=8) and concentration studies (infinite) were appropriate according to OECD guidelines.
Domain 5: Outcome Assessment				

Continued on next page ...

... continued from previous page

Study Citation:		Frasch, H. F., Barbero, A. M., Alachkar, H., Mcdougal, J. N. (2007). Skin penetration and lag times of neat and aqueous diethyl phthalate, 1,1-Dichloroethane and naphthalene. Cutaneous and Ocular Toxicology 26(2):147-160.		
HERO ID:		1313327		
		EVALUATION		
Domain	Metric	Rating	Comments	
	Metric 14:	Outcome assessment methodology	High	Outcome assessment methodology was appropriate. 1.0 ml of an infinite dose was added to the donor compartment. The diameter of the exposed skin was 11.28 mm. The area of the skin was calculated by the reviewer to be 1.0 cm ² , therefore a 1000 uL/cm ² of saturated solution was studied and in line with guidance. 20uL of receptor volume was removed at 15–30-minute intervals for 4-6 hours and replaced with fresh buffer. “The solubility of each chemical in HBSS was determined to assure that was not a limiting factor in the flux experiment”.
	Metric 15:	Consistency of outcome assessment	High	The outcome assessment was carried out consistently across study groups. 20 ul samples were taken from the receptor compartment at specified and consistent timepoints. Receptor volume was replaced with the same volume of buffer. Assessment of concentration in receptor fluid was analyzed consistently with gas chromatography.
	Metric 16:	Sampling adequacy and sensitivity	High	The sampling intervals were adequate to allow for accurate graphical representation of results (15-30 minutes).
Domain 6: Confounding/Variable Control				
	Metric 17:	Confounding variables in test design and procedures	Low	The thickness of the skin was consistent. The study did not measure skin integrity by any acceptable measure (checking electrical resistance; checking trans-epidermal water loss is in normal range; or measure penetration characteristics of a reference material).
	Metric 18:	Confounding variables in outcomes unrelated to exposure	High	The study authors states “The solubility of each chemical in HBSS was determined to assure that was not a limiting factor in the flux experiment.” The solubility at 20oC in buffer was reported to be 87,000 ug/ml (determined by this lab)..
Domain 7: Data Presentation and Analysis				
	Metric 19:	Data analysis	High	Statistical methods and data analysis were clearly described.
	Metric 20:	Data interpretation	Medium	Permeability (Kp) was calculated using infinite concentration, however a non-linear regression was used for deriving Kp/flux values. The study also does not report what timepoints were used for Kp/flux calculation, although presumably it was the timepoints beginning from the end of lag time through the end of the experiment.
	Metric 21:	Reporting of data	Medium	Study reports calculated permeability (kp), steady state flux, and lag time. Mass accumulation vs time plots are shown. Data is presented as mean +/- SD (n=8). Methods state samples were collected for 4-6 hours, however data is only presented for 2 hours, without an explanation.

Overall Quality Determination

Medium

Study Citation: Frasch, H. F., Barbero, A. M., Alachkar, H., Mcdougal, J. N. (2007). Skin penetration and lag times of neat and aqueous diethyl phthalate, 1,1-Dichloroethane and naphthalene. Cutaneous and Ocular Toxicology 26(2):147-160.
HERO ID: 1313327

EXTRACTION

Parameter	Data
Extraction ID; Chemical:	Aqueous from the Frasch lab; 1,1-Dichloroethane-Parent compound
Skin Material/Species; Skin Preparation; Skin Thickness (um); Diffusion Cell Exposure Setup Type:	Other; Split thickness; 315; Static; Notes: Hairless guinea pig
Occlusion Type; Donor Chamber Vehicle; Concentration of Test Substance in Vehicle (enter as percent):	Occluded; modified Hanks's Balanced Salt Solution (saturated); 0.5
Mass per Surface Area on Skin (mg/cm2); Duration of Test Substance on Skin:	4.2; Other; 1.5 hrs
Duration of Absorbance Measured; Frequency of Samples:	Other; 0, 10, 20, 30, 45, 60, 90 minutes; Notes: 1.5 hours
Time Skin was Washed and Method used; Radiolabel Presence:	Not Reported; No
Total Recovery (percent); Dose Type:	Not Reported; Infinite
Percent Found in Skin Depot After Washing and Tape Stripping; Comments:	Not Reported; Notes: no skin depot - flux only
Percent Found in All Tape Strips, Excluding the Upper Two Strips; Comments:	Not Reported; Notes: no skin depot - flux only
Percent Found in Receptor Fluid and Receptor Fluid Rinse; Comments:	Not Reported; Notes: no rinse
Total Percent Absorbed:	0
Steady State Permeability Coefficient (Kp) (cm/hr); Steady State Permeability Coefficient (Comments); Steady State Flux (ug/cm2/hr); Steady State Flux (Comments); Maxium Permeability Coefficient (Kp) (cm/hr); Maxium Permeability Coefficient (Comments); Maximum Flux (ug/cm2/hr); Maximum Flux (Comments):	0.201; Notes: 17%; Not Reported; Notes: Not Reported; 1076; Notes: 17%; Not Reported; Notes: Not Reported

EVALUATION

Domain	Metric	Rating	Comments
Domain 1: Test Substance			
	Metric 1: Test substance identity	High	Test substance was identified as 1,1-Dichloroethane; CAS RN 107-06-2.
	Metric 2: Test substance source	High	The source of the test substance was Sigma-Aldrich (St. Louis, MO, USA). Lot or batch number were not provided, but this is not required for a rating of high.
	Metric 3: Test substance purity	High	The purity of the test substance was reported to be 99.8%.
Domain 2: Test Design			
	Metric 4: Reference compounds	Low	A concurrent reference compound was not tested along with the test substance.

Continued on next page ...

... continued from previous page

Study Citation:		Frasch, H. F., Barbero, A. M., Alachkar, H., Mcdougal, J. N. (2007). Skin penetration and lag times of neat and aqueous diethyl phthalate, 1,1-Dichloroethane and naphthalene. Cutaneous and Ocular Toxicology 26(2):147-160.		
HERO ID:		1313327		
Domain	Metric	EVALUATION Rating	Comments	
	Metric 5:	Assay procedures	Medium	The assay procedure was partially described. The study used a Franz static water-jetted cell apparatus with a 9 mm diameter opening for the skin (0.636 cm ² exposure area). The temperature of the receptor compartment was maintained at 37oC, and the donor compartment at 32oC; humidity was not reported. The donor compartment was 0.5 ml of saturated solution (in a HEPES-buffered Hanks' balanced salt solution). The receptor compartment was filled with 5 ml HEPES-buffered Hanks' balanced salt solution. The test substance is non-polar and lipophilic, therefore there is the potential for inadequate solubility in receptor fluid. To address this the study should have added a modifier to the solution (6% polyethylene glycol 20 oleyl ether or 5% bovine serum albumin as described in OECD guidance document 28). The receptor compartment fluid was stirred at ~1000 rpm. The donor compartment was covered with parafilm, donor solutions were replaced periodically to maintain "infinite" dose conditions. Concentration in receptor fluid was analyzed using gas chromatography and method adequately reported. Although not explicitly stated, from methods it can be inferred harvested skin was studied the same day.
	Metric 6:	Standards for tests	Low	Harvested skin was not stored and used the same day. The integrity of the skin was not tested prior to testing (via electrical resistance, trans-epidermal water loss is in normal range, or tritiated water). The Coefficient of variation was <25%.
Domain 3: Exposure Characterization				
	Metric 7:	Preparation and storage of test substance (chemical)	Medium	Saturated aqueous solutions were made fresh prior to the experiment. Excess chemical was added to HBSS buffer and vortexed for approximately 24 hours at room temperature before being centrifuged for 30 minutes at 4,000 rpms. Storage conditions of stock solution was not reported, however since excess chemical was added to buffer, any evaporation of stock solution that may have occurred is not likely to impact results.
	Metric 8:	Consistency of exposure administration	Medium	The diameter of skin studied was consistent (9mm) and volume of donor and receptor compartments were consistent. Dorsal skin dermatomed at 315 um (after the removal of underlying muscle and fat). Coefficient of variation was <25% for permeability.
	Metric 9:	Reporting of concentrations	Medium	The donor solution is a saturated solution. The study does not measure the concentration, rather uses the measured a measured solubility in buffer (5,347 ug/ml) as the assumed concentration for calculations.
	Metric 10:	Exposure frequency	Low	The duration of 1.5 hours was much shorter than typical occupational exposure scenarios and may be too short to establish a reliable steady-state flux, although the shallow absorption curve does appear relatively linear by visual inspection.
	Metric 11:	Number of exposure groups and concentration spacing	High	The number of exposure groups was appropriate for study design. An infinite dose was studied to determine Kp, in agreement with OECD guidelines.
Domain 4: Test Model				
	Metric 12:	Test model (skin)	Low	Dorsal skin from a hairless guinea pig skin was harvested, dermatomed to 315 um and used directly. Skin integrity was not measured. The model and skin thickness were appropriate. Lack of measuring integrity could impact results.
	Metric 13:	Number/Replicates per group	Medium	The number of replicates/group (n=8) and concentration studies (infinite) were appropriate according to OECD guidelines.

Continued on next page ...

... continued from previous page

Study Citation: Frasch, H. F., Barbero, A. M., Alachkar, H., Mcdougal, J. N. (2007). Skin penetration and lag times of neat and aqueous diethyl phthalate, 1,1-Dichloroethane and naphthalene. Cutaneous and Ocular Toxicology 26(2):147-160.

HERO ID: 1313327

		EVALUATION		
Domain	Metric		Rating	Comments
Domain 5: Outcome Assessment				
	Metric 14:	Outcome assessment methodology	Low	The aqueous saturated solution did not meet the definition of infinite dose. Based on solubility, the concentration was calculated as only 0.5% 1,2DCE, which results in only 2.5ul of 1,2DCE from the 500ul dosed saturated solution. For 0.636 cm2 surface area, this results in loading of only 3.9 ul/cm2, which is well below the recommended loading of 100 ul/cm2. This metric was not scored as unacceptable because a steady-state dose was able to be derived based on regression analysis, however it was nonlinear regression and the absorption curve is very shallow which makes visual inspection difficult. The timing of sampling the receptor compartment (0, 10, 20, 30, 45, 60, and 90 minutes) was appropriate based on the absorption curve, although longer duration would be preferred to establish steady-state. 100 uL of receptor solution was removed and replaced with the same volume fresh buffer and aqueous donor solution was replaced every 15 minutes. .
	Metric 15:	Consistency of outcome assessment	High	The outcome assessment was carried out consistently across study groups. 100 ul samples were taken from the receptor compartment at specified and consistent timepoints. Receptor volume was replaced with the same volume of buffer. Assessment of concentration in receptor fluid was analyzed consistently with gas chromatography.
	Metric 16:	Sampling adequacy and sensitivity	High	The sampling intervals were adequate to allow for accurate graphical representation of results (0, 10, 20, 30, 45, 60, and 90 minutes).
Domain 6: Confounding/Variable Control				
	Metric 17:	Confounding variables in test design and procedures	Low	The thickness of the skin was consistent. The study did not measure skin integrity by any acceptable measure (checking electrical resistance; checking trans-epidermal water loss is in normal range; or measure penetration characteristics of a reference material).
	Metric 18:	Confounding variables in outcomes unrelated to exposure	High	Solubility in the receptor fluid was not expected to substantially impact results. The receptor compartment volume was 5 ml and the donor compartment volume was 0.5 ml. The solubility is reported to be 5,347 ug/ml at 22oC (determined by this lab), therefore is in not likely to be a limiting factor in the receptor fluid.
Domain 7: Data Presentation and Analysis				
	Metric 19:	Data analysis	High	Statistical methods and data analysis were clearly described.
	Metric 20:	Data interpretation	Low	Permeability (Kp) was calculated using less-than infinite concentration, and a non-linear regression was used for deriving Kp/flux values. The study also does not report what timepoints were used for Kp/flux calculation, although presumably it was the timepoints beginning from the end of lag time through the end of the experiment.
	Metric 21:	Reporting of data	High	Study reports calculated permeability (kp), steady state flux, and lag time. Mass accumulation vs time plots are shown. Data is presented as mean +/- SD (n=8).

Overall Quality Determination**Medium**

Study Citation: Gajjar, R. M., Kasting, G. B. (2014). Absorption of ethanol, acetone, benzene and 1,1-Dichloroethane through human skin in vitro: a test of diffusion model predictions. Toxicology and Applied Pharmacology 281(1):109-117.
HERO ID: 4141541

EXTRACTION

Parameter	Data
Extraction ID; Chemical:	31.5 mg/cm2 group; 1,1-Dichloroethane-Parent compound
Skin Material/Species; Skin Preparation; Skin Thickness (um); Diffusion Cell Exposure Setup Type:	ex vivo human; Split thickness; 300; Static; Notes: Not Reported
Occlusion Type; Donor Chamber Vehicle; Concentration of Test Substance in Vehicle (enter as percent):	Unoccluded; None specified; 99
Mass per Surface Area on Skin (mg/cm2); Duration of Test Substance on Skin:	31.5; 24 hrs; Not Reported
Duration of Absorbance Measured; Frequency of Samples:	24 hrs; Samples were taken at 5, 10, 15, 20, 40 minutes and 1, 2, 4, 8, 24-hours post-dose.; Notes: Not Reported
Time Skin was Washed and Method used; Radiolabel Presence:	Skin was not washed; Yes
Total Recovery (percent); Dose Type:	Not Reported; Finite
Percent Found in Skin Depot After Washing and Tape Stripping; Comments:	0.08; Notes: Less than 0.08% was retained in the skin (global statement not specific exposure group). Exact values were not provided. The study authors considered the amount to be negligible.
Percent Found in All Tape Strips, Excluding the Upper Two Strips; Comments:	0.08; Notes: Less than 0.08% was retained in the skin (global statement not specific exposure group). Exact values were not provided. The study authors considered the amount to be negligible.
Percent Found in Receptor Fluid and Receptor Fluid Rinse; Comments:	0.163; Notes: Calculated based on a the reported accumulated amount permeated at 24 hours (51.4 ± 8.1 ug/cm2)
Total Percent Absorbed:	0
Steady State Permeability Coefficient (Kp) (cm/hr); Steady State Permeability Coefficient (Comments); Steady State Flux (ug/cm2/hr); Steady State Flux (Comments); Maxium Permeability Coefficient (Kp) (cm/hr); Maxium Permeability Coefficient (Comments); Maximum Flux (ug/cm2/hr); Maximum Flux (Comments):	Not Reported; Notes: Not Reported; Not Reported; Notes: Not Reported; Not Reported; Notes: Not Reported; Not Reported; Notes: The text qualitatively stated that the maximum flux was reached within 10 minutes post-dose for all doses tested and values ranged from 37 to 193 ug/cm2/hour. The exact values could not be extracted from figures shown.

EVALUATION

Domain	Metric	Rating	Comments
Domain 1: Test Substance			
	Metric 1: Test substance identity	High	The test substance was identified as 1,1-Dichloroethane (radiolabeled 14C-1,2-DCE). The test substance is sold in liquid form (as per supplier websites). The radiolabeled test substance was a neat liquid in break seal or ethanol solvent. An unlabeled neat test substance was also mentioned in the methods but was not mentioned further.
	Metric 2: Test substance source	High	The sources of the test substance were clearly reported although batch/lot numbers were not provided. The radiolabeled test substance was purchased from American Radiolabeled. The unlabeled test substance was purchased from Sigma Aldrich.

Continued on next page ...

... continued from previous page

Study Citation:		Gajjar, R. M., Kasting, G. B. (2014). Absorption of ethanol, acetone, benzene and 1,1-Dichloroethane through human skin in vitro: a test of diffusion model predictions. Toxicology and Applied Pharmacology 281(1):109-117.		
HERO ID:		4141541		
Domain	Metric	EVALUATION Rating	Comments	
	Metric 3:	Test substance purity	High	The purity of the radiolabeled chemical was 99.0%. It is not clear which unlabeled product was purchased from Sigma-Aldrich, so the purity cannot be definitively determined. Because the study did not specify further use of the unlabeled material, this is not considered to be relevant.
Domain 2: Test Design	Metric 4:	Reference compounds	Low	The study did not include any of the specified concurrent controls (caffeine, testosterone, or benzoic acid). The study did cite other publications on their diffusion skin model, but it is not known whether they previously performed skin absorption studies. Other solvents including ethanol, acetone, and benzene were also tested in this study.
	Metric 5:	Assay procedures	Low	The study adequately described the diffusion cell setup. The receptor fluid was PBS which is acceptable for a non-lipophilic compound; however, 1,2-DCE is lipophilic, and the use of a saline solution may not be appropriate (OECD No. 156 Guidance, 2019). It is unclear what effect this may have on the study results. Upon collection, the receptor fluid was immediately placed into scintillation vials to minimize loss and radioactivity was measured within an hour. The temperature was reported and was appropriate (32 degrees C); humidity was not specified. The study was conducted in a fume hood; therefore, the humidity was likely low. A wind velocity was 0.92 m/s was reported. The static set-up was continuously stirred with a magnetic stir bar. The glass tops were non-occluded. The doses per skin surface area were clearly reported. The volumes applied, up to 40 ul per 0.79 cm ² cell, were higher than recommended (up to 10 uL/cm ²). The volume of the receptor fluid was not reported. Receptor fluid was collected at 5, 10, 15, 20, and 40 minutes, and at 1, 2, 4, 8, and 24 hours post-dosing. The study did not use tape strips but measured the total amount of radioactivity in the tissue 24 hours after application. The skin was not washed, but due to evaporation, it is unlikely that there was any unabsorbed test material left on the skin surface. The study did not indicate the use of a carbon trap filter for a volatile substance.
	Metric 6:	Standards for tests	Low	The study did not conduct all of the recommended QC criteria. Skin integrity was assessed prior to the start of exposure; any samples with permeation values > 1.2uL/cm ² were discarded. The study did not explicitly report the overall percentage recovered. The cumulative permeated over 24 hours (ranging from 0.21% to 0.13%), as well as the amount of radioactivity remaining in the tissue (reported to be <0.08% -data not shown) was noted. The study authors did not justify why carbon filter traps were not utilized. Without complete measurements of recovery, it cannot be excluded that, for example, there wasn't loss due to absorbed material onto experimental equipment. CV values were not explicitly reported. Data were presented as means ± SE were reported for data from three donors (with an n = 2-7 per donor). All data were first averaged by dose for each donor, and the results represent the average across donors (indicating an n =3). Assuming an n of 3, CV values can be calculated; these values are > 25% for the 7.9, 15.8, and 31.5 mg/cm ² groups, but < 25% for the 63.1 mg/cm ² group. Further discussion of CVs is in Metric 19.
Domain 3: Exposure Characterization				

Continued on next page ...

... continued from previous page

Study Citation:		Gajjar, R. M., Kasting, G. B. (2014). Absorption of ethanol, acetone, benzene and 1,1-Dichloroethane through human skin in vitro: a test of diffusion model predictions. <i>Toxicology and Applied Pharmacology</i> 281(1):109-117.		
HERO ID:		4141541		
		EVALUATION		
Domain	Metric	Rating	Comments	
	Metric 7: Preparation and storage of test substance (chemical)	Low	Details of the test solution preparation and storage of the volatile test substance were not provided. The stability and solubility of the test substance were not discussed, and the properties of the test substance (insoluble, lipophilic) suggest solubility may be poor. It is presumed that the substance was applied neat (no vehicle), but it is unclear whether the radiolabeled substance was diluted with the non-labelled substance. Long-term storage could have an impact on the specific activity; the specific activity of the radiolabeled test substance at purchase was reported. It is unclear if the reported specific activity at the time of dosing (0.05 uCi/uL) was measured in a scintillation counter or simply based on the activity at purchase (5.0 mCi/mmol).	
	Metric 8: Consistency of exposure administration	Low	Different volumes (ranging from 5 to 40 uL) were applied across groups to obtain different exposure concentrations. The different volumes applied likely had an impact on the study results. The skin thickness and area of the skin were uniform across groups.	
	Metric 9: Reporting of concentrations	Low	The exposure concentrations were clearly reported. Volumes of 5, 10, 20, and 40 uL of the radiolabeled test substance were added per 0.79 cm ² cell, with a specific activity of 0.05 uCi/uL, yielding concentrations of 7.9, 15.8, 31.5, and 63.1 mg/cm ² . The concentrations were not analytically verified, and it was not reported whether the specific activity of the test solutions was measured.	
	Metric 10: Exposure frequency	High	The skin was exposed to a finite dose and receptor solutions were collected up to 24 hours post-dosing. The purpose of the study was to present a method for estimating the absorption of small doses of liquid VOCs that are splashed on the skin, so the finite dose and 24-hour post-exposure monitoring period were appropriate for the purposes of the study. However, due to the volatility of the test substance, there was clearly a significant amount of evaporation and essentially no test material was present in the receptor fluid after 1-hour post-exposure, and max absorption rates were obtained within 10-20 min. A shorter duration in this case may also have been appropriate.	
	Metric 11: Number of exposure groups and concentration spacing	Low	The study tested 4 exposure concentrations. The concentrations and spacing were not explicitly justified by the study authors. The goal was to mimic small doses of liquid VOCs that are splashed on the skin, but it was not specified what those expected doses/volumes would be. The doses used exceeded the typically recommended dose for an in vitro dermal absorption test (1-5 mg/cm ² for finite dose experiments as per WHO2006, OECD TG 428 etc.,). An inverse dose-response (decreased absorption with increasing dose) was obtained. It is unclear if max absorption would be obtained at even lower concentrations not tested in this study.	
Domain 4: Test Model	Metric 12: Test model (skin)	High	The test used dermatomed split-thickness human cadaver skin (~0.3 mm thick) either from the back, abdomen, or thigh. These are acceptable sites for skin samples. The source of the skin samples was provided. The skin came treated with a 10% glycerol solution and was stored at -80 degrees C. The skin samples were washed prior to use and cut into 1.5 cm x 1.5 cm squares that were mounted onto modified Franz diffusion cells. According to OECD No. 28, skin should not be stored at low temperatures because storage at -80 degrees C can enhance permeability. However, skin integrity was tested pre-study, and only samples with water permeation values less than 1.2 uL/cm ² were used.	

Continued on next page ...

... continued from previous page

Study Citation:		Gajjar, R. M., Kasting, G. B. (2014). Absorption of ethanol, acetone, benzene and 1,1-Dichloroethane through human skin in vitro: a test of diffusion model predictions. Toxicology and Applied Pharmacology 281(1):109-117.		
HERO ID:		4141541		
Domain	Metric	EVALUATION Rating	Comments	
	Metric 13:	Number/Replicates per group	Medium	Permeation experiments were done using skin from 3 donors with an n = 2-7 per donor. The text did indicate that in most cases, conditions were tested with 4-5 replicates for each of the 3 donors, yielding 12-15 replicates per dose. This meets or exceeds the guideline recommendation of 4 replicates per test preparation.
Domain 5: Outcome Assessment				
	Metric 14:	Outcome assessment methodology	Low	The purpose of the study was to develop a methodology for estimating absorption from small doses of liquid VOCs. The model estimates were compared to experimental values generated by using human cadaver skin. The outcome methodology was generally appropriate for the purposes of the study; a skin surface of 0.79 cm ² was loaded with different volumes (5, 10, 20, and 40 uL) of the neat test material. The volumes applied at higher doses were greater than guideline recommendations (up to 10 ul/cm ²). Cumulative skin permeation was determined by measuring radioactivity in receptor fluid over a 24-hour period and in the skin after 24 hours. Carbon traps were not used to capture the evaporation of a volatile chemical. The study was conducted in a fume hood rather than under ambient conditions, although the authors indicated fume hood was used for safety reasons. However, since most of the applied dose presumably evaporated from the skin surface, the higher airflow may not have been appropriate. The study authors often point to the higher airflow as justification for the low permeation percentages. Solubility in the receptor fluid was not tested. This is considered to be a significant flaw as PBS is not generally recommended for a lipophilic compound. The study did not wash the skin sample, which deviates from OECD TG 428 recommendations.
	Metric 15:	Consistency of outcome assessment	High	Details of the outcome assessment protocol were reported, and the outcomes were consistently assessed across groups
	Metric 16:	Sampling adequacy and sensitivity	Medium	Limited details on scintillation counting were provided. Receptor fluid was poured into scintillation vials containing 12 mL of scintillation fluid and vials were measured within 1 hour of collection. Skin samples were dissolved in Soluene and analyzed by LSC. The number of scintillation counts per sample, signal to noise ratio/background levels were not reported. It is not clear that any background measurements were taken. The number of evaluations per exposure group in the data table indicates 2-7 per donor per dose, and there were 3 donors.
Domain 6: Confounding/Variable Control				
	Metric 17:	Confounding variables in test design and procedures	Low	The skin batch/lot numbers were not reported. The number of skin samples per group varied (2-7 samples per donor, 3 donors/ group). Skin integrity was measured based on 3H ₂ O permeation. It was specified that samples with water permeation > 1.2 uL/cm ² were discarded, therefore there were no differences in the quality of the tissue. The results were not expressed as Kp values or % of the applied dose. Electrical resistance and TEWL measurements were not conducted. It is unknown whether the skin sample collection sites (back, abdomen, or shoulder) were consistent within or across groups. The distribution of the samples in each group was not specified, and this missing information is likely to have a significant impact on the study results.
Continued on next page ...				

... continued from previous page

Study Citation:		Gajjar, R. M., Kasting, G. B. (2014). Absorption of ethanol, acetone, benzene and 1,1-Dichloroethane through human skin in vitro: a test of diffusion model predictions. Toxicology and Applied Pharmacology 281(1):109-117.		
HERO ID:		4141541		
Domain	Metric	EVALUATION		Comments
			Rating	
	Metric 18:	Confounding variables in outcomes unrelated to exposure	Low	Solubility of the test substance in the receptor fluid is a potential concern. The test substance has low solubility in water (8.6 g/L) and is lipophilic. The receptor fluid (PBS) is generally appropriate for water soluble compounds but may not be appropriate for lipophilic compounds. The solubility of the test substance in the receptor fluid was not reported, and it is unclear if the permeation rate was influenced by solubility, and whether this had a confounding impact, particularly at the higher doses.
Domain 7: Data Presentation and Analysis				
	Metric 19:	Data analysis	Uninformative	The cumulative amount of permeated test substance was reported across time the data were reported as means donors \pm SE. The percent of dose permeated was graphically displayed and reported numerically in the text. Data were first averaged by dose for each donor, and then averaged across donors to obtain the mean values. It is unknown whether the study accounted for any outliers. The study did not report the percentage absorbed, but this could be determined by dividing the cumulative amount permeated by the applied dose, although it should be taken into consideration that it is presumed that the majority of the applied dose evaporated. Based on the data provided for the cumulative amount permeated (at 24hrs), the standard deviations can first be determined ($SD = SE \times \sqrt{n}$, where $n = 3$). The CV values can then be calculated (SD relative to the mean). The calculated CV values are 35%, 44%, 27%, and 23%, for the 7.9, 15.8, 31.5, and 63.1 mg/cm ² groups, respectively. CV values >25% are considered to be unacceptable.
	Metric 20:	Data interpretation	Medium	The percent recovery of the applied test substance was not explicitly reported, although it is presumed to be low because the total cumulative penetration amount ranged from 0.21% to 0.13% of the applied dose. The study authors presumed the remainder evaporated; however, carbon traps were not used to definitively account for evaporated material. The skin was not washed (no measurements of washes), and no measurements of amounts in the donor chamber were collected. Only percentage absorption in the skin from a finite dose is reported and is not defined by dose group (quantitative values not reported). It is not specified whether the absorption estimates were normalized, but it is highly likely that the missing material evaporated and therefore, was unlikely to have been absorbed (under the conditions of this test); the inclusion of evaporated material may not be required. The study also reported the maximum flux of the VOC into the receptor solution. The study authors believed that the data implied that mild lipid barrier disruption likely occurred.
	Metric 21:	Reporting of data	Low	Overall cumulative absorption for each exposure group, at each collection point, was reported as Means \pm SE, and the percentage of the applied dose was noted in the text for some, but not all exposure groups. These values presumably represent primarily the amounts measured in receptor fluid. The amount in the skin after 24 hours was simply reported to be less than 0.08% (presumably of the applied dose), and was not described by exposure group. It is not clear if this amount was included in the reported cumulative values. Separate measurements for receptor fluid alone were not reported.

Overall Quality Determination**Uninformative**

Study Citation: Gajjar, R. M., Kasting, G. B. (2014). Absorption of ethanol, acetone, benzene and 1,1-Dichloroethane through human skin in vitro: a test of diffusion model predictions. *Toxicology and Applied Pharmacology* 281(1):109-117.

HERO ID: 4141541

EXTRACTION

Parameter	Data
Extraction ID; Chemical:	63.1 mg/cm2 group; 1,1-Dichloroethane-Parent compound
Skin Material/Species; Skin Preparation; Skin Thickness (um); Diffusion Cell Exposure Setup Type:	ex vivo human; Split thickness; 300; Static; Notes: Not Reported
Occlusion Type; Donor Chamber Vehicle; Concentration of Test Substance in Vehicle (enter as percent):	Unoccluded; None specified; 99
Mass per Surface Area on Skin (mg/cm2); Duration of Test Substance on Skin:	63.1; 24 hrs; Not Reported
Duration of Absorbance Measured; Frequency of Samples:	24 hrs; Absorption was measured for 24 hours following a single application.; Notes: Not Reported
Time Skin was Washed and Method used; Radiolabel Presence:	Skin was not washed; Yes
Total Recovery (percent); Dose Type:	Not Reported; Finite
Percent Found in Skin Depot After Washing and Tape Stripping; Comments:	0.08; Notes: Less than 0.08% was retained in the skin (global statement not specific exposure group). Exact values were not provided. The study authors considered the amount to be negligible.
Percent Found in All Tape Strips, Excluding the Upper Two Strips; Comments:	0.08; Notes: Less than 0.08% was retained in the skin (global statement not specific exposure group). Exact values were not provided. The study authors considered the amount to be negligible.
Percent Found in Receptor Fluid and Receptor Fluid Rinse; Comments:	0.133; Notes: Calculated based on a the reported accumulated amount permeated at 24 hours (84.4 ± 11.4 ug/cm2)
Total Percent Absorbed:	0
Steady State Permeability Coefficient (Kp) (cm/hr); Steady State Permeability Coefficient (Comments); Steady State Flux (ug/cm2/hr); Steady State Flux (Comments); Maxium Permeability Coefficient (Kp) (cm/hr); Maxium Permeability Coefficient (Comments); Maximum Flux (ug/cm2/hr); Maximum Flux (Comments):	Not Reported; Notes: Not Reported; Not Reported; Notes: Not Reported; Not Reported; Notes: Not Reported; Not Reported; Notes: The text qualitatively stated that the maximum flux was reached within 10 minutes post-dose for all doses tested and values ranged from 37 to 193 ug/cm2/hour. The exact values could not be extracted from figures shown.

EVALUATION

Domain	Metric	Rating	Comments
Domain 1: Test Substance			
	Metric 1:	High	Test substance identity
	Metric 2:	High	Test substance source
	Metric 3:	High	Test substance purity

Continued on next page ...

... continued from previous page

Study Citation: Gajjar, R. M., Kasting, G. B. (2014). Absorption of ethanol, acetone, benzene and 1,1-Dichloroethane through human skin in vitro: a test of diffusion model predictions. *Toxicology and Applied Pharmacology* 281(1):109-117.

HERO ID: 4141541

		EVALUATION		
Domain	Metric	Rating	Comments	
Domain 2: Test Design				
	Metric 4:	Reference compounds	Low	The study did not include any of the specified concurrent controls (caffeine, testosterone, or benzoic acid). The study did cite other publications on their diffusion skin model, but it is not known whether they previously performed skin absorption studies. Other solvents including ethanol, acetone, and benzene were also tested in this study.
	Metric 5:	Assay procedures	Low	The study adequately described the diffusion cell setup. The receptor fluid was PBS which is acceptable for a non-lipophilic compound; however, 1,2-DCE is lipophilic, and the use of a saline solution may not be appropriate (OECD No. 156 Guidance, 2019). It is unclear what effect this may have on the study results. Upon collection, the receptor fluid was immediately placed into scintillation vials to minimize loss and radioactivity was measured within an hour. The temperature was reported and was appropriate (32 degrees C); humidity was not specified. The study was conducted in a fume hood; therefore, the humidity was likely low. A wind velocity was 0.92 m/s was reported. The static set-up was continuously stirred with a magnetic stir bar. The glass tops were non-occluded. The doses per skin surface area were clearly reported. The volumes applied, up to 40 ul per 0.79 cm ² cell, were higher than recommended (up to 10 uL/cm ²). The volume of the receptor fluid was not reported. Receptor fluid was collected at 5, 10, 15, 20, and 40 minutes, and at 1, 2, 4, 8, and 24 hours post-dosing. The study did not use tape strips but measured the total amount of radioactivity in the tissue 24 hours after application. The skin was not washed, but due to evaporation, it is unlikely that there was any unabsorbed test material left on the skin surface. The study did indicate the use of a carbon trap filter for a volatile substance.
	Metric 6:	Standards for tests	Low	The study did not conduct all of the recommended QC criteria. Skin integrity was assessed prior to the start of exposure; any samples with permeation values > 1.2uL/cm ² were discarded. The study did not explicitly report the overall percentage recovered. The cumulative permeated over 24 hours (ranging from 0.21% to 0.13%), as well as the amount of radioactivity remaining in the tissue (reported to be <0.08% -data not shown) was noted. The study authors did not justify why carbon filter traps were not utilized. Without complete measurements of recovery, it cannot be excluded that, for example, there wasn't loss due to absorbed material onto experimental equipment. CV values were not explicitly reported. Data were presented as means ± SE were reported for data from three donors (with an n = 2-7 per donor). All data were first averaged by dose for each donor, and the results represent the average across donors (indicating an n =3). Assuming an n of 3, CV values can be calculated; these values are > 25% for the 7.9, 15.8, and 31.5 mg/cm ² groups, but < 25% for the 63.1 mg/cm ² group. Further discussion of CVs is in Metric 19.
Domain 3: Exposure Characterization				
Continued on next page ...				

... continued from previous page

Study Citation: Gajjar, R. M., Kasting, G. B. (2014). Absorption of ethanol, acetone, benzene and 1,1-Dichloroethane through human skin in vitro: a test of diffusion model predictions. *Toxicology and Applied Pharmacology* 281(1):109-117.

HERO ID: 4141541

EVALUATION				
Domain	Metric	Rating	Comments	
	Metric 7:	Preparation and storage of test substance (chemical)	Low	Details of the test solution preparation and storage of the volatile test substance were not provided. The stability and solubility of the test substance were not discussed, and the properties of the test substance (insoluble, lipophilic) suggest solubility may be poor. It is presumed that the substance was applied neat (no vehicle), but it is unclear whether the radiolabeled substance was diluted with the non-labelled substance. Long-term storage could have an impact on the specific activity; the specific activity of the radiolabeled test substance at purchase was reported. It is unclear if the reported specific activity at the time of dosing (0.05 uCi/uL) was measured in a scintillation counter or simply based on the activity at purchase (5.0 mCi/mmol).
	Metric 8:	Consistency of exposure administration	Low	Different volumes (ranging from 5 to 40 uL) were applied across groups to obtain different exposure concentrations. The different volumes applied likely had an impact on the study results. The skin thickness and area of the skin were uniform across groups.
	Metric 9:	Reporting of concentrations	Low	The exposure concentrations were clearly reported. Volumes of 5, 10, 20, and 40 uL of the radiolabeled test substance were added per 0.79 cm ² cell, with a specific activity of 0.05 uCi/uL, yielding concentrations of 7.9, 15.8, 31.5, and 63.1 mg/cm ² . The concentrations were not analytically verified, and it was not reported whether the specific activity of the test solutions was measured.
	Metric 10:	Exposure frequency	High	The skin was exposed to a finite dose and receptor solutions were collected up to 24 hours post-dosing. The purpose of the study was to present a method for estimating the absorption of small doses of liquid VOCs that are splashed on the skin, so the finite dose and 24-hour post-exposure monitoring period were appropriate for the purposes of the study. However, due to the volatility of the test substance, there was clearly a significant amount of evaporation and essentially no test material was present in the receptor fluid after 1-hour post-exposure, and max absorption rates were obtained within 10-20 min. A shorter duration in this case may also have been appropriate.
	Metric 11:	Number of exposure groups and concentration spacing	Low	The study tested 4 exposure concentrations. The concentrations and spacing were not explicitly justified by the study authors. The goal was to mimic small doses of liquid VOCs that are splashed on the skin, but it was not specified what those expected doses/volumes would be. The doses used exceeded the typically recommended dose for an in vitro dermal absorption test (1-5 mg/cm ² for finite dose experiments as per WHO2006, OECD TG 428 etc.,). An inverse dose-response (decreased absorption with increasing dose) was obtained. It is unclear if max absorption would be obtained at even lower concentrations not tested in this study.
Domain 4: Test Model	Metric 12:	Test model (skin)	High	The test used dermatomed split-thickness human cadaver skin (~0.3 mm thick) either from the back, abdomen, or thigh. These are acceptable sites for skin samples. The source of the skin samples was provided. The skin came treated with a 10% glycerol solution and was stored at -80 degrees C. The skin samples were washed prior to use and cut into 1.5 cm x 1.5 cm squares that were mounted onto modified Franz diffusion cells. According to OECD No. 28, skin should not be stored at low temperatures because storage at -80 degrees C can enhance permeability. However, skin integrity was tested pre-study, and only samples with water permeation values less than 1.2 uL/cm ² were used.

Continued on next page ...

... continued from previous page

Study Citation: Gajjar, R. M., Kasting, G. B. (2014). Absorption of ethanol, acetone, benzene and 1,1-Dichloroethane through human skin in vitro: a test of diffusion model predictions. *Toxicology and Applied Pharmacology* 281(1):109-117.

HERO ID: 4141541

		EVALUATION		
Domain	Metric	Rating	Comments	
	Metric 13:	Number/Replicates per group	Medium	Permeation experiments were done using skin from 3 donors with an n = 2-7 per donor. The text did indicate that in most cases, conditions were tested with 4-5 replicates for each of the 3 donors, yielding 12-15 replicates per dose. This meets or exceeds the guideline recommendation of 4 replicates per test preparation.
Domain 5: Outcome Assessment				
	Metric 14:	Outcome assessment methodology	Low	The purpose of the study was to develop a methodology for estimating absorption from small doses of liquid VOCs. The model estimates were compared to experimental values generated by using human cadaver skin. The outcome methodology was generally appropriate for the purposes of the study; a skin surface of 0.79 cm ² was loaded with different volumes (5, 10, 20, and 40 uL) of the neat test material. The volumes applied at higher doses were greater than guideline recommendations (up to 10 ul/cm ²). Cumulative skin permeation was determined by measuring radioactivity in receptor fluid over a 24-hour period and in the skin after 24 hours. Carbon traps were not used to capture the evaporation of a volatile chemical. The study was conducted in a fume hood rather than under ambient conditions, although the authors indicated fume hood was used for safety reasons. However, since most of the applied dose presumably evaporated from the skin surface, the higher airflow may not have been appropriate. The study authors often point to the higher airflow as justification for the low permeation percentages. Solubility in the receptor fluid was not tested. This is considered to be a significant flaw as PBS is not generally recommended for a lipophilic compound. The study did not wash the skin sample, which deviates from OECD TG 428 recommendations.
	Metric 15:	Consistency of outcome assessment	High	Details of the outcome assessment protocol were reported, and the outcomes were consistently assessed across groups
	Metric 16:	Sampling adequacy and sensitivity	Medium	Limited details on scintillation counting were provided. Receptor fluid was poured into scintillation vials containing 12 mL of scintillation fluid and vials were measured within 1 hour of collection. Skin samples were dissolved in Soluene and analyzed by LSC. The number of scintillation counts per sample, signal to noise ratio/background levels were not reported. It is not clear that any background measurements were taken. The number of evaluations per exposure group in the data table indicates 2-7 per donor per dose, and there were 3 donors.
Domain 6: Confounding/Variable Control				
	Metric 17:	Confounding variables in test design and procedures	Low	The skin batch/lot numbers were not reported. The number of skin samples per group varied (2-7 samples per donor, 3 donors/ group). Skin integrity was measured based on 3H ₂ O permeation. It was specified that samples with water permeation > 1.2 uL/cm ² were discarded, therefore there were no differences in the quality of the tissue. The results were not expressed as Kp values or % of the applied dose. Electrical resistance and TEWL measurements were not conducted. It is unknown whether the skin sample collection sites (back, abdomen, or shoulder) were consistent within or across groups. The distribution of the samples in each group was not specified, and this missing information is likely to have a significant impact on the study results.

Continued on next page ...

... continued from previous page

Study Citation: Gajjar, R. M., Kasting, G. B. (2014). Absorption of ethanol, acetone, benzene and 1,1-Dichloroethane through human skin in vitro: a test of diffusion model predictions. *Toxicology and Applied Pharmacology* 281(1):109-117.

HERO ID: 4141541

EVALUATION				
Domain	Metric	Rating	Comments	
	Metric 18:	Confounding variables in outcomes unrelated to exposure	Low	Solubility of the test substance in the receptor fluid is a potential concern. The test substance has low solubility in water (8.6 g/L) and is lipophilic. The receptor fluid (PBS) is generally appropriate for water soluble compounds but may not be appropriate for lipophilic compounds. The solubility of the test substance in the receptor fluid was not reported, and it is unclear if the permeation rate was influenced by solubility, and whether this had a confounding impact, particularly at the higher doses.
Domain 7: Data Presentation and Analysis				
	Metric 19:	Data analysis	Low	The cumulative amount of permeated test substance was reported across time the data were reported as means \pm SE. The percent of dose permeated was graphically displayed and reported numerically in the text. Data were first averaged by dose for each donor, and then averaged across donors to obtain the mean values. It is unknown whether the study accounted for any outliers. The study did not report the percentage absorbed, but this could be determined by dividing the cumulative amount permeated by the applied dose, although it should be taken into consideration that it is presumed that the majority of the applied dose evaporated. Based on the data provided for the cumulative amount permeated (at 24hrs), the standard deviations can first be determined ($SD = SE \times \sqrt{n}$, where $n = 3$). The CV values can then be calculated (SD relative to the mean). The calculated CV values are 35%, 44%, 27%, and 23%, for the 7.9, 15.8, 31.5, and 63.1 mg/cm ² groups, respectively. CV values >25% are considered to be unacceptable. Only the high dose group had an acceptable CV (this form).
	Metric 20:	Data interpretation	Medium	The percent recovery of the applied test substance was not explicitly reported, although it is presumed to be low because the total cumulative penetration amount ranged from 0.21% to 0.13% of the applied dose. The study authors presumed the remainder evaporated; however, carbon traps were not used to definitively account for evaporated material. The skin was not washed (no measurements of washes), and no measurements of amounts in the donor chamber were collected. Only percentage absorption in the skin from a finite dose is reported and is not defined by dose group (quantitative values not reported). It is not specified whether the absorption estimates were normalized, but it is highly likely that the missing material evaporated and therefore, was unlikely to have been absorbed (under the conditions of this test); the inclusion of evaporated material may not be required. The study also reported the maximum flux of the VOC into the receptor solution. The study authors believed that the data implied that mild lipid barrier disruption likely occurred.
	Metric 21:	Reporting of data	Low	Overall cumulative absorption for each exposure group, at each collection point, was reported as Means \pm SE, and the percentage of the applied dose was noted in the text for some, but not all exposure groups. These values presumably represent primarily the amounts measured in receptor fluid. The amount in the skin after 24 hours was simply reported to be less than 0.08% (presumably of the applied dose), and was not described by exposure group. It is not clear if this amount was included in the reported cumulative values. Separate measurements for receptor fluid alone were not reported.

Overall Quality Determination

Low

Study Citation: Schenk, L., Rauma, M., Fransson, M. N., Johanson, G. (2018). Percutaneous absorption of thirty-eight organic solvents in vitro using pig skin. PLoS ONE 13(10):e0205458.
HERO ID: 4940676

EXTRACTION

Parameter	Data
Extraction ID; Chemical:	Not Reported; 1,1-Dichloroethane-Parent compound
Skin Material/Species; Skin Preparation; Skin Thickness (um); Diffusion Cell Exposure Setup Type:	ex vivo pig; Full thickness; Not Reported; Static; Notes: Not Reported
Occlusion Type; Donor Chamber Vehicle; Concentration of Test Substance in Vehicle (enter as percent):	Occluded; Deionized water; 0.8
Mass per Surface Area on Skin (mg/cm ²); Duration of Test Substance on Skin:	Not Reported; Other; 4-9 hours
Duration of Absorbance Measured; Frequency of Samples:	Other; Every 10 min for the first hour, every 20 min for the second hour, and every 30 min thereafter; Notes: 4 to 9 hours
Time Skin was Washed and Method used; Radiolabel Presence:	Not Reported; No
Total Recovery (percent); Dose Type:	Not Reported; Infinite
Percent Found in Skin Depot After Washing and Tape Stripping; Comments:	Not Reported; Notes: Not Reported
Percent Found in All Tape Strips, Excluding the Upper Two Strips; Comments:	Not Reported; Notes: Not Reported
Percent Found in Receptor Fluid and Receptor Fluid Rinse; Comments:	Not Reported; Notes: Not Reported
Total Percent Absorbed:	0
Steady State Permeability Coefficient (Kp) (cm/hr); Steady State Permeability Coefficient (Comments); Steady State Flux (ug/cm ² /hr); Steady State Flux (Comments); Maximum Permeability Coefficient (Kp) (cm/hr); Maximum Permeability Coefficient (Comments); Maximum Flux (ug/cm ² /hr); Maximum Flux (Comments):	0.0594; Notes: Coefficient of variation = 24%. The standard error of the mean was 0.00594 cm/hr.; Not Reported; Notes: Not Reported; 596; Notes: The steady state flux was reported as 0.000596 g/cm ² /hr. The standard error of the mean was 0.0000595 g/cm ² /hr = 59.5 ug/cm ² /hr.; Not Reported; Notes: Not Reported

EVALUATION

Domain	Metric	Rating	Comments
Domain 1: Test Substance			
	Metric 1: Test substance identity	High	The test substance was definitively identified by name and CASRN.
	Metric 2: Test substance source	High	The source of the test substance was reported as a manufacturer (Fluka).
	Metric 3: Test substance purity	High	The purity of the test substance was >= 99.5%.

Domain 2: Test Design

Continued on next page ...

... continued from previous page

Study Citation:		Schenk, L., Rauma, M., Fransson, M. N., Johanson, G. (2018). Percutaneous absorption of thirty-eight organic solvents in vitro using pig skin. PLoS ONE 13(10):e0205458.		
HERO ID:		4940676		
Domain	Metric	EVALUATION Rating	Comments	
	Metric 4:	Reference compounds	Medium	The study did not report use of a reference compound (or cite historical data for a reference compound like caffeine, benzoic acid, or testosterone); however, the study evaluated the dermal absorption of 36 compounds of different lipophilicity (including other compounds with similar lipophilicity as 1,2-DCA). The log P indicated for 1,2-DCA was 1.83 (cited to EPI Suite); note: the log P identified for 1,2-DCA in the TSCA Scoping Document was 1.48. The study also noted that the results that they obtained for some compounds were similar to results for the same compounds in other studies.
	Metric 5:	Assay procedures	Medium	Assay procedures specified in the study report included: diffusion cell set-up (static), temperature, identity of and rationale for using PBS as the receptor fluid, receptor volume (5.0-5.4 mL), volume of the test substance applied (i.e., excess chemical; dosing volume was specified as approximately 1 mL), and skin exposure area (0.64 cm ²). The donor compartment was capped with a glass stopper (presumably to prevent evaporation); aliquots of receptor fluid were sampled regularly (every 10 min for the first hour, every 20 min in the second hour, and then every 30 minutes). Levels in the receptor fluid aliquots were determined via gas chromatographic analysis and a standard curve. The limits of detection were reported. Any omissions (i.e., humidity only) are unlikely to substantially impact the study results.
	Metric 6:	Standards for tests	Low	The study indicated that the cut-off resistance for skin integrity was 50 ohms; that cut-off value was validated against in-house measurements of intact and damaged skin. The actual resistance values were not reported. The coefficient of variation (CV) for neat 1,2-DCA was reported (32%); no cut-off for acceptability was specified. The CV was 11% to 120% across the chemicals tested in the study (and below 75% in 29/36 cases); this was interpreted by the study authors as "low variability." Further discussion of CV is provided in Metric 19. Few QC criteria were provided to define the boundaries of the test.
Domain 3: Exposure Characterization				
	Metric 7:	Preparation and storage of test substance (chemical)	Low	Details regarding test substance preparation and storage were not provided.
	Metric 8:	Consistency of exposure administration	Medium	The study authors suggested that consistency of exposure administration within and across chemicals was one of the strengths of their study. The same area of skin (same area of pig) was used for all experiments. The volume of chemical (1 mL neat) used in the test was reported. The study authors noted when dermatomed skin, rather than full thickness skin, was used. Although some samples used in the study were split skin whereas others were full thickness skin, both experiments that involved 1,2-DCA (this experiment and the one described on a separate evaluation form) used full thickness skin (i.e., this was consistent within the 1,2-DCA experiments).
	Metric 9:	Reporting of concentrations	Medium	The test chemical concentrations were reported without ambiguity (i.e., neat 1,2-DCA); nominal concentrations were used to determine the 'apparent' K _p .

Continued on next page ...

... continued from previous page

Study Citation:		Schenk, L., Rauma, M., Fransson, M. N., Johanson, G. (2018). Percutaneous absorption of thirty-eight organic solvents in vitro using pig skin. PLoS ONE 13(10):e0205458.		
HERO ID:		4940676		
Domain	Metric	EVALUATION Rating	Comments	
	Metric 10: Exposure frequency	Low	The study did not explicitly indicate the exposure duration for each chemical (only a range of 4 to 9 hours for experiments conducted using 36 solvents, including 1,2-DCA). OECD TG 28 indicated that longer exposure periods may be required for infinite dosing protocols; standard curves were established for each chemical, and these data were used to measure the apparent permeability coefficient (Kp). The test chemical is expected to be volatile. There was no definitive evidence based on the study details provided that the exposure duration was consistent across the two 1,2-DCA experiments (i.e., the experiment for neat 1,2-DCA evaluated on this form and the experiment for diluted 1,2-DCA evaluated on a separate form).	
	Metric 11: Number of exposure groups and concentration spacing	Low	Fewer than 3 concentrations were tested; the study evaluated permeability of the neat chemical (this evaluation form) and a 0.8% dilution of 1,2-DCA in water (on a separate evaluation form).	
Domain 4: Test Model				
	Metric 12: Test model (skin)	Low	The test model and descriptive information were reported (i.e., skin from the back and the flank regions of newborn Duroc piglets that died of natural causes). Details regarding tissue storage and preparation were provided. A rationale for using this specific skin type was provided based on characteristics that make this model a suitable comparison to human skin (i.e., "pig skin is similar to human skin with respect to stratum corneum and epidermal thickness, as well as permeability" and similar flux values were observed using human and pig skin in other studies). Although human or animal skin can be used; viable skin is preferable for this study type. The study authors provided some indication that the skin samples used, from dead pigs, were viable (a requirement for skin resistance >50 ohms). As per applicable guidelines, full thickness may be used so long as excessive thickness is avoided; the thickness of the skin sample used for this experiment (and the experiment using diluted 1,2-DCA) was not explicitly specified.	
	Metric 13: Number/Replicates per group	Medium	The number of replicates per group (n = 6) was reported and appropriate.	
Domain 5: Outcome Assessment				
	Metric 14: Outcome assessment methodology	High	An infinite dose of the test substance was used to determine the Kp. It was indicated that about 1 mL was used for 0.64 cm ² of skin; this exceeds the requirement of at least 100 uL/cm ² for infinite dosing protocols as per OECD 28. The measurement technique and timing of measurements appeared appropriate.	
	Metric 15: Consistency of outcome assessment	High	Based on the information provided, it appeared that outcomes were assessed consistently across groups. The same receptor fluid was used, the same sampling times were evaluated, etc.	
	Metric 16: Sampling adequacy and sensitivity	Medium	The number of total samples per replicate was not explicitly specified; the study stated that (for all chemicals evaluated in the study), aliquots were collected every 10 min for the first hour, every 20 min for the second hours, and every 30 minutes thereafter, and that experiments ranged from 4-9 hours. However, based on the timing of sampling (and knowledge about the standard curve for 1,2-DCA), sampling appeared appropriate.	
Domain 6: Confounding/Variable Control				
Continued on next page ...				

... continued from previous page

Study Citation:		Schenk, L., Rauma, M., Fransson, M. N., Johanson, G. (2018). Percutaneous absorption of thirty-eight organic solvents in vitro using pig skin. PLoS ONE 13(10):e0205458.		
HERO ID:		4940676		
Domain	Metric	EVALUATION Rating	Comments	
	Metric 17: Confounding variables in test design and procedures	Medium	No differences were reported with respect to the size or integrity of skin samples among groups (but it was not explicitly specified that there were no differences). Skin integrity was measured using a preferable method (i.e., electrical resistance) and the study criteria were met for all replicates (i.e., at least 50 ohms for each skin sample). The study did not indicate how the skin samples correlated to chemical testing (number of pigs from which skin samples were obtained to perform the experiments).	
	Metric 18: Confounding variables in outcomes unrelated to exposure	High	The study indicated that the maximum solubility of the receptor fluid was not exceeded for 1,2-DCA (or any of the chemicals tested). There was no evidence to suggest that the test material interfered with the assay conditions.	
Domain 7: Data Presentation and Analysis	Metric 19: Data analysis	Uninformative	Calculation of the Kp was described adequately in the methods. Kp measurements were based on the linear/steady-state of the curve; pre- and post-steady-state values were excluded from analyses. [Correlations between Kp values and other factors (molecular weight, number of carbons) across chemicals were analyzed statistically.] However, the CV using neat 1,2-DCA was 32%, which is outside the range for acceptability (according to OECD 156, a CV <25% indicates that variability is relatively low, while a CV equal or greater than 25% indicates that variability is high). The correlation of experimental Kp values with predicted values (using EPI Suite) was generally poor for both neat (separate evaluation) and diluted (this evaluation) substances evaluated in this study (not just 1,2-DCA).	
	Metric 20: Data interpretation	High	An infinite dosing protocol was used to calculate permeability. Results were interpreted adequately based on the assay set-up/conditions.	
	Metric 21: Reporting of data	High	Data were reported by exposure group for all measured outcomes. Table 1 included data for the test substance (neat = this evaluation or diluted = evaluated on a separate form, molecular weight, log P), the skin used (full thickness for 1,2-DCA experiments), number of replicates (n = 6) as well values determined experimentally, including lag time, flux at steady state, and coefficients of variation.	

Overall Quality Determination

Uninformative

Study Citation: Schenk, L., Rauma, M., Fransson, M. N., Johanson, G. (2018). Percutaneous absorption of thirty-eight organic solvents in vitro using pig skin. PLoS ONE 13(10):e0205458.
HERO ID: 4940676

EXTRACTION**Parameter****Data**

Extraction ID; Chemical: Not Reported; 1,1-Dichloroethane-Parent compound
 Skin Material/Species; Skin Preparation; Skin Thickness (um); Diffusion Cell Exposure Setup Type: ex vivo pig; Full thickness; Not Reported; Static; Notes: Not Reported
 Occlusion Type; Donor Chamber Vehicle; Concentration of Test Substance in Vehicle (enter as percent): Occluded; None; 100
 Mass per Surface Area on Skin (mg/cm²); Duration of Test Substance on Skin: Not Reported; Other; 4 to 9 hours
 Duration of Absorbance Measured; Frequency of Samples: Other; Receptor fluid swabs sampled every 10 min for the first hour, every 20 min for the second hour, and every 30 min thereafter; Notes: 4 to 9 hours
 Time Skin was Washed and Method used; Radiolabel Presence: Not Reported; No
 Total Recovery (percent); Dose Type: Not Reported; Infinite
 Percent Found in Skin Depot After Washing and Tape Stripping; Comments: Not Reported; Notes: Not Reported
 Percent Found in All Tape Strips, Excluding the Upper Two Strips; Comments: Not Reported; Notes: Not Reported
 Percent Found in Receptor Fluid and Receptor Fluid Rinse; Comments: Not Reported; Notes: Not Reported
 Total Percent Absorbed: 0
 Steady State Permeability Coefficient (Kp) (cm/hr); Steady State Permeability Coefficient (Comments); Steady State Flux (ug/cm²/hr); Steady State Flux (Comments); Maximum Permeability Coefficient (Kp) (cm/hr); Maximum Permeability Coefficient (Comments); Maximum Flux (ug/cm²/hr); Maximum Flux (Comments): 0.00109; Notes: Coefficient of variation = 32%. The standard error of the mean was 0.000143 cm/hr.; Not Reported; Notes: Not Reported; 1360; Notes: The steady state flux was reported as 0.00136 g/cm²/hr. The standard error of the mean was 0.000179 g/cm²/hr = 179 ug/cm²/hr.; Not Reported; Notes: Not Reported

EVALUATION

Domain	Metric	Rating	Comments
Domain 1: Test Substance			
	Metric 1: Test substance identity	High	The test substance was definitively identified by name and CASRN.
	Metric 2: Test substance source	High	The source of the test substance was reported as a manufacturer (Fluka).
	Metric 3: Test substance purity	High	The purity of the test substance was $\geq 99.5\%$.
Domain 2: Test Design			

Continued on next page ...

... continued from previous page

Study Citation:		Schenk, L., Rauma, M., Fransson, M. N., Johanson, G. (2018). Percutaneous absorption of thirty-eight organic solvents in vitro using pig skin. PLoS ONE 13(10):e0205458.		
HERO ID:		4940676		
Domain	Metric	EVALUATION Rating	Comments	
	Metric 4:	Reference compounds	Medium	The study did not report use of a reference compound (or cite historical data for a reference compound like caffeine, benzoic acid, or testosterone); however, the study evaluated the dermal absorption of 36 compounds of different lipophilicity (and including other compounds with similar lipophilicity as 1,2-DCA). The log P indicated for 1,2-DCA was 1.83 (cited to EPI Suite); note: the log P identified for 1,2-DCA in the TSCA Scoping Document was 1.48. The study also noted that the results that they obtained for some compounds were similar to results for the same compounds in other studies.
	Metric 5:	Assay procedures	Medium	Assay procedures specified in the study report included: diffusion cell set-up (static), temperature, identity of and rationale for using PBS as the receptor fluid, receptor volume (5.0-5.4 mL), volume of the test substance applied (i.e., excess chemical; dosing volume was specified as approximately 1 mL), and skin exposure area (0.64 cm ²). The donor compartment was capped with a glass stopper (presumably to prevent evaporation); aliquots of receptor fluid were sampled regularly (every 10 min for the first hour, every 20 min in the second hour, and then every 30 minutes). Levels in the receptor fluid aliquots were determined via gas chromatographic analysis and a standard curve. The limits of detection were reported. Any omissions (i.e., humidity only) are unlikely to substantially impact the study results.
	Metric 6:	Standards for tests	Low	The study indicated that the cut-off resistance for skin integrity was 50 ohms; that cut-off value was validated against in-house measurements of intact and damaged skin. The actual resistance values were not reported. The coefficient of variation (CV) for diluted 1,2-DCA was reported (24%); no cut-off for acceptability was specified. The CV was 11% to 120% across the chemicals tested in the study (and below 75% in 29/36 cases); this was interpreted by the study authors as "low variability." Further discussion of CV is provided in Metric 19. Few QC criteria were provided to define the boundaries of the test.
Domain 3: Exposure Characterization				
	Metric 7:	Preparation and storage of test substance (chemical)	Low	Details regarding test substance preparation and storage were not provided.
	Metric 8:	Consistency of exposure administration	Medium	The study authors suggested that consistency of exposure administration within and across chemicals was one of the strengths of their study. The same area of skin (same area of pig) was used for all experiments. The volume of chemical (1 mL of 0.8% 1,2-DCA) used in the test was reported. The study authors noted when dermatomed skin, rather than full thickness skin, was used. Although some samples used in the study were split skin whereas others were full thickness skin, both experiments that involved 1,2-DCA (this experiment and the one described on a separate evaluation form) used full thickness skin (i.e., this was consistent within the 1,2-DCA experiments). In addition, although most of the chemicals evaluated in the study were diluted in water, a few (that were not miscible in water) were diluted in PBS. This would not affect findings for 1,2-DCA because only the neat chemical (evaluated in a separate evaluation) or the diluted chemical (0.8% 1,2-DCA in water, this evaluation) were used; there was no inconsistency with respect to the 1,2-DCA experiments.
	Metric 9:	Reporting of concentrations	Medium	The test chemical concentrations were reported without ambiguity (i.e., 0.8% 1,2-DCA diluted in water); nominal concentrations were used to determine the 'apparent' K _p .

Continued on next page ...

... continued from previous page

Study Citation:		Schenk, L., Rauma, M., Fransson, M. N., Johanson, G. (2018). Percutaneous absorption of thirty-eight organic solvents in vitro using pig skin. PLoS ONE 13(10):e0205458.		
HERO ID:		4940676		
		EVALUATION		
Domain	Metric	Rating	Comments	
	Metric 10:	Exposure frequency	Low	The study did not explicitly indicate the exposure duration for each chemical (only a range of 4 to 9 hours for experiments conducted using 36 solvents, including 1,2-DCA). OECD TG 28 indicated that longer exposure periods may be required for infinite dosing protocols; standard curves were established for each chemical, and these data were used to measure the apparent permeability coefficient (Kp). The test chemical is expected to be volatile. There was no definitive evidence based on the study details provided that the exposure duration was consistent across the two 1,2-DCA experiments (i.e., the experiment for neat 1,2-DCA evaluated on a separate form and the experiment for diluted 1,2-DCA evaluated on this form).
	Metric 11:	Number of exposure groups and concentration spacing	Low	Fewer than 3 concentrations were tested; the study evaluated permeability of the neat chemical (evaluated on a separate form) and a 0.8% dilution of 1,2-DCA in water (evaluated on this form).
Domain 4: Test Model				
	Metric 12:	Test model (skin)	Low	The test model and descriptive information were reported (i.e., skin from the back and the flank regions of newborn Duroc piglets that died of natural causes). Details regarding tissue storage and preparation were provided. A rationale for using this specific skin type was provided based on characteristics that make this model a suitable comparison to human skin (i.e., "pig skin is similar to human skin with respect to stratum corneum and epidermal thickness, as well as permeability" and similar flux values were observed using human and pig skin in other studies). Although human or animal skin can be used; viable skin is preferable for this study type. The study authors provided some indication that the skin samples used, from dead pigs, were viable (a requirement for skin resistance >50 ohms). As per applicable guidelines, full thickness may be used so long as excessive thickness is avoided; the thickness of the skin sample used for this experiment (and the experiment using neat 1,2-DCA) was not explicitly specified.
	Metric 13:	Number/Replicates per group	Medium	The number of replicates per group (n = 6) was reported and appropriate.
Domain 5: Outcome Assessment				
	Metric 14:	Outcome assessment methodology	High	An infinite dose of the test substance was used to determine the Kp. It was indicated that about 1 mL was used for 0.64 cm ² of skin; this exceeds the requirement of at least 100 uL/cm ² for infinite dosing protocols as per OECD 28. The measurement technique and timing of measurements appeared appropriate.
	Metric 15:	Consistency of outcome assessment	High	Based on the information provided, it appeared that outcomes were assessed consistently across groups. The same receptor fluid was used, the same sampling times were evaluated, etc.).
	Metric 16:	Sampling adequacy and sensitivity	Medium	The number of total samples per replicate was not explicitly specified; the study stated that (for all chemicals evaluated in the study), aliquots were collected every 10 min for the first hour, every 20 min for the second hours, and every 30 minutes thereafter, and that experiments ranged from 4-9 hours. However, based on the timing of sampling (and knowledge about the standard curve for 1,2-DCA), sampling appeared appropriate.
Domain 6: Confounding/Variable Control				

Continued on next page ...

... continued from previous page

Study Citation:	Schenk, L., Rauma, M., Fransson, M. N., Johanson, G. (2018). Percutaneous absorption of thirty-eight organic solvents in vitro using pig skin. PLoS ONE 13(10):e0205458.			
HERO ID:	4940676			
	EVALUATION			
Domain	Metric	Rating	Comments	
	Metric 17:	Confounding variables in test design and procedures	Medium	No differences were reported with respect to the size or integrity of skin samples among groups (but it was not explicitly specified that there were no differences). Skin integrity was measured using a preferable method (i.e., electrical resistance) and the study criteria were met for all replicates (i.e., at least 50 ohms for each skin sample). The study did not indicate how the skin samples correlated to chemical testing (number of pigs from which skin samples were obtained to perform the experiments).
	Metric 18:	Confounding variables in outcomes unrelated to exposure	High	The study indicated that the maximum solubility of the receptor fluid was not exceeded for 1,2-DCA (or any of the chemicals tested). There was no evidence to suggest that the test material interfered with the assay conditions.
Domain 7: Data Presentation and Analysis				
	Metric 19:	Data analysis	High	Calculation of the Kp was described adequately in the methods. Kp measurements were based on the linear/steady-state of the curve; pre- and post-steady-state values were excluded from analyses. [Correlations between Kp values and other factors (molecular weight, number of carbons) across chemicals were analyzed statistically.] The CV using diluted 1,2-DCA was 24%, which is within the range for acceptability (according to OECD 156, a CV of <25% indicates that variability is relatively low, while a CV equal or greater than 25% indicates that variability is high). The correlation of experimental Kp values with predicted values (using EPI Suite) was generally poor for both neat (separate evaluation) and diluted (this evaluation) substances evaluated in this study (not just 1,2-DCA).
	Metric 20:	Data interpretation	High	An infinite dosing protocol was used to calculate permeability. Results were interpreted adequately based on the assay set-up/conditions.
	Metric 21:	Reporting of data	High	Data were reported by exposure group for all measured outcomes. Table 1 included data for the test substance (diluted = this evaluation or neat = evaluated on a separate form, molecular weight, log P), the skin used (full thickness for 1,2-DCA experiments), number of replicates (n = 6) as well values determined experimentally, including lag time, flux at steady state, and coefficients of variation.

Overall Quality Determination

Medium

Study Citation: Jakobson, I., Wahlberg, J. E., Holmberg, B., Johansson, G. (1982). Uptake via the blood and elimination of 10 organic solvents following epicutaneous exposure of anesthetized guinea pigs. *Toxicology and Applied Pharmacology* 63(2):181-187.
HERO ID: 94881

EXTRACTION

Parameter	Data
Extraction ID; Chemical:	1,2-DCE - uptake; 1,1-Dichloroethane-Parent compound
Species; Comments:	Guinea pig; Not Reported; Notes: Not Reported
Sex; Covering used in Test System:	Both; Occluded
Vehicle; Concentration of Test Substance in Vehicle (percent):	None specified; Not Reported
Mass per Surface Area on Skin (mg/cm ²); Dose (include units (e.g., mg/kg bw)):	403; 1 ml/3.1 cm ² - no additional information was provided.
Test Substance on Skin; Exposure Repeated (Days); Test Substance on Skin (Comments):	Other; The exposure was continuous for 12 hours; Notes: 12 hours
Time of Absorption Measured; Frequency; Time of Absorption Measured (Comments):	Other; Every 5th and 12th minute.; Notes: 12 hours
Time Skin was Washed and Method used; Radiolabel Presence:	Skin was not washed; No
Total Recovery (percent); Dose Type:	Not Reported; Infinite
Percent Found in Skin Depot after Washing and Tape Stripping:	Not Reported; Notes: The concentration on skin was not measured.
Percent Found in Urine ; Comments:	Not Reported; Notes: The concentration on skin was not measured.
Percent Found in Feces ; Comments:	Not Reported; Notes: The concentration on skin was not measured.
Percent Found in Blood/Serum ; Comments:	Not Reported; Notes: The concentration on skin was not measured.
Percent Found in Air ; Comments:	Not Reported; Notes: The concentration in expired air was not measured
Percent Found in Cage Wash ; Comments:	Not Reported; Notes: The concentration in expired air was not measured
Percent Found in All Tape Strips, Excluding the Upper Two Strips:	Not Reported; Notes: The concentration on skin was not measured.
Total Percent Absorbed:	0
Steady State Permeability Coefficient (Kp) (cm/hr); Steady State Permeability Coefficient (Comments); Steady State Flux (ug/cm ² /hr); Steady State Flux (Comments); Maximum Permeability Coefficient (Kp) (cm/hr); Maximum Permeability Coefficient (Comments); Maximum Flux (ug/cm ² /hr); Maximum Flux (Comments); Additional Comments:	Not Reported; Notes: The concentration in expired air was not measured; Not Reported; Notes: The concentration in expired air was not measured; Not Reported; Notes: The concentration in expired air was not measured; Notes: Not Reported

EVALUATION

Domain	Metric	Rating	Comments
Domain 1: Test Substance	Metric 1: Test substance identity	High	The test substance was identified definitively as 1, 2-dichloroethane. The form (liquid or solid) of the test substance was not specified. It was not radiolabeled.

Continued on next page ...

... continued from previous page

Study Citation:		Jakobson, I., Wahlberg, J. E., Holmberg, B., Johansson, G. (1982). Uptake via the blood and elimination of 10 organic solvents following epicutaneous exposure of anesthetized guinea pigs. Toxicology and Applied Pharmacology 63(2):181-187.		
HERO ID:		94881		
		EVALUATION		
Domain	Metric	Rating	Comments	
	Metric 2:	Test substance source	Low	The test material was sourced from Merck, A.G., Darmstadt, West Germany. A batch or lot number was not specified. Certification was not provided. The exact material purchased from the supplier could not be determined (multiple products listed), therefore, the chemical could not be verified on the supplier's website.
	Metric 3:	Test substance purity	Medium	The test material was reported to be spectroscopic grade, an actual purity was not reported, but chemicals of this grade are generally highly pure.
Domain 2: Test Design				
	Metric 4:	Randomized allocation of animals	Low	The study did not specify whether animals were randomized.
	Metric 5:	Standards for Tests	Uninformative	The study authors did not report whether the test met any pre-established criteria. This study did not employ the use of metabolism cages. Percutaneous uptake over time in an infinite exposure model was evaluated by measuring concentrations of the test substance in the blood. The study did not calculate or determine percent recovery. Graphs, presumably showing mean blood concentrations over time did not include any measures of variance. Measures of variance also were not provided in a table reporting quantitative values. This lack of data prevents the ability to determine the variance across means making it impossible to determine the validity of the test.
Domain 3: Exposure Characterization				
	Metric 6:	Preparation and storage of test substance (chemical)	Low	Details of the test substance preparation and storage were not provided. No dilutions or use of vehicles were mentioned, so it is assumed that the materials were applied neat. Storage conditions of the purchased material were not reported, and the test material is a volatile chemical.
	Metric 7:	Consistency of exposure administration	High	Details of exposure administration were reported and based on the information provided, exposures were administered consistently across study groups. Glass rings (area 3.1 cm ²) were glued onto the clipped skin on the backs of guinea pigs. A cover glass with a central hole was attached, and 1 mL of the test material was injected. The holes were covered to prevent leakage or evaporation.
	Metric 8:	Reporting of concentrations	Medium	The study reported the volume applied (1mL per depot) and the skin surface area (3.1 cm ²); equivalent to 0.322 mL/cm ² . Using the density of the test material (1.25 g/mL), a dose in mg/cm ² can be determined. (1.25 g/mL) x 0.322 mL/cm ² = 0.403 g/cm ² (or 403 mg/cm ²). There is no indication that the purchased test material was analytically verified, but because the material was applied neat, and had spectral grade purity, the lack of measurement is not expected to have a significant impact on the study results.
	Metric 9:	Exposure duration	High	Based on Figure 3 in the results, the exposure duration was 12 hours. The study authors did not provide justification for the duration of an uptake experiment, However, the purpose of the study was simply to measure concentrations in blood during exposure, and the authors did note (based on unpublished data) that "the concentration might eventually become fatal if the exposure is extended."

Continued on next page ...

... continued from previous page

Study Citation:		Jakobson, I., Wahlberg, J. E., Holmberg, B., Johansson, G. (1982). Uptake via the blood and elimination of 10 organic solvents following epicutaneous exposure of anesthetized guinea pigs. <i>Toxicology and Applied Pharmacology</i> 63(2):181-187.		
HERO ID:		94881		
		EVALUATION		
Domain	Metric	Rating	Comments	
	Metric 10:	Number of exposure groups and concentration spacing	Low	The study included two groups. One group of animals had a single skin depot with 1 mL of the test substance; the other group had two skin depots, each containing 1 mL of the test substance. Separate groups of animals were used for elimination experiments (separate form). OECD TG 427 only specifies that a study will "normally involve several groups;" therefore, the number of groups in this study seems to be appropriate. It is unclear whether the concentrations/dosing was appropriate. Exposure should mimic potential human exposure, which is typically up to 10 uL/cm ² for a liquid. This study used 1 mL/3.1 cm ² , or 322.5 uL/cm ² which is significantly higher than recommended in OECD 427.
Domain 4: Test Model	Metric 11:	Test animal characteristics	Low	The study used male and female guinea pigs (distribution within groups not specified) with starting body weights ranging between 600 and 1,000 g (a mean weight was not reported), and it is not known whether the weights of the animals within a group were within 20% of the mean weight. The animal strain, age, and source were not reported. Guinea pigs are an appropriate model as per OECD 427, although rats are more commonly used.
	Metric 12:	Adequacy and consistency of animal husbandry conditions	Low	No animal husbandry details were provided.
	Metric 13:	Number of animals per group	Low	The study used a total of 4 animals per group. The methods indicated that both males and females were used, but the number of each sex/group was not specified. OECD TG 427 indicates that at least 4 animals of a single-sex should be used.
Domain 5: Outcome Assessment	Metric 14:	Outcome assessment methodology	Uninformative	The outcome assessment methodology deviated significantly from OECD 427 guidelines. Based on the information provided (volume applied and surface area), a dose of 700 mg/cm ² was determined using a test substance density of 2.17 g/mL, indicating infinite dosing. The surface area (3.1 cm ²) is lower than the guideline recommendations and does not cover 10% of the animal surface area. The study did not evaluate anything other than the presence of the chemical in blood. It did not evaluate 1,2-DCA remaining in the excreta, carcass, skin, skin wash, or exhaled air. Therefore it cannot be used to determine skin absorption.
	Metric 15:	Consistency of outcome assessment	Medium	The outcome assessment methodology was reported, and based on the available information was mostly consistent across groups. There was one animal that was not anaesthetized; however, this animal showed a similar uptake curve, and the difference is not expected to have a significant impact on results.
	Metric 16:	Sampling adequacy and sensitivity	Medium	None of the figures or data tables reported an "n." It is assumed that the data were generated from all 4 animals per group, but this was not explicitly stated. It is unclear if the frequency of blood collection is appropriate because it necessitated replacement with Marcodex.
Domain 6: Confounding/Variable Control				

Continued on next page ...

... continued from previous page

Study Citation:		Jakobson, I., Wahlberg, J. E., Holmberg, B., Johansson, G. (1982). Uptake via the blood and elimination of 10 organic solvents following epicutaneous exposure of anesthetized guinea pigs. Toxicology and Applied Pharmacology 63(2):181-187.		
HERO ID:		94881		
Domain	Metric	EVALUATION Rating	Comments	
	Metric 17:	Confounding variables in test design and procedures	Medium	The study did not report all information to determine whether confounding bias occurred (test substance batch, body weight changes, or body weight variations greater than 20%, compared to the mean). Animals were anaesthetized throughout the experiment. Based on the results with one animal that was not anaesthetized, it was determined that the injection of pentobarbital (anaesthesia) did not significantly change the concentration of the test material in the blood.
	Metric 18:	Confounding variables in outcomes unrelated to exposure	Medium	There was no information either to support or dismiss the suggestion that there were differences among groups in animal attrition, health outcomes unrelated to exposure, or solubility that could influence the outcome assessment.
Domain 7: Data Presentation and Analysis				
	Metric 19:	Data analysis	Uninformative	This study did not determine absorption estimates. The only results were blood concentrations over time and these data were not statistically analyzed.
	Metric 20:	Data interpretation	Uninformative	The study described the observed absorption pattern for the test material in blood but is inadequate for quantitatively determining skin absorption. Concentrations in the blood increased throughout the entire exposure, except for a short dip at 1 hr. In animals with a single skin depot, the maximum concentration in blood (at 12 hours) was approximately 20 ug/mL. This study did not calculate the amount of test the chemical absorbed per cm ² of skin over time. It is unclear how useful this study is due to significant deviations from standard guidelines.
	Metric 21:	Reporting of Data	Uninformative	The data for some, but not all specified outcomes were presented. The plot of blood concentrations by time for the group of animals exposed to 2 skin depots was not provided. Only concentrations at 0.5 and 6 hours as provided in table format. None of the data provided in the study included measures of variance. Data were inadequate for evaluating the fraction of the chemical absorbed through skin.

Overall Quality Determination**Uninformative**

Study Citation: Jakobson, I., Wahlberg, J. E., Holmberg, B., Johansson, G. (1982). Uptake via the blood and elimination of 10 organic solvents following epicutaneous exposure of anesthetized guinea pigs. *Toxicology and Applied Pharmacology* 63(2):181-187.

HERO ID: 94881

EXTRACTION

Parameter	Data
Extraction ID; Chemical:	1,2-DCE - Elimination; 1,1-Dichloroethane-Parent compound
Species; Comments:	Guinea pig; Not Reported; Notes: Not Reported
Sex; Covering used in Test System:	Both; Occluded
Vehicle; Concentration of Test Substance in Vehicle (percent):	None specified; Not Reported
Mass per Surface Area on Skin (mg/cm ²); Dose (include units (e.g., mg/kg bw)):	403; 1 ml/3.1 cm ² - no additional information was provided.
Test Substance on Skin; Exposure Repeated (Days); Test Substance on Skin (Comments):	Other; The exposure was continuous for 4 hours.; Notes: 4 hours
Time of Absorption Measured; Frequency; Time of Absorption Measured (Comments):	6 hrs; The methods state every 5th and 12th minute, but it is unclear if this was the frequency for all experiments.; Notes: Not Reported
Time Skin was Washed and Method used; Radiolabel Presence:	Skin was not washed; No
Total Recovery (percent); Dose Type:	Not Reported; Infinite
Percent Found in Skin Depot after Washing and Tape Stripping:	Not Reported; Notes: The concentration in the skin depot was not measured.
Percent Found in Urine ; Comments:	Not Reported; Notes: The concentration in the skin depot was not measured.
Percent Found in Feces ; Comments:	Not Reported; Notes: The concentration in the skin depot was not measured.
Percent Found in Blood/Serum ; Comments:	Not Reported; Notes: The concentration in the skin depot was not measured.
Percent Found in Air ; Comments:	Not Reported; Notes: The concentration in expired air was not measured
Percent Found in Cage Wash ; Comments:	Not Reported; Notes: The concentration in expired air was not measured
Percent Found in All Tape Strips, Excluding the Upper Two Strips:	Not Reported; Notes: The concentration in the skin depot was not measured.
Total Percent Absorbed:	0
Steady State Permeability Coefficient (Kp) (cm/hr); Steady State Permeability Coefficient (Comments); Steady State Flux (ug/cm ² /hr); Steady State Flux (Comments); Maximum Permeability Coefficient (Kp) (cm/hr); Maximum Permeability Coefficient (Comments); Maximum Flux (ug/cm ² /hr); Maximum Flux (Comments); Additional Comments:	Not Reported; Notes: The concentration in expired air was not measured; Not Reported; Notes: The concentration in expired air was not measured; Not Reported; Notes: The concentration in expired air was not measured; Not Reported; Notes: The concentration in expired air was not measured; Notes: Not Reported

EVALUATION

Domain	Metric	Rating	Comments	
Domain 1: Test Substance				
	Metric 1:	Test substance identity	High	The test substance was identified definitively as 1, 2-dichloroethane. The form (liquid or solid) of the test substance was not specified. It was not radiolabeled.
	Metric 2:	Test substance source	Low	The test material was sourced from Merck, A.G., Darmstadt, West Germany. A batch or lot number was not specified. Certification was not provided. The exact material purchased from the supplier could not be determined (multiple products listed), therefore, the chemical could not be verified on the supplier's website.

Continued on next page ...

... continued from previous page

Study Citation:		Jakobson, I., Wahlberg, J. E., Holmberg, B., Johansson, G. (1982). Uptake via the blood and elimination of 10 organic solvents following epicutaneous exposure of anesthetized guinea pigs. Toxicology and Applied Pharmacology 63(2):181-187.		
HERO ID:		94881		
Domain	Metric	EVALUATION Rating	Comments	
	Metric 3:	Test substance purity	Medium	The test material was reported to be spectroscopic grade, an actual purity was not reported, but chemicals of this grade are generally highly pure.
Domain 2: Test Design	Metric 4:	Randomized allocation of animals	Low	The study did not specify whether animals were randomized.
	Metric 5:	Standards for Tests	Uninformative	The study authors did not report whether the test met any pre-established criteria. This study did not employ the use of metabolism cages. Percutaneous uptake over time in an infinite exposure model was evaluated by measuring concentrations of the test substance in the blood. The study did not calculate or determine percent recovery. Graphs, presumably showing mean blood concentrations over time did not include any measures of variance. Measures of variance also were not provided in a table reporting quantitative values. This lack of data prevents the ability to determine the variance across means making it impossible to determine the validity of the test.
Domain 3: Exposure Characterization	Metric 6:	Preparation and storage of test substance (chemical)	Low	Details of the test substance preparation and storage were not provided. No dilutions or use of vehicles were mentioned, so it is assumed that the materials were applied neat. Storage conditions of the purchased material were not reported, and the test material is a volatile chemical.
	Metric 7:	Consistency of exposure administration	High	Details of exposure administration were reported and based on the information provided, exposures were administered consistently across study groups. Glass rings (area 3.1 cm ²) were glued onto the clipped skin on the backs of guinea pigs. A cover glass with a central hole was attached, and 1 mL of the test material was injected. During elimination experiments, the seal was broken and the excess solvent was aspirated and the skin was dried with compressed air.
	Metric 8:	Reporting of concentrations	Medium	The study reported the volume applied (1mL per depot) and the skin surface area (3.1 cm ²); equivalent to 0.322 mL/cm ² . Using the density of the test material (1.25 g/mL), a dose in mg/cm ² can be determined. (1.25 g/mL) x 0.322 mL/cm ² = 0.403 g/cm ² (or 403 mg/cm ²). There is no indication that the purchased test material was analytically verified, but because the material was applied neat, and had spectral grade purity, the lack of measurement is not expected to have a significant impact on the study results.
	Metric 9:	Exposure duration	High	For the elimination part of the study, animals were percutaneously exposed for 4 hours. Blood samples were then taken during the following 2 hours. The duration was not justified but seemed to be appropriate for this part of the test.
	Metric 10:	Number of exposure groups and concentration spacing	Low	Elimination experiments (this form) were only done with animals exposed to one skin depot. OECD TG 427 only specifies that a study will "normally involve several groups;" therefore, the number of groups in this study seems to be appropriate. It is unclear whether the concentrations/dosing was appropriate. Exposure should mimic potential human exposure, which is typically up to 10 uL/cm ² for a liquid. This study used 1 mL/3.1 cm ² , or 322.5 uL/cm ² which is significantly higher than recommended in OECD 427.
Domain 4: Test Model				

Continued on next page ...

... continued from previous page

Study Citation:		Jakobson, I., Wahlberg, J. E., Holmberg, B., Johansson, G. (1982). Uptake via the blood and elimination of 10 organic solvents following epicutaneous exposure of anesthetized guinea pigs. Toxicology and Applied Pharmacology 63(2):181-187.		
HERO ID:		94881		
		EVALUATION		
Domain	Metric	Rating	Comments	
	Metric 11: Test animal characteristics	Low	The study used male and female guinea pigs (distribution within groups not specified) with starting body weights ranging between 600 and 1,000 g (a mean weight was not reported), and it is not known whether the weights of the animals within a group were within 20% of the mean weight. The animal strain, age, and source were not reported. Guinea pigs are an appropriate model as per OECD 427, although rats are more commonly used. .	
	Metric 12: Adequacy and consistency of animal husbandry conditions	Low	No animal husbandry details were provided.	
	Metric 13: Number of animals per group	Low	The number of animals per group was not specified in the methods. A table suggests 4 animals per group, but this table is reporting results from the uptake experiment. It is unclear if the same number of animals were used for the elimination experiments. The methods indicated that both males and females were used, but the number of each sex/group was not specified. OECD TG 427 indicates that at least 4 animals of a single-sex should be used.	
Domain 5: Outcome Assessment				
	Metric 14: Outcome assessment methodology	Uninformative	The outcome assessment methodology deviated significantly from OECD 427 guidelines. Based on the information provided (volume applied and surface area), a dose of 403 mg/cm ² was determined using a test substance density of 1.25 g/mL, indicating infinite dosing. The surface area (3.1 cm ²) is lower than the guideline recommendations and does not cover 10% of the animal surface area. The study did not evaluate anything other than the presence of the chemical in blood. It did not evaluate the chemical remaining in the excreta, carcass, skin, skin wash, or exhaled air. Therefore it cannot be used to determine skin absorption.	
	Metric 15: Consistency of outcome assessment	High	The outcome assessment methodology was reported, and, based on the available information was consistent across groups.	
	Metric 16: Sampling adequacy and sensitivity	Medium	None of the figures or data tables reported an "n." It is assumed that the data were generated from all 4 animals per group, but this was not explicitly stated. It is unclear if the frequency of blood collection is appropriate because it necessitated replacement with Marcodex.	
Domain 6: Confounding/Variable Control				
	Metric 17: Confounding variables in test design and procedures	Medium	The study did not report all information to determine whether confounding bias occurred (test substance batch, body weight changes, or body weight variations greater than 20%, compared to the mean). Animals were anaesthetized throughout the experiment.	
	Metric 18: Confounding variables in outcomes unrelated to exposure	Medium	There was no information either to support or dismiss the suggestion that there were differences among groups in animal attrition, health outcomes unrelated to exposure, or solubility that could influence the outcome assessment.	
Domain 7: Data Presentation and Analysis				

Continued on next page ...

... continued from previous page

Study Citation: Jakobson, I., Wahlberg, J. E., Holmberg, B., Johansson, G. (1982). Uptake via the blood and elimination of 10 organic solvents following epicutaneous exposure of anesthetized guinea pigs. *Toxicology and Applied Pharmacology* 63(2):181-187.

HERO ID: 94881

EVALUATION

Domain	Metric	EVALUATION Rating	Comments
	Metric 19: Data analysis	Uninformative	Data analysis methods were described. Nonlinear regression analysis was conducted on values obtained for the concentration of the test material in the blood. The study referenced Holmberg et al. 1977 for additional details (open access, reviewed for this assessment). This cited reference did not provide a significant amount of additional details. The data were fitted to an equation representing a simplified two-compartment kinetic model (Snyder et al. 1981). The study, however, cannot be used to measure absorption based on a lack of appropriate information.
	Metric 20: Data interpretation	Uninformative	The study specified studying elimination, but only measured concentrations in blood for 2 hours post-exposure, and not measure concentrations in other excreta. The study also did not calculate clearance from plasma, or report any elimination kinetics. This part of the study does not provide any useful information on absorption.
	Metric 21: Reporting of Data	Uninformative	A figure shows elimination curves from blood drawn according to a two-compartment model. The "n" was not reported and the data points had no measures of variance. Data were inadequate for evaluating the fraction of the chemical absorbed through skin.

Overall Quality Determination**Uninformative**

Study Citation: Morgan, D. L., Cooper, S. W., Carlock, D. L., Sykora, J. J., Sutton, B., Mattie, D. R., Mcdougal, J. N. (1991). Dermal absorption of neat and aqueous volatile organic chemicals in the Fischer 344 rat. Environmental Research 55(1):51-63.

HERO ID: 200487

EXTRACTION

Parameter	Data
Extraction ID; Chemical:	1-2-Dichloroethane - NEAT; 1,1-Dichloroethane-Parent compound
Species; Comments:	Rat; Not Reported; Notes: Not Reported
Sex; Covering used in Test System:	Male; Occluded
Vehicle; Concentration of Test Substance in Vehicle (percent):	No vehicle was used.; Not Reported
Mass per Surface Area on Skin (mg/cm ²); Dose (include units (e.g., mg/kg bw)):	1,399; Neat, concentration not reported, but is assumed to be taken at its density (2.17 g/mL)
Test Substance on Skin; Exposure Repeated (Days); Test Substance on Skin (Comments):	24 hrs; Not Reported; Notes: Not Reported
Time of Absorption Measured; Frequency; Time of Absorption Measured (Comments):	24 hrs; Blood samples were measured at 0, 0.5, 1, 2, 4, 8, and 24 hours; Notes: Not Reported
Time Skin was Washed and Method used; Radiolabel Presence:	Skin was not washed; No
Total Recovery (percent); Dose Type:	Not Reported; Infinite
Percent Found in Skin Depot after Washing and Tape Stripping:	Not Reported; Notes: The skin depot was analyzed
Percent Found in Urine ; Comments:	Not Reported; Notes: The skin depot was analyzed
Percent Found in Feces ; Comments:	Not Reported; Notes: The skin depot was analyzed
Percent Found in Blood/Serum ; Comments:	Not Reported; Notes: The skin depot was analyzed
Percent Found in Air ; Comments:	Not Reported; Notes: Concentrations in expelled air were not measured.
Percent Found in Cage Wash ; Comments:	Not Reported; Notes: Concentrations in expelled air were not measured.
Percent Found in All Tape Strips, Excluding the Upper Two Strips:	Not Reported; Notes: The skin depot was analyzed
Total Percent Absorbed:	0
Steady State Permeability Coefficient (Kp) (cm/hr); Steady State Permeability Coefficient (Comments); Steady State Flux (ug/cm ² /hr); Steady State Flux (Comments); Maximum Permeability Coefficient (Kp) (cm/hr); Maximum Permeability Coefficient (Comments); Maximum Flux (ug/cm ² /hr); Maximum Flux (Comments); Additional Comments:	Not Reported; Notes: Concentrations in expelled air were not measured.; Not Reported; Notes: Concentrations in expelled air were not measured.; Not Reported; Notes: Concentrations in expelled air were not measured.; Notes: Not Reported

EVALUATION

Domain	Metric	Rating	Comments
Domain 1: Test Substance	Metric 1: Test substance identity	High	The test substance was definitively identified as 1,1-Dichloroethane, it was not radiolabeled.

Continued on next page ...

... continued from previous page

Study Citation:		Morgan, D. L., Cooper, S. W., Carlock, D. L., Sykora, J. J., Sutton, B., Mattie, D. R., Mcdougal, J. N. (1991). Dermal absorption of neat and aqueous volatile organic chemicals in the Fischer 344 rat. Environmental Research 55(1):51-63.		
HERO ID:		200487		
		EVALUATION		
Domain	Metric	Rating	Comments	
	Metric 2:	Test substance source	High	The test substance was sourced from Aldrich Chemical Co. The exact product purchased from the supplier was not specified, but the study did perform GC analysis to confirm purity, and all of the products sold by Aldrich are certified.
	Metric 3:	Test substance purity	High	The purity was >99% and was confirmed by gas chromatography before study initiation. The text indicated that the test substance came stabilized with 0.05% low alkyl epoxides.
Domain 2: Test Design				
	Metric 4:	Randomized allocation of animals	Medium	The method of animal allocation was not specified, although it was indicated that some normalization to body weight occurred (body weights of rats within the same dose group were all within a 10 g range).
	Metric 5:	Standards for Tests	Low	The study authors did not report whether the test met any pre-established criteria. This study did not employ the use of metabolism cages. Percutaneous uptake over time in an intended infinite exposure model was evaluated by measuring concentrations of the test substance in the blood. The study did not calculate or determine percent recovery. It was mentioned that there was no loss to head space in the exposure cells, or to evaporation. The study did not explicitly report coefficients of variance. Figures showing mean blood concentrations over time did include SEM, but SD, and thus CVs cannot be determined because the "n" was provided as a range (n = 6-10). The only endpoint in which CVs can be determined was the volume of the chemical solution absorbed. These CV values (calculated for this review and noted in Metric 19) were > 25%.
Domain 3: Exposure Characterization				
	Metric 6:	Preparation and storage of test substance (chemical)	High	No preparation of the neat chemical was required. Saturated aqueous solutions were prepared by mixing 10-20 mL of neat chemical with 200 mL of HPLC-grade water. Details of mixing and removal of any remaining immiscible chemicals were provided. The samples were stored at 5-10 degrees C in zero head-space vials for 24 -48 hours prior to use. The study text suggests that stability tests were performed, and the solutions were stable for at least three weeks. The saturated solutions were diluted with HPLC-grade water.
	Metric 7:	Consistency of exposure administration	High	All animals were consistently exposed using the same protocol. Briefly, glass exposure cells (20 mm diameter, skin area 3.1 cm ²) were attached to the shaved backs of rats the day before exposure. 2 mL volumes of the test solution were added to each cell under occlusive conditions (2 mL/3.1 cm ² or 0.645 mL/cm ²), and animals were exposed for 24 hours.
	Metric 8:	Reporting of concentrations	High	Animals were exposed to neat, 1/3 saturated (2,270 ug/mL), 2/3 saturated (4,290 ug/mL), or saturated (6,738 ug/mL) aqueous solutions. Saturated solutions were analyzed by GC to ensure that the concentrations were in the range of literature values for saturated aqueous solutions. The concentration of the neat test material was not specified, but it is assumed it can be taken at its density (2.17 g/mL).
	Metric 9:	Exposure duration	High	Animals were exposed for 24 hours, which is consistent with OECD 427 guidelines.
	Metric 10:	Number of exposure groups and concentration spacing	High	The study included 4 exposure groups. The concentration spacing was not explicitly justified, but the study wanted to determine if there were differences in the volume absorbed and in blood concentrations attained from different dilutions.

Continued on next page ...

...continued from previous page

Study Citation:		Morgan, D. L., Cooper, S. W., Carlock, D. L., Sykora, J. J., Sutton, B., Mattie, D. R., Mcdougal, J. N. (1991). Dermal absorption of neat and aqueous volatile organic chemicals in the Fischer 344 rat. Environmental Research 55(1):51-63.		
HERO ID:		200487		
Domain	Metric	EVALUATION Rating	Comments	
Domain 4: Test Model				
	Metric 11:	Test animal characteristics	Medium	The study used male Fisher 344 rats sourced from Charles River Breeding Laboratories. An initial body weight range (at purchase) was reported to be 201-215g. Animals were acclimated for a week. Body weights prior to exposure were reported to be between 215 and 300 g, with all rats within one dose group within a 10g range. The variation between groups is not clear. Animal age was not reported.
	Metric 12:	Adequacy and consistency of animal husbandry conditions	Medium	Housing, animals per cage, and food and water availability were reported. Additional animal husbandry details (e.g., temperature, humidity) were not specified.
	Metric 13:	Number of animals per group	Low	The number of animals per group was not specified in the methods. Based on the data tables, the sample size ranged from 6 to 10. It is unclear whether all groups initially had 10 animals per group, or if the number per group was a range from 6-10 animals. OECD guidelines only require 4 animals per group, so overall, the number was more than required.
Domain 5: Outcome Assessment				
	Metric 14:	Outcome assessment methodology	Low	The outcome assessment methodology was sufficient to address the intentions of this non-guideline study, but deviated from a traditional dermal absorption study as per OECD 427. In brief, this study attempted infinite exposure under occlusion. The mass per area was > 10 mg/cm ² for all exposure groups; however, exposure was only about 1% of the total rat skin surface area, which is less than the suggested 5-10% (OECD 28). Blood samples were collected during exposure at times 0, 0.5, 1, 2, 4, 8, and 24 hours. The collection vials were sealed and stored at 5 degrees until GC analysis. They were stable for at least 3 weeks. At the end of exposure, the volume of test solution remaining in the exposure cell was measured. The remaining volume was subtracted from the initial 2 mL used to obtain the volume absorbed. This study did not determine the % recovery or the percent absorbed, but these were not intended endpoints for this study. The study did not conduct any skin washes or a wash of the dermal cell. Concentrations in the skin, other bodily fluids, or in exhaled air were not analyzed.
	Metric 15:	Consistency of outcome assessment	Low	Blood sample volumes were reported to be 50 or 100 uL, it is unclear if different volumes were collected from animals within a group, or across groups for a single chemical. This inconsistency could have an impact on the study results. Other aspects of outcome assessment (e.g., the timing of blood collection and study termination) were consistent across groups.
	Metric 16:	Sampling adequacy and sensitivity	Low	The sampling sizes were not specified in the methods but were provided in the data tables and figures. The figures reported sample sizes "n" as a range from 6-10. For the volume of aqueous chemical absorbed endpoint, the data table indicated exact sample sizes. There is some uncertainty and concern over the differing sample sizes. The number of animals per group was not reported. It is not clear whether the sample sizes represent different numbers of animals per group, or if the differences, instead, whether only a portion of the full number of animals dosed were sampled, which could lead to some biases. In some instances, the volume remaining in the neat samples was low, and the authors suspected leakage. These samples were not included in the calculations of volume absorbed, but this does not account for the different sample sizes in other groups.
Domain 6: Confounding/Variable Control				

Continued on next page ...

... continued from previous page

Study Citation:		Morgan, D. L., Cooper, S. W., Carlock, D. L., Sykora, J. J., Sutton, B., Mattie, D. R., Mcdougal, J. N. (1991). Dermal absorption of neat and aqueous volatile organic chemicals in the Fischer 344 rat. Environmental Research 55(1):51-63.		
HERO ID:		200487		
Domain	Metric	EVALUATION Rating	Comments	
	Metric 17:	Confounding variables in test design and procedures	Medium	Animal body weights at the start of exposure remained within 10 g within a group. Overall, weights ranged from 215-300 g. A mean was not specified, so it is not known if these ranges fall within 20% of the mean. It was noted that some exposure cells with low volumes of neat test chemicals were assumed to have leaked. The study tested multiple chemicals and did not specify what chemicals or how many samples were affected by this. It was noted that these cells were not used in calculating the mean absorption volume. The text also indicated that there was a small amount of headspace in the exposure cells; however, the authors said the headspace was too small to lower the exposure concentration. They confirmed this by exposing one group of rats to saturated 1,1,1-trichloroethane using a cell completely filled (no headspace). After 24 hours, the concentration of 1,1,1-trichloroethane was still depleted to an amount similar to samples that had a headspace, indicating no loss to headspace occurred.
	Metric 18:	Confounding variables in outcomes unrelated to exposure	Medium	There was no information either to support or dismiss the suggestion that there were differences among groups in animal attrition, health outcomes unrelated to exposure, or solubility that could influence the outcome assessment.
Domain 7: Data Presentation and Analysis				
	Metric 19:	Data analysis	Uninformative	The coefficient of variation could only be determined for the volume of chemical solution absorbed. The study authors noted a significant amount of variation, and the CV (calculated for this review) was 26%, 76%, 33%, and 31% for the neat, saturated, 2/3 saturated, and 1/3 saturated solutions, respectively, making this study critically deficient.
	Metric 20:	Data interpretation	Uninformative	This study was not conducted according to OECD 427, or in a manner allowing the determination of recovery or the calculation of absorption estimates. The only tissue compartment analyzed was blood/plasma. Absorption volumes were also determined, but not the total percent absorbed. The authors found no correlation between blood levels and absorption volumes. Although the authors intended to create an infinite exposure scenario, less than 1% of the initial concentration was purportedly present in the exposure cells after 24 hours (chemical-specific quantitative values were not provided). This indicated rapid absorption and depletion from the aqueous solutions. The authors believed evaporation of the chemical was unlikely because the exposure cells were sealed, and the Teflon-coated caps were found not to absorb the test chemicals. The usefulness of this study is limited to demonstrating absorption through the skin occurs upon exposure to lower dilutions in aqueous solutions.
	Metric 21:	Reporting of Data	Low	Data were reported for all of the outcomes specified in the methods and included concentrations in blood over time and volume of test solution absorbed for each group. Data were presented as means ± either SE or SD. The number of samples was provided either as a range, or a specific numerical value. Because a range was reported for some data, CV values could not be calculated for those endpoints, which significantly reduces the ability to interpret the data.

Overall Quality Determination**Uninformative**